



HAL
open science

Spin contamination and noncollinearity in general complex Hartree-Fock wave functions

Patrick Cassam-Chenaï

► **To cite this version:**

Patrick Cassam-Chenaï. Spin contamination and noncollinearity in general complex Hartree-Fock wave functions. 2013. hal-00790674v4

HAL Id: hal-00790674

<https://hal.univ-cotedazur.fr/hal-00790674v4>

Preprint submitted on 9 Aug 2015 (v4), last revised 4 Sep 2015 (v5)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spin contamination and noncollinearity in general complex Hartree-Fock wave functions

Patrick Cassam-Chenai

August 9, 2015

Abstract An expression for the square of the spin operator expectation value, $\langle S^2 \rangle$, is obtained for a general complex Hartree-Fock (GCHF) wave function and decomposed into four contributions: The main one whose expression is formally identical to the restricted (open-shell) Hartree-Fock expression. A spin contamination one formally analogous to that found for spin unrestricted Hartree-Fock wave functions. A noncollinearity contribution related to the fact that the wave function is not an eigenfunction of the spin- S_z operator. A perpendicularity contribution related to the fact that the spin density is not constrained to be zero in the xy-plane. All these contributions are evaluated and compared for the H_2O^+ system. The optimization of the collinearity axis is also considered.

Keywords spin contamination · collinearity · general complex Hartree-Fock

Patrick Cassam-Chenai

Univ. Nice Sophia Antipolis, CNRS, LJAD, UMR 7351, 06100 Nice, France

Tel.: +33-4-92076260

Fax: +33-4-93517974

E-mail: cassam@unice.fr

1 Introduction

Particle-independent models based on single Slater determinant wave functions, have enjoyed considerable interest in quantum chemistry, since the pioneering works of Hartree, Slater and Fock [1–3].

When a quantum system is described by a spin-free Hamiltonian, which obviously commutes with the spin operators S_z and S^2 , a spin-symmetry respectful way of using the Hartree-Fock method consists in:

- 1) Using spin-orbitals of pure α - or β -spin, so that the HF optimized Slater determinant is an eigenfunction of S_z ;
- 2) Imposing the spin-equivalence restriction [4], which means that paired α - and β -spin-orbitals are formed from the same set of linearly independent orbitals. We have proved mathematically [5–7] that this additional constraint is a necessary and sufficient condition to insure that a Slater determinantal wave function is an eigenfunction of the spin operator S^2 . In other words, we have shown that relaxing the S^2 -symmetry constraint exactly amounts to allow different “paired orbitals”, in the sense of Refs. [8,9], to have different spins. This equivalence enabled us to characterize the variational space explored by the restricted open-shell Hartree-Fock (ROHF) method [10], which precisely consists in optimizing a Slater determinant subject to constraints 1) and 2) (plus spatial-symmetry constraints if any) [11]. The equivalence was also discovered independently [12] by optimizing a Slater determinant with a Lagrange multiplier, enforcing $\langle S^2 \rangle$ to be arbitrarily close to the ROHF value, instead of applying the spin-equivalence restriction. Not surprisingly, the determinant was approaching the ROHF solution.

Similar to the spin-free case is the “complex-free” one: When a quantum system is described by a real Hamiltonian, which obviously commutes with complex conjugation, one can restrict oneself to the calculation of real eigenfunctions. Then, it is also possible to employ only real spin-orbitals to construct the HF Slater determinant [13]. (However, difficulties may occur when the symmetry group of the molecule cannot be represented over real numbers, and nonetheless, one wishes the spin-orbitals to be adapted to spatial-symmetry).

However, it has been proposed by various authors to relax some or all of the above-mentioned constraints, to gain variational freedom. For example, the different orbitals for different spins method (DODS) of Refs. [14,15], (which is usually just called “unrestricted Hartree-Fock” (UHF), but in this paper we use “DODS” to avoid confusions), relaxes the spin-equivalence restriction, hence the HF

solution is no longer an eigenfunction of S^2 . Other authors [16–18] have advocated the use of general spin-orbitals, mixing α -spin and β -spin parts, in conjunction with the use of projectors [19].

Along the same line of thought, the use of complex spin-orbitals has been proposed [17, 20] to increase variational freedom in the case of real Hamiltonian. Prat and Lefebvre went a step further with so-called “hypercomplex” spin-orbitals to construct Slater determinants of arbitrary accuracy [21]. However, the coefficients of their spin-orbitals were elements of a Clifford algebra of dimension 2^{2n} , that was not a normed division algebra, also known as Cayley algebra, for arbitrary values of n . This was unfortunate, since such a structure appears to be a minimal requirement for a quantum formalism, if, for example, Born’s interpretation of the wave function is to hold firmly. For $n = 1$, the Clifford algebra of Prat et al. was actually the non-commutative field of Quaternions, therefore, *a fortiori*, a normed division algebra. The only larger normed division algebra is the Octonion algebra. It is a Clifford algebra of dimension 8, which has also been proposed in a quantum mechanical context [22], but this algebra is neither commutative nor associative. The lack of these properties rises difficulties for its use for multipartite quantum systems, nevertheless these difficulties can be overcome by keeping the product of octonion coefficients in the form of a tensor product. So, octonion-unrestricted HF appears to be the largest Clifford algebra-unrestricted single determinantal method that can be considered in the spirit of Prat and Lefebvre’s proposal. However, octonions seem incompatible with the desirable requirement that the algebra of quantum observables be what is now called a formally real Jordan algebra [23] acting on a vector space of arbitrarily large dimension. Octonions are also ruled out by the requirement of orthomodularity in infinite dimension according to Solèr’s theorem [24, 25], which restricts quantum Hilbert spaces to be real, complex or at most quaternionic.

The first HF molecular calculations with general complex spin-orbitals, without projecting out the symmetry-breaking part of the wave function, are maybe those of Ref. [26]. It was found on the BH molecule around its equilibrium geometry that the general complex Hartree-Fock (GCHF) energy was indeed lower than the DODS one, which itself was lower than the restricted Hartree-Fock (RHF) solution. So necessarily, the corresponding GCHF wave functions had S^2 -spin contamination and S_z -spin contamination, that is to say, the expectation values of these operators were different from 0, the value expected for a singlet ground state. (It is not clear whether complex numbers were used for this molecule, but the authors did mention that they performed complex calculations for 2-electron systems.)

Relaxing the “ S_z -constraint” hence the “collinearity constraint”, becomes perfectly legitimate when hyperfine or spin-orbit couplings are considered, since the operator S_z no longer commutes with the

Hamiltonian. As a matter of fact, real physical systems do exhibit either light [27,28] or strong [29, 30] noncollinearity of their spin densities. Similarly, the use of complex spin-orbitals is natural, when considering relativistic corrections resulting in a complex Hamiltonian operator. So, in such a context, one should use no less than general complex spin-orbitals in HF calculations [31]. The “spin-same-orbit” coupling term used in these calculations does not commute with the S^2 -operator. Therefore, one cannot strictly speak of “ S^2 -spin contamination” in relativistic GCHF wave functions. However, calculating the expectation value of S^2 , a *bona fide* quantum observable, can still provide valuable physical information about the system.

A general expression for the expectation value of S^2 has been obtained in the DODS case [8], and has served as a measure of S^2 -spin contamination. However, as far as we are aware, no such formula has been published in the case of a GCHF wave function. This gap will be filled in the next section.

Studying departure from collinearity is more difficult because of arbitrariness in the quantification axis. One possible way to overcome the difficulty would be to apply an external magnetic field to fix the z -axis but small enough not to perturb the GCHF solution. However, an elegant alternative has been proposed recently by Small et al. [32]. It is based on studying the lowest eigenvalue of a (3×3) -matrix built from expectation values of spin operator components and their products. In the GCHF case, the authors provided the expressions required to compute the matrix elements in a compact form. In the third section, we give a more extended formula in terms of molecular orbital overlap matrix elements. We also illustrate the connections between spin contamination, noncollinearity and its correlative: “perpendicularity” on the H_2O^+ cation example. We sum up our conclusions in the last section.

2 Spin contamination in GCHF

A General Complex Hartree Fock (GCHF) wave function

$$\Phi_{GCHF} = \phi_1 \wedge \cdots \wedge \phi_{N_e} \quad (1)$$

is the antisymmetrized product (or wedge product, denoted by \wedge) of orthonormal spinorbitals, or “two-component spinors”,

$$\phi_i = \begin{pmatrix} \phi_{i\alpha} \\ \phi_{i\beta} \end{pmatrix}, \quad (2)$$

$$\langle \phi_i | \phi_j \rangle = \delta_{i,j}, \quad (3)$$

where the scalar product $\langle \cdot | \cdot \rangle$ means integration over space variables and summation (i.e. taking the trace) over spin variables. The same symbol will also be used for the scalar product between orbitals (i.e. no trace taken). We define the “number of α -spin electrons” (respectively “number of β -spin electrons”) as $N_\alpha := \sum_{i=1}^{N_e} \langle \phi_{i\alpha} | \phi_{i\alpha} \rangle$ (respectively, $N_\beta := \sum_{i=1}^{N_e} \langle \phi_{i\beta} | \phi_{i\beta} \rangle$). Note that they need not be integer numbers, however, $N_\alpha + N_\beta = N_e$.

Let us work out the expectation value of the spin operator,

$$S^2 = S_z^2 + \frac{1}{2}(S^+ S^- + S^- S^+), \quad (4)$$

on a general GCHF wave function.

The action of S_z is given by,

$$S_z \Phi_{GCHF} = \frac{1}{2} \sum_{i=1}^{N_e} \hat{\Phi}_{GCHF}^i, \quad (5)$$

where,

$$\hat{\Phi}_{GCHF}^i = \phi_1 \wedge \cdots \wedge \phi_{i-1} \wedge \hat{\phi}_i \wedge \phi_{i+1} \wedge \cdots \wedge \phi_{N_e}, \quad (6)$$

and,

$$\hat{\phi}_i = \begin{pmatrix} +\phi_{i\alpha} \\ -\phi_{i\beta} \end{pmatrix}. \quad (7)$$

Note that,

$$\langle \hat{\phi}_i | \hat{\phi}_j \rangle = \langle \phi_i | \phi_j \rangle = \delta_{i,j}. \quad (8)$$

So, the expectation value of S_z^2 is,

$$\begin{aligned}
\langle \Phi_{GCHF} | S_z^2 | \Phi_{GCHF} \rangle &= \langle S_z \Phi_{GCHF} | S_z \Phi_{GCHF} \rangle \\
&= \frac{1}{4} \sum_{i,j=1}^{N_e} \langle \hat{\Phi}_{GCHF}^i | \hat{\Phi}_{GCHF}^j \rangle \\
&= \frac{1}{4} \left(\sum_{i=1}^{N_e} \langle \hat{\Phi}_{GCHF}^i | \hat{\Phi}_{GCHF}^i \rangle + \sum_{i \neq j} \langle \hat{\Phi}_{GCHF}^i | \hat{\Phi}_{GCHF}^j \rangle \right) \\
&= \frac{1}{4} \sum_{i=1}^{N_e} \left(\langle \hat{\phi}_i | \hat{\phi}_i \rangle + \sum_{j \neq i} (-1) |\langle \hat{\phi}_i | \phi_j \rangle|^2 + \langle \hat{\phi}_i | \phi_i \rangle \langle \phi_j | \hat{\phi}_j \rangle \right) \\
&= \frac{1}{4} \left(N_e + \sum_{i=1}^{N_e} \sum_{j \neq i} (-1) |\langle \phi_{i\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{i\beta} | \phi_{j\beta} \rangle|^2 + (\langle \phi_{i\alpha} | \phi_{i\alpha} \rangle - \langle \phi_{i\beta} | \phi_{i\beta} \rangle) (\langle \phi_{j\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{j\beta} | \phi_{j\beta} \rangle) \right) \\
&= \frac{1}{4} \left(N_e + \sum_{i,j=1}^{N_e} (\langle \phi_{i\alpha} | \phi_{i\alpha} \rangle - \langle \phi_{i\beta} | \phi_{i\beta} \rangle) (\langle \phi_{j\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{j\beta} | \phi_{j\beta} \rangle) - |\langle \phi_{i\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{i\beta} | \phi_{j\beta} \rangle|^2 \right) \\
&= \left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} \right)^2 + \frac{1}{4} \left(N_e - \sum_{i,j=1}^{N_e} |\langle \phi_{i\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{i\beta} | \phi_{j\beta} \rangle|^2 \right). \tag{9}
\end{aligned}$$

This equation reduces to $\left(\frac{N_\alpha}{2} - \frac{N_\beta}{2}\right)^2$ in the case of a DODS wave function. So, the second term on the right-hand side (rhs), which is $(\langle \Phi_{GCHF} | S_z^2 | \Phi_{GCHF} \rangle - \langle \Phi_{GCHF} | S_z | \Phi_{GCHF} \rangle^2)$ is directly related to relaxation of the S_z -constraint and will be called the “ z -noncollinearity” contribution. Note, however, that for a GCHF wave function, the first term on the rhs does not necessarily correspond to an eigenvalue of S_z^2 , according to the definition of N_α and N_β .

The action of S^+ is given by,

$$S^+ \Phi_{GCHF} = \sum_{i=1}^{N_e} \acute{\Phi}_{GCHF}^i, \tag{10}$$

where,

$$\acute{\Phi}_{GCHF}^i = \phi_1 \wedge \cdots \wedge \phi_{i-1} \wedge \acute{\phi}_i \wedge \phi_{i+1} \wedge \cdots \wedge \phi_{N_e}, \tag{11}$$

and,

$$\acute{\phi}_i = \begin{pmatrix} +\phi_{i\beta} \\ 0 \end{pmatrix}. \tag{12}$$

Similarly, the action of S^- is given by,

$$S^- \Phi_{GCHF} = \sum_{i=1}^{N_e} \grave{\Phi}_{GCHF}^i, \tag{13}$$

where,

$$\dot{\Phi}_{GCHF}^i = \phi_1 \wedge \cdots \wedge \phi_{i-1} \wedge \dot{\phi}_i \wedge \phi_{i+1} \wedge \cdots \wedge \phi_{N_e}, \quad (14)$$

and,

$$\dot{\phi}_i = \begin{pmatrix} 0 \\ +\phi_{i\alpha} \end{pmatrix}. \quad (15)$$

So, the expectation value of $S^- S^+$ is,

$$\begin{aligned} \langle \Phi_{GCHF} | S^- S^+ | \Phi_{GCHF} \rangle &= \langle S^+ \Phi_{GCHF} | S^+ \Phi_{GCHF} \rangle \\ &= \sum_{i,j=1}^{N_e} \langle \dot{\Phi}_{GCHF}^i | \dot{\Phi}_{GCHF}^j \rangle \\ &= \sum_{i=1}^{N_e} \langle \dot{\Phi}_{GCHF}^i | \dot{\Phi}_{GCHF}^i \rangle + \sum_{i \neq j} \langle \dot{\Phi}_{GCHF}^i | \dot{\Phi}_{GCHF}^j \rangle \\ &= \sum_{i=1}^{N_e} \left(\langle \dot{\phi}_i | \dot{\phi}_i \rangle + \sum_{j \neq i} (-1) |\langle \dot{\phi}_i | \dot{\phi}_j \rangle|^2 + \langle \dot{\phi}_i | \dot{\phi}_i \rangle \langle \dot{\phi}_j | \dot{\phi}_j \rangle \right) \\ &= \sum_{i=1}^{N_e} \left(\langle \phi_{i\beta} | \phi_{i\beta} \rangle + \sum_{j \neq i} (-1) |\langle \phi_{i\beta} | \phi_{j\alpha} \rangle|^2 + \langle \phi_{i\beta} | \phi_{i\alpha} \rangle \langle \phi_{j\alpha} | \phi_{j\beta} \rangle \right) \\ &= N_\beta + \sum_{i,j=1}^{N_e} \langle \phi_{i\beta} | \phi_{i\alpha} \rangle \langle \phi_{j\alpha} | \phi_{j\beta} \rangle - \langle \phi_{i\beta} | \phi_{j\alpha} \rangle \langle \phi_{j\alpha} | \phi_{i\beta} \rangle. \end{aligned} \quad (16)$$

Similarly, the expectation value of $S^+ S^-$ is,

$$\begin{aligned} \langle \Phi_{GCHF} | S^+ S^- | \Phi_{GCHF} \rangle &= \langle S^- \Phi_{GCHF} | S^- \Phi_{GCHF} \rangle \\ &= \sum_{i=1}^{N_e} \left(\langle \dot{\phi}_i | \dot{\phi}_i \rangle + \sum_{j \neq i} (-1) |\langle \dot{\phi}_i | \dot{\phi}_j \rangle|^2 + \langle \dot{\phi}_i | \dot{\phi}_i \rangle \langle \dot{\phi}_j | \dot{\phi}_j \rangle \right) \\ &= \sum_{i=1}^{N_e} \left(\langle \phi_{i\alpha} | \phi_{i\alpha} \rangle + \sum_{j \neq i} (-1) |\langle \phi_{i\alpha} | \phi_{j\beta} \rangle|^2 + \langle \phi_{i\alpha} | \phi_{i\beta} \rangle \langle \phi_{j\beta} | \phi_{j\alpha} \rangle \right) \\ &= N_\alpha + \sum_{i,j=1}^{N_e} \langle \phi_{i\alpha} | \phi_{i\beta} \rangle \langle \phi_{j\beta} | \phi_{j\alpha} \rangle - \langle \phi_{i\alpha} | \phi_{j\beta} \rangle \langle \phi_{j\beta} | \phi_{i\alpha} \rangle. \end{aligned} \quad (17)$$

Using Eq.(4) and putting together Eqs.(9), (16) and (17), one obtains the expectation value of S^2 ,

$$\begin{aligned} \langle \Phi_{GCHF} | S^2 | \Phi_{GCHF} \rangle &= \left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} \right)^2 + \frac{N_\alpha}{2} + \frac{N_\beta}{2} + \frac{1}{4} \left(N_e - \sum_{i,j=1}^{N_e} |\langle \phi_{i\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{i\beta} | \phi_{j\beta} \rangle|^2 \right) \\ &+ \sum_{i,j=1}^{N_e} \langle \phi_{i\alpha} | \phi_{i\beta} \rangle \langle \phi_{j\beta} | \phi_{j\alpha} \rangle - \langle \phi_{i\alpha} | \phi_{j\beta} \rangle \langle \phi_{j\beta} | \phi_{i\alpha} \rangle. \end{aligned} \quad (18)$$

The expression reduces to the known formula in the case of a DODS wave function. Assuming, without loss of generality, that $N_\alpha \geq N_\beta$, we rewrite Eq. (18) as,

$$\begin{aligned} \langle \Phi_{GCHF} | S^2 | \Phi_{GCHF} \rangle &= \left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} \right) \left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} + 1 \right) + \frac{1}{4} \left(N_e - \sum_{i,j=1}^{N_e} |\langle \phi_{i\alpha} | \phi_{j\alpha} \rangle - \langle \phi_{i\beta} | \phi_{j\beta} \rangle|^2 \right) \\ &+ \left(N_\beta - \sum_{i,j=1}^{N_e} \langle \phi_{i\alpha} | \phi_{j\beta} \rangle \langle \phi_{j\beta} | \phi_{i\alpha} \rangle \right) + \left| \sum_{i=1}^{N_e} \langle \phi_{i\beta} | \phi_{i\alpha} \rangle \right|^2. \end{aligned} \quad (19)$$

In this formula we identify four contributions: The first term is formally identical to the ROHF expression also found in the DODS case. However, care must be taken that it is actually different, because the numbers of α - and β -electrons are not good quantum numbers in the GCHF case. The second term on the first line is the “ z -noncollinearity” contribution. The third term on the second line is formally analogous to the “spin contamination” of a DODS wave function as defined in [7,8]. Finally, the last term on the second line arises from the release of the S_z -constraint, which allows for the two components of a given, general spin-orbital to be non zero. But it originates from S^+S^- and S^-S^+ , and is maximal when $\phi_{i\beta} = \exp(i\theta)\phi_{i\alpha}$ for all i , that is to say when the ϕ_i ’s are eigenfunctions of $\cos\theta S_x + \sin\theta S_y$ for some angle θ . It is related to the emergence of a non-zero spin density in the x, y -plane, correlatively to the loss of z -collinearity. We tentatively call this term the “ x, y -perpendicularity” contribution.

The present formulas have been implemented in the code TONTO [33] and applied in a recent article (third column of Tab. 4 in [34]). Let us discuss further the different contributions to S^2 for a H_2O^+ GCHF calculation similar to that reported in [34]. The z -quantification axis was the axis perpendicular to the plane of the molecule. The results, see Tab. 1, shows that the main contribution to $\langle \Phi_{GCHF} | S^2 | \Phi_{GCHF} \rangle$ beside the reference expression (first term in eq.(19)) is the so-called spin contamination contribution (we set $\hbar = 1$ throughout the paper). The x, y -perpendicularity and z -noncollinearity contributions are of the same order of magnitude and more than one order of magnitude smaller. Added to $\left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} \right) \left(\frac{N_\alpha}{2} - \frac{N_\beta}{2} + 1 \right)$, they almost make up the reference value of +0.75. So, the spin contamination value of 0.007033 amounts almost exactly to the difference between the exact expectation value $\langle \Phi_{GCHF} | S^2 | \Phi_{GCHF} \rangle$ and this reference “ROHF value”.

3 Collinearity in GCHF

In the previous section, we have encountered a z -(non)collinearity measure, $col_z := (\langle S_z^2 \rangle - \langle S_z \rangle^2)$. This quantity can be generalized to an arbitrary quantization direction defined by a unit vector $\mathbf{u} = \begin{pmatrix} u_x \\ u_y \\ u_z \end{pmatrix}$ of the unit sphere \mathcal{S}^2 of \mathbb{R}^3 by replacing S_z by $\mathbf{u} \cdot \mathbf{S} = \sum_{\mu \in \{x,y,z\}} u_\mu S_\mu$. Then, \mathbf{u} -(non)collinearity is measured by:

$$col(\mathbf{u}) := \sum_{\mu, \nu \in \{x,y,z\}} u_\mu u_\nu (\langle S_\mu S_\nu \rangle - \langle S_\mu \rangle \langle S_\nu \rangle). \quad (20)$$

Small et al. [32] defined a (non)collinearity measure by:

$$col := \min_{\mathbf{u} \in \mathcal{S}^2} col(\mathbf{u}), \quad (21)$$

which corresponds to the lowest eigenvalue of the matrix A whose elements are given by,

$$A_{\mu\nu} = \Re(\langle S_\mu S_\nu \rangle) - \langle S_\mu \rangle \langle S_\nu \rangle, \quad (22)$$

where $\Re(z)$ is the real part of z . The associated eigenvector gives the optimal collinearity direction. Setting $x\tilde{\phi} = \frac{1}{2}(\phi + \dot{\phi})$, $y\tilde{\phi} = \frac{-i}{2}(\phi - \dot{\phi})$ and $z\tilde{\phi} = \hat{\phi}$, we have in these notation,

$$\forall \mu, \nu \in \{x, y, z\} \quad A_{\mu\nu} = \delta_{\mu\nu} \frac{N_e}{4} - \sum_{i,j=1}^{N_e} \langle \tilde{\phi}_i | \phi_j \rangle \langle \phi_j | \tilde{\phi}_i \rangle, \quad (23)$$

where $\delta_{\mu\nu}$ is the Krönecker symbol.

Returning to the H_2O^+ example and applying these formulae, we obtain

$$A = \begin{pmatrix} +0.253128 + 0.000145 - 0.009774 \\ +0.000145 + 0.253451 + 0.003745 \\ -0.009774 + 0.003745 + 0.000461 \end{pmatrix} \quad (24)$$

The diagonalization of the A -matrix gives the optimal collinear direction: $\mathbf{u}_0^t = (+0.0385908, -0.014789, +0.999146)$, which is only slightly tilted with respect to the z -direction, and the system is quasi-collinear in this direction since $col = 0.000028$ is very close to zero. This shows that the noncollinearity contribution to $\langle \Phi_{GCHF} | S^2 | \Phi_{GCHF} \rangle$ could be further reduced by more than one order of magnitude by selecting the optimal quantization axis corresponding to \mathbf{u}_0 instead of the spatial z -axis. The perpendicularity contribution would decrease accordingly.

4 Conclusion

We have decomposed the expectation value of the spin operator S^2 into (i) a term formally identical to its expression for a ROHF reference wave function, (ii) a term called “spin contamination” because it is formally analogous to that derived by Amos and Hall [8] for DODS wave functions, (iii) a noncollinear contribution which can be minimized by following a procedure recently introduced [32], (iv) a term called the “perpendicularity contribution” which arises from the release of the z -collinearity constraint but which should rather be regarded as arising from the release of the “nonperpendicularity constraint” on the spin-density. The collinearity and nonperpendicularity constraints are correlatives.

We have evaluated these four different contributions for a GCHF calculation on the H_2O^+ cation. Note that we used the IOTC relativistic Hamiltonian [36] so that the term “spin contamination” is not really appropriate in this context, departure from the ROHF reference value being legitimate. However, the so-called spin contamination contribution has been found to dominate the noncollinearity and perpendicularity ones. This could be made even more so, by tilting the quantification axis to the optimal collinearity direction.

Acknowledgements

We acknowledge Dr. Lukáš Bučinský for drawing our attention to the problem of the derivation of GCHF spin contamination, and to the fact that S^2 does not commute with the “spin-same-orbit” coupling term, usually used in quantum chemistry. The referees and the editor are acknowledged for suggesting many improvements to the manuscript. This article is a tribute to Prof. P. Surjàn and is also dedicated to the memory of the late Prof. Gaston Berthier, who introduced to the author in the course of vivid discussions, many of the references listed below.

References

1. D. R. Hartree, Proc. Cambridge Phil. Soc. 24 (1928) 89.
2. J. C. Slater, Phys. Rev. 34 (1929) 1293.
3. V. Fock, Z. Physik 61 (1930) 126.
4. G. Berthier, in *Molecular orbitals in Chemistry, Physics and Biology*, (P. O. Löwdin, B. Pullman ed., Acad. Press New York, 1964), p.57.
5. P. Cassam-Chenaï, *Algèbre fermionique et chimie quantique*, Ph.D thesis, Université de Paris 6, (1992).

6. P. Cassam-Chenaï, G.S. Chandler, *Sur les fonctions de Hartree-Fock sans contrainte*, Comptes-rendus de l'acadmie des sciences srie II, **314**, p.755-757, (1992).
7. P. Cassam-Chenaï et G.S. Chandler, *Int. J. Quantum Chem.* 46 (1993) 593-607.
8. A. T. Amos, G. G. Hall, *Proc. Roy. Soc. A*263 (1961) 483.
9. P. O. Löwdin, *J. Appl. Phys. Suppl.* 33 (1962) 251.
10. P. Cassam-Chenaï, *J. Math. Chem.* 15 (1994) 303.
11. C. C. J. Roothaan, *Rev. Mod. Phys.* 32 (1960) 179.
12. J. S. Andrews, D. Jayatilaka, R. G. A. Bone, N. C. Handy, R. D. Amos, *Chem. Phys. Lett.* 183 (1991) 423.
13. E. Brändas, *J. Mol. Spectrosc.* 27 (1968) 236.
14. G. Berthier, *Comptes-Rendus de l'Acadmie des Sciences* 238 (1954) 91-93.
15. J. A. Pople, R. K. Nesbet, *J. Chem. Phys* 22 (1954) 571.
16. C. Bunge, *Phys. Rev.* 154 (1967) 70.
17. R. Lefebvre, Y. G. Smyers, *Int. J. Quantum Chem.* 1 (1967) 403.
18. S. Lunell, *Chem. Phys. Lett.* 13 (1972) 93.
19. P. O. Löwdin, *Phys. Rev.* 97 (1955) 509.
20. J. Hendeković, *Int. J. Quantum Chem.* 8 (1974) 799.
21. R. E. Prat, R. Lefebvre, *Int. J. Quantum Chem.* 3 (1969) 503.
22. R. Penney, *American J. Phys.* 36 (1968) 871.
23. P. Jordan, J. V. Neumann, E. Wigner *Annals of Math.* 35 (1934) 29-64.
24. M. P. Solèr, *Comm. Algebra* 23 (1995) 219.
25. S. S. Holland, *Bull. Amer. Math. Soc.* 32 (1995) 205.
26. I. Mayer, P. O. Löwdin, *Chem. Phys. Lett.* 202 (1993) 1.
27. P. Cassam-Chenaï, *J. Chem. Phys* 116 (2002) 8677-8690.
28. P. Cassam-Chenaï, D. Jayatilaka, *J. Chem. Phys* 137 (2012) 064107. And supplementary material.
29. J. M. D. Coey, *Revue canadienne de physique* 65 (1987) 1210.
30. E. Libby, J. K. McCusker , E. A. Schmitt , K. Folting , D. N. Hendrickson , G. Christou, *Inorg. Chem.* 30 (1991) 3486.
31. D. Jayatilaka, *J. Chem. Phys* 108 (1998) 7587.
32. D. W. Small, E. J. Sundstrom, M. Head-Gordon *J. Chem. Phys* 142 (2015) 094112.
33. D. Jayatilaka and D. J. Grimwood, in *Computational Science - ICCS 2003*, (Lectures Notes in Computer Sciences Vol. 2660, Sloot P., Abramson D., Bogdanov A., Gorbachev Y., Dongarra J., and Zomaya A. editors, Springer, Berlin and Heidelberg, 2003) p.142-151.
34. L. Bučinský, M. Malček, S. Biskupič, D. Jayatilaka b, G. E. Büchel, Vl. B. Arion, *Comp. Theor. Chem.* 1065 (2015) 27.
35. T. H. Dunning, *J. Chem. Phys* 90 (1989) 1007.
36. M. Barysz, A.J. Sadlej, *J. Mol. Struct. (THEOCHEM)* 573 (2001) 181.

N_α	+4.999546
N_β	+4.000454
$(\frac{N_\alpha}{2} - \frac{N_\beta}{2})(\frac{N_\alpha}{2} - \frac{N_\beta}{2} + 1)$	+0.749091
z -noncollinearity	+0.000461
x, y -nonperpendicularity	+0.000427
spin contamination	+0.007033
$\langle S^2 \rangle$	+0.757013

Table 1 Expectation value of $\langle S^2 \rangle$ and related quantities for an H_2O^+ GCHF optimized wave function. The geometry parameters were $r_{OH} = 0.99192\text{\AA}$, $\widehat{HOH} = 101.411\text{deg}$. The basis set consisted of the primitives Gaussian functions left uncontracted of Dunning's cc-pVDZ hydrogen and oxygen basis sets [35]. The infinite-order two-component (IOTC) relativistic Hamiltonian of Barysz and Sadlej [36] was employed.