



HAL
open science

STABILITY OF THE OPTIMAL FILTER IN CONTINUOUS TIME: BEYOND THE BENEŠ FILTER

van Bien Bui, Sylvain Rubenthaler

► **To cite this version:**

van Bien Bui, Sylvain Rubenthaler. STABILITY OF THE OPTIMAL FILTER IN CONTINUOUS TIME: BEYOND THE BENEŠ FILTER. 2016. hal-01301157v3

HAL Id: hal-01301157

<https://hal.univ-cotedazur.fr/hal-01301157v3>

Preprint submitted on 27 Jan 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Stability of the optimal filter in continuous time: beyond the Beneš filter

Bui, Van Bien & Rubenthaler, Sylvain

January 27, 2020

Laboratoire J. A. Dieudonné, Université de Nice Sophia Antipolis, Parc Valrose, 06108 Nice cedex 2, FRANCE (bienmaths@gmail.com, rubentha@unice.fr)

Abstract

We are interested in the optimal filter in a continuous time setting. We want to show that the optimal filter is stable with respect to its initial condition. We reduce the problem to a discrete time setting and apply truncation techniques coming from [OR05]. Due to the continuous time setting, we need a new technique to solve the problem. In the end, we show that the forgetting rate is at least a power of the time t . The results can be re-used to prove the stability in time of a numerical approximation of the optimal filter.

filtering, signal detection, inference from stochastic processes.

1 Introduction

1.1 Exposition of the problem

We are given a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. We are interested in the processes $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$, solutions of the following SDE's in \mathbb{R}

$$X_t = X_0 + \int_0^t f(X_s) ds + V_t,$$
$$Y_t = \int_0^t h X_s ds + W_t \quad (h \neq 0),$$

where V, W are two independent standard Brownian motions, X_0 is a random variable in \mathbb{R} , of law π_0 . We set $(\mathcal{F}_t)_{t \geq 0}$ to be the filtration associated to (V_t, W_t) . For $t \geq 0$, we call optimal filter at time t the law of X_t knowing $(Y_s)_{0 \leq s \leq t}$, and we denote this law by π_t . Let $\tau > 1$, this parameter will be adjusted later. For any $t > 0$, we set Q_t to be the transition kernel of the Markov chain $(X_{kt})_{k \geq 0}$. We set $Q = Q_\tau$.

Before going further, we mention that we believe the results we have could be transposed to processes in \mathbb{R}^d for any d . This would result in a more technical paper and we chose not to pursue this goal.

Hypothesis 1. *We suppose that f is C^1 and that $\|f\|_\infty, \|f'\|_\infty$ are bounded by a constant M .*

Hypothesis 2. *We suppose that there exists positive constants v_1, v_2 such that, for all nonnegative Δ ,*

$$\pi_0([- \Delta, \Delta]^{\mathbb{C}}) \leq v_1 e^{-v_2 \Delta^2}.$$

(This means that the tail of π_0 is sub-Gaussian.)

We are interested in the stability of $(\pi_t)_{t \geq 0}$ with respect to its initial condition. As explained below in Equation (1.2), for all t , π_t can be written as a functional of $(Y_s)_{0 \leq s \leq t}$ and π_0 . Suppose now we plug a probability π'_0 instead of π_0 into this functional, we then obtain what is called a “wrongly initialized filter” π'_t (Equation (1.3)). It is natural to wonder whether $\pi_t - \pi'_t$ goes to zero when t goes to infinity in any sense, and at which rate it does so. We would then say that the filter (π_t) is stable with respect to its initial condition. We give such a result with a rate of convergence, in Theorem 1.2, under Hypothesis 1 and 2. This question has been answered for more general processes (X_t) and (Y_t) evolving in continuous time. Here is a brief review and the existing results. We stress the differences with our setting (see [CR11] for other references).

- There is a proof of stability with respect to the initial condition in [AZ97]. In this paper, the process (X_t) has to stay in a compact space. The rate is exponential.
- There are stability results in [Ata98], [KV08]. There, the process (X_t) has to satisfy some ergodicity conditions. The rate is exponential.
- In [BCL04, CvH07], the process (X_t) has to take values in a finite state space.
- The article [Sta05] is the closest to our results. There are some assumptions on the drift coefficient $f(\dots)$ (see Section 1.3 of [Sta05]), f would have to be of the form φ'/φ , $\varphi \in \mathcal{C}^2(\mathbb{R})$, which are different from ours. The rate is exponential.

In the case of an exponential rate, the filter would be called “exponentially stable” (with respect to its initial condition). The widespread idea is that exponential stability induces that a numerical approximation of the optimal filter would not deteriorate in time. Such an approximation is usually based on a time-recursive computation and it is believed that exponential stability will prevent an accumulation of errors. In order to use a stability result in a proof concerning a numerical scheme, one might like the distance between π_t and π'_t to be expressed in term of the distance between π_0 and π'_0 , and there is no such result, at least when the time is continuous. We do not prove such a result in this paper. We explain below that there is another way to prove that a numerical approximation of the optimal filter does not deteriorate in time.

Again, our aim in this paper is to show stability in such a way that the results can be used in a proof that a numerical scheme remains good uniformly in time. This has not been done in the literature we cited above. We follow [OR05] by introducing a “robust filter” restricted to compact spaces. We show that this filter remains close to the optimal filter uniformly in time and this is enough to prove the stability of the optimal filter with respect to its initial condition. As in [OR05], we do not show that the optimal filter is exponentially stable, nor can we write the dependency in π_0 , π'_0 in the stability result. However, in a future work, we will use the stability properties of the robust filter to show that there exists a numerical approximation that remains uniformly good in time.

In the case where f satisfies

$$f'(x) + f(x)^2 + h^2 x^2 = P(x), \quad (1.1)$$

where $P(x)$ is a second-order polynomial with positive leading-order coefficient, then π_t is called the Beneš filter (see [Ben81], [BC09]) and there exists an explicit formula for the density of π_t , for all t . The study of the Beneš filter is developed in [Oco99]. Under Hypothesis 1, we have (1.1) if and only if $f(x) = f(0)(1 + xf(0))^{-1}$. What we present here is a case in the neighborhood of the Beneš filter.

We present the main results and our strategy in Section 1.2 below. Before this, we have to go through some definitions.

For all $t > 0$, the law of $Y_{0:t}$ under \mathbb{P} and conditionally on X_0 , X_t has a density denoted by $y_{0:t} \mapsto \psi_t(y_{0:t}, X_0, X_t)$ with respect to the Wiener measure (see Lemma 2.2 below). The Kallianpur-Striebel formula (see [BC09], p. 57) gives us the following result (see the proof in Section 7.1).

Lemma 1.1. For all $t > 0$ and all bounded continuous φ ,

$$\pi_t(\varphi) = \frac{\int_{\mathbb{R}} \varphi(y) Q_t(x, dy) \psi_t(Y_{0:t}, x, y) \pi_0(dx)}{\int_{\mathbb{R}} Q_t(x, dy) \psi_t(Y_{0:t}, x, y) \pi_0(dx)}. \quad (1.2)$$

For any probability law π'_0 on \mathbb{R} , we define the wrongly initialized filter (with initial condition π'_0) by, for any $t > 0$,

$$\pi'_t(\varphi) = \frac{\int_{\mathbb{R}} \varphi(y) Q_t(x, dy) \psi_t(Y_{0:t}, x, y) \pi'_0(dx)}{\int_{\mathbb{R}} Q_t(x, dy) \psi_t(Y_{0:t}, x, y) \pi'_0(dx)}. \quad (1.3)$$

1.2 Main results and outline of the proofs

All the results are given under Hypothesis 1 and 2. The first main result is the following ($\|\dots\|$ is the total variation norm when applied to measures).

Theorem 1.2. There exists $\nu_0 > 0$ such that, if π_0 and π'_0 are comparable,

$$\mathbb{E}(\|\pi_t - \pi'_t\|) = O(t^{-\nu_0}),$$

when $t \rightarrow +\infty$ (see Section 3.1 for the definition of comparable).

Remark 1.3. We have here a polynomial stability whereas filters in discrete time are usually exponentially stable. We believe this result is not sharp (mainly because the proof is very convoluted).

In order to prove this Theorem, we first reduce the problem to a problem in discrete time by looking at the sequences $(\pi_0, \pi_\tau, \pi_{2\tau}, \dots)$, $(\pi'_0, \pi'_\tau, \pi'_{2\tau}, \dots)$. We then introduce a truncated filter $(\pi_{k\tau}^\Delta)_{k \geq 0}$ and a wrongly initialized truncated filter $((\pi'_{k\tau})^\Delta)_{k \geq 0}$, where Δ is a truncation parameter. We do not get into more details since to get a closed definition of the truncated filter, one would have to look at Equations (4.29), (4.28), (3.1), (4.25), (4.19), (4.20), (4.26), (4.3), (7.19), (7.4), (7.21), (2.6), (2.7), (2.8), (2.5), (2.10), (2.13) and Lemma 2.3. The truncated filter can be viewed as a restriction of $(\pi_{k\tau})_{k \geq 0}$ to some compacts. Equation (4.29) tells us that

$$\pi_{k\tau}^\Delta = \overline{R}_k^\Delta \overline{R}_{k-1}^\Delta \dots \overline{R}_1^\Delta(\pi_0), \text{ for all } k \geq 1,$$

for some operators $\overline{R}_{\dots}^\Delta$. We define

$$(\pi'_{k\tau})^\Delta = \overline{R}_k^\Delta \overline{R}_{k-1}^\Delta \dots \overline{R}_1^\Delta(\pi'_0), \text{ for all } k \geq 1.$$

We first prove that the operators $\overline{R}_{\dots}^\Delta$ are contracting (see Section 5, Equations (5.3), (5.8) for the definitions of ϵ_{\dots} , ϵ'_{\dots}).

Lemma 1.4. We suppose that τ and h are chose such that Equation (4.4) holds. For $n \geq 1$ and k in $\{1, 2, \dots, n-1\}$, for all μ, μ' in $\mathcal{P}(\mathbb{R})$,

$$\begin{aligned} & \left\| \overline{R}_n^\Delta \overline{R}_{n-1}^\Delta \dots \overline{R}_{k+1}^\Delta(\mu) - \overline{R}_n^\Delta \overline{R}_{n-1}^\Delta \dots \overline{R}_{k+1}^\Delta(\mu') \right\| \\ & \leq \prod_{i=1}^{\lfloor (n-k)/2 \rfloor} (1 - \epsilon_{k+2i-1}^2 (\epsilon'_{k+2i})^2) \times 4 \inf \left(1, \frac{\|\mu - \mu'\|}{(\epsilon'_{k+2})^2 \epsilon_{k+1}^4} \right). \end{aligned}$$

We then write a telescopic sum (Equation (6.2)) for the optimal filter

$$\|\pi_{n\tau} - \pi_{n\tau}^\Delta\| \leq \|\pi_{n\tau} - \overline{R}_n^\Delta(\pi_{(n-1)\tau})\| + \sum_{1 \leq k \leq n-1} \|\overline{R}_{n:k+1}^\Delta(\pi_{k\tau}) - \overline{R}_{n:k+1}^\Delta(\overline{R}_k^\Delta(\pi_{(k-1)\tau}))\|$$

where, for all $n \geq k$, $\overline{R}_{n:k+1}^\Delta = \overline{R}_n^\Delta \circ \overline{R}_{n-1}^\Delta \circ \dots \circ \overline{R}_{k+1}^\Delta$ is a composition of operators (and we write a similar sum for the wrongly initialized filter). Due to the contractivity of the operators $\overline{R}_{\dots}^\Delta$, we are then able to prove that the truncation error is uniform in time (see Section 2.5 for the definition of \preceq).

Proposition 1.5. *There exist a function of h , $\lambda'_1(h)$ and a constant τ_∞ such that, if $\tau \geq \tau_\infty$, we have*

$$\sup_{n \geq 0} \log(\mathbb{E}(\|\pi_{n\tau} - \pi_{n\tau}^\Delta\|)) \underset{\Delta, c}{\leq} -\Delta^2 \times \lambda'_1(h); \quad (1.4)$$

and, if π_0 and π'_0 are comparable,

$$\sup_{n \geq 0} \log(\mathbb{E}(\|\pi'_{n\tau} - (\pi')_{n\tau}^\Delta\|)) \underset{\Delta, c}{\leq} -\Delta^2 \times \lambda'_1(h). \quad (1.5)$$

(See Section 3.1 for a definition of comparable.)

The proof of Theorem 1.2 then start with the inequality

$$\|\pi_{n\tau} - \pi'_{n\tau}\| \leq \|\pi_{n\tau} - \pi_{n\tau}^\Delta\| + \|\pi_{n\tau}^\Delta - (\pi')_{n\tau}^\Delta\| + \|(\pi')_{n\tau}^\Delta - \pi'_{n\tau}\|.$$

We choose Δ as a function of n to make sure all the terms on the right go to zero when n goes to infinity. After some technicalities, we are able to get rid of the discrete time index $n\tau$ and write a result for any $t \geq 0$.

The second main result of the paper is that these results can be used to show that a numerical approximation of the filter does not deteriorate in time. Suppose we have a numerical approximation $\pi_{n\tau}^{N, \Delta}$ of $\pi_{n\tau}^\Delta$ for all $n \geq 0$ (with $\pi_0^{N, \Delta} = \pi_0^\Delta$); Δ is again the truncation parameter and N is the approximation parameter (one can think of a number of particles in case of a particle filter approximation). We write

$$\|\pi_{n\tau} - \pi_{n\tau}^{N, \Delta}\| \leq \|\pi_{n\tau} - \pi_{n\tau}^\Delta\| + \|\pi_{n\tau}^\Delta - \pi_{n\tau}^{N, \Delta}\|, \text{ for all } n \geq 0.$$

We already have the result that $\|\pi_{n\tau} - \pi_{n\tau}^\Delta\|$ is small when Δ goes to infinity (Proposition 1.5). For the second term, we write a telescopic sum

$$\|\pi_{n\tau}^{N, \Delta} - \pi_{n\tau}^\Delta\| \leq \|\pi_{n\tau}^{N, \Delta} - \bar{R}_n^\Delta(\pi_{(n-1)\tau}^{N, \Delta})\| + \sum_{1 \leq k \leq n-1} \|\bar{R}_{n:k+1}^\Delta(\pi_{k\tau}^{N, \Delta}) - \bar{R}_{n:k+1}^\Delta(\bar{R}_k^\Delta(\pi_{(k-1)\tau}^{N, \Delta}))\|.$$

Using Lemma 1.4 again, we would be able to show that $\|\pi_{n\tau}^{N, \Delta} - \pi_{n\tau}^\Delta\|$ remains small uniformly in n (when N is big). Choosing Δ as a function of N in the same way as in [OR05] would then allow to show that $\|\pi_{n\tau} - \pi_{n\tau}^{N, \Delta}\|$ remains small uniformly in the time n , when N is big.

The fact that stability results have the potential to be used to show that a numerical approximation does not deteriorate in time is a novelty compared to the other papers in the area.

Another novelty lies in the proof of Lemma 1.4. In Proposition 5.6, we exhibit a representation of the truncated filter $\bar{R}_{n:1}^\Delta(\pi_0)$ as the n -th point of a Feynman-Kac sequence based on a mixing Markov kernel (see Section 3 for the definition of the term Feynman-Kac sequence). This is not straightforward because the observations are given as a process in continuous time. Whatever transformation we adopt, the problem remains more difficult than in discrete time. We will write more comments about the novelty of the proof in Section 5 (it requires more notations to develop on the subject).

The outline of the paper is as follows. In Section 2, we compute useful estimates concerning ψ_t . In Section 3, we recall useful notions on filtering. In Section 4 and 5, we introduce the truncated filter and its properties. At the beginning of Section 5, we elaborate on our strategy. In Section 6, we prove that the optimal filter can be approximated by robust filters uniformly in time (Proposition 1.5), and that the optimal filter is stable with respect to its initial condition (Theorem 1.2).

2 Computations around ψ_t

2.1 Estimation of the transition density and the likelihood

We begin by bounding the transition density by above and below (see the proof in Section 7.1).

Lemma 2.1. For all $x, y \in \mathbb{R}$, $Q(x, dy)$ has a density $Q(x, y)$ with respect to the Lebesgue measure and

$$\frac{e^{-\frac{(y-x)^2}{2\tau}}}{\sqrt{2\pi\tau}} e^{-M|y-x|-\tau\left(\frac{M}{2}+\frac{M^2}{2}\right)} \leq Q(x, y) \leq \frac{\exp^{-\frac{(y-x)^2}{2\tau}}}{\sqrt{2\pi\tau}} e^{M|y-x|+\frac{M\tau}{2}}.$$

Following [BC09] (Chapter 6, Section 6.1), we define a new probability $\widehat{\mathbb{P}}$ by (for all $t \geq 0$)

$$\widehat{V}_t = V_t + \int_0^t f(X_s) ds,$$

$$\frac{d\widehat{\mathbb{P}}}{d\mathbb{P}} \Big|_{\mathcal{F}_t} = \widehat{Z}_t = \exp \left(\int_0^t f(X_s) d\widehat{V}_s - \frac{1}{2} \int_0^t f(X_s)^2 ds + \int_0^t hX_s dY_s - \frac{1}{2} \int_0^t h^2 X_s^2 ds \right)$$

We define, for all $0 \leq s \leq t$,

$$Y_{s:t} = (Y_u)_{s \leq u \leq t}.$$

We set

$$\widehat{\psi}_t(y_{0:t}, x_0, x_1) = \mathbb{E}^{\widehat{\mathbb{P}}} \left(\exp \left(\int_0^t hX_s dY_s - \frac{1}{2} \int_0^t (hX_s)^2 ds \right) \Big| X_0 = x_0, X_t = x_1, Y_{0:t} = y_{0:t} \right). \quad (2.1)$$

We set

$$\psi = \psi_\tau, \widehat{\psi} = \widehat{\psi}_\tau.$$

We have the following bounds for the likelihood (see the proof in Section 7.1).

Lemma 2.2. For all $t > 0$, the law of $Y_{0:t}$ under \mathbb{P} and conditionally on X_0, X_t has a density denoted by $y_{0:t} \mapsto \psi_t(y_{0:t}, X_0, X_t)$ with respect to the Wiener measure. This density satisfies, for all $x, z \in \mathbb{R}$ and any continuous trajectory $y_{0:t}$

$$\widehat{\psi}_t(y_{0:t}, x, z) e^{-2M|z-x|-\tau\left(M+\frac{M^2}{2}\right)} \leq \psi_t(y_{0:t}, x, z) \leq e^{2M|z-x|+\tau\left(M+\frac{M^2}{2}\right)} \widehat{\psi}_t(y_{0:t}, x, z).$$

2.2 Change of measure

Under $\widehat{\mathbb{P}}$, \widehat{V} is a standard Brownian motion. So, using a standard representation of a Brownian bridge, we can rewrite $\widehat{\psi}$ as

$$\widehat{\psi}(y_{0:\tau}, x, z) = \mathbb{E} \left(\exp \left(\int_0^\tau h \left(x \left(1 - \frac{s}{\tau} \right) + z \frac{s}{\tau} + \left(B_s - \frac{s}{\tau} B_\tau \right) \right) dy_s - \frac{h^2}{2} \int_0^\tau \left(x \left(1 - \frac{s}{\tau} \right) + z \frac{s}{\tau} + \left(B_s - \frac{s}{\tau} B_\tau \right) \right)^2 ds \right) \right),$$

where B is a standard Brownian motion (under \mathbb{P}). As we want to compute the above integral, where B is the only random variable involved, we can suppose that B is adapted to the filtration \mathcal{F} . We have (using the change of variable $s' = s/\tau$ and the scaling property of the Brownian motion)

$$\begin{aligned} \widehat{\psi}(y_{0:\tau}, x, z) &= \mathbb{E} \left(\exp \left(\int_0^1 h \left(x(1-s') + zs' + B_{\tau s'} - s' B_\tau \right) dy_{\tau s'} - \frac{h^2 \tau}{2} \int_0^1 \left(x(1-s') + zs' + B_{\tau s'} - s' B_\tau \right)^2 ds' \right) \right) \\ &= \mathbb{E} \left(\exp \left(\int_0^1 h \left(x(1-s') + zs' + \sqrt{\tau} (B_{s'} - s' B_1) \right) dy_{\tau s'} \right) \right) \end{aligned}$$

$$-\frac{h^2\tau}{2} \int_0^1 (x(1-s') + zs' + \sqrt{\tau}(B_{s'} - s'B_1))^2 ds') \Big) \Big)$$

In the spirit of [MY08] (Section 2.1), we define a new probability \mathbb{Q} by (for all t)

$$\frac{d\mathbb{Q}}{d\mathbb{P}} \Big|_{\mathcal{F}_t} = \exp \left(-\frac{h^2\tau^2}{2} \int_0^1 B_s^2 ds - h\tau \int_0^1 B_s dB_s \right).$$

By Girsanov's theorem, under the probability \mathbb{Q} , the process

$$\beta_t = B_t + \int_0^t h\tau B_s ds, \forall t \geq 0 \quad (2.2)$$

is a Brownian motion. We get

$$\begin{aligned} \widehat{\psi}(y_{0:\tau}, x, z) &= \exp \left(\int_0^1 h(x(1-s) + zs) dy_{\tau s} - \frac{h^2\tau}{2} \int_0^1 (x(1-s) + zs)^2 ds \right) \\ &\times \mathbb{E}^{\mathbb{Q}} \left(\exp \left(\int_0^1 h\sqrt{\tau}(B_s - sB_1) dy_{\tau s} - h^2\tau^{3/2} \int_0^1 (x(1-s) + zs)(B_s - sB_1) ds \right. \right. \\ &\quad \left. \left. - \frac{h^2\tau^2}{2} \int_0^1 s^2 B_1^2 - 2sB_s B_1 ds + h\tau \int_0^1 B_s dB_s \right) \right). \quad (2.3) \end{aligned}$$

Using the integration by parts formula, we can rewrite the last expectation as

$$\begin{aligned} \mathbb{E}^{\mathbb{Q}} \left(\exp \left(-h\sqrt{\tau} \int_0^1 (y_{\tau s} - \int_0^1 y_{\tau u} du) dB_s + h^2\tau^{3/2} \int_0^1 \left(-x \frac{(1-s)^2}{2} + z \frac{s^2}{2} + \frac{x}{6} - \frac{z}{6} \right) dB_s \right. \right. \\ \left. \left. + h^2\tau^2 B_1 \left(\frac{B_1}{3} - \int_0^1 \frac{s^2}{2} dB_s \right) + h\tau \left(\frac{B_1^2}{2} - \frac{1}{2} \right) \right) \right) = \\ \mathbb{E}^{\mathbb{Q}} \left(\exp \left(-h\sqrt{\tau} \int_0^1 (y_{\tau s} - \int_0^1 y_{\tau u} du) dB_s + h^2\tau^{3/2} x \int_0^1 s dB_s + \frac{h^2\tau^{3/2}}{2} (z-x) \int_0^1 s^2 dB_s \right. \right. \\ \left. \left. - h^2\tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) B_1 + \left(\frac{h^2\tau^2}{3} + \frac{h\tau}{2} \right) B_1^2 - \frac{h^2\tau^2}{2} B_1 \int_0^1 s^2 dB_s - \frac{h\tau}{2} \right) \right). \quad (2.4) \end{aligned}$$

2.3 Covariances computation

The last expectation contains an exponential of a polynomial of degree 2 of 4 Gaussians:

$$G_1 = B_1, G_2 = \int_0^1 s dB_s, G_3 = \int_0^1 s^2 dB_s, G_4 = \int_0^1 \left(y_{\tau s} - \int_0^1 y_{\tau u} du \right) dB_s.$$

So this expectation can be expressed as a function of the covariance matrix of these Gaussians. We compute here the covariances, which do not depend on $y_{0:\tau}$. We set

$$\theta = h\tau. \quad (2.5)$$

Lemma 2.3. *We have:*

$$\begin{aligned} \text{Var}^{\mathbb{Q}}(G_1) &= \frac{1 - e^{-2\theta}}{2\theta}, \\ \text{Var}^{\mathbb{Q}}(G_2) &= \left(1 + \frac{1}{\theta}\right)^2 \frac{(1 - e^{-2\theta})}{2\theta} + \frac{1}{\theta^2} - \left(\frac{2}{\theta^2} + \frac{2}{\theta^3}\right) (1 - e^{-\theta}), \\ \text{Var}^{\mathbb{Q}}(G_3) &= \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right)^2 \frac{(1 - e^{-2\theta})}{2\theta} + \left(\frac{2}{\theta} + \frac{2}{\theta^2}\right)^3 \frac{\theta}{6} - \frac{8}{6\theta^5} - \frac{4}{\theta^2} \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) \end{aligned}$$

$$\begin{aligned}\text{Cov}^{\mathbb{Q}}(G_1, G_2) &= \left(\frac{1}{2\theta} + \frac{1}{2\theta^2}\right) (1 - e^{-2\theta}) + \frac{e^{-\theta} - 1}{\theta^2}, \\ \text{Cov}^{\mathbb{Q}}(G_1, G_3) &= \left(\frac{1}{2\theta} + \frac{1}{\theta^2} + \frac{1}{\theta^3}\right) (1 - e^{-2\theta}) - \frac{2}{\theta^2}, \\ \text{Cov}^{\mathbb{Q}}(G_2, G_3) &= \left(1 + \frac{1}{\theta}\right) \left(\frac{1}{2\theta} + \frac{1}{\theta^2} + \frac{1}{\theta^3}\right) (1 - e^{-2\theta}) - \left(\frac{1}{\theta^2} + \frac{2}{\theta^3} + \frac{2}{\theta^4}\right) (1 - e^{-\theta}) - \frac{1}{\theta^2}.\end{aligned}$$

See the proof in Section 7.2

Let U_1, U_2, U_3, U_4 be i.i.d. of law $\mathcal{N}(0, 1)$. We can find $\alpha, \beta, \gamma, a, b, c, \lambda_1, \lambda_2, \lambda_3, \lambda_4 \in \mathbb{R}$ such that (under \mathbb{Q})

$$\begin{pmatrix} G_1 \\ G_3 \\ G_2 \\ G_4 \end{pmatrix} \stackrel{\text{law}}{=} \begin{pmatrix} \alpha U_1 \\ \beta U_1 + \gamma U_2 \\ a U_1 + b U_2 + c U_3 \\ \lambda_1 U_1 + \lambda_2 U_2 + \lambda_3 U_3 + \lambda_4 U_4 \end{pmatrix}.$$

(There is no mistake here, we do intend to look at the vector (G_1, G_3, G_2, G_4) .) Indeed, we take

$$\alpha = \sqrt{\text{Var}^{\mathbb{Q}}(G_1)}, \quad \beta = \frac{\text{Cov}^{\mathbb{Q}}(G_1, G_3)}{\alpha}, \quad \gamma = \sqrt{\text{Var}^{\mathbb{Q}}(G_3) - \beta^2}, \quad (2.6)$$

$$a = \frac{\text{Cov}^{\mathbb{Q}}(G_1, G_2)}{\alpha}, \quad b = \frac{\text{Cov}^{\mathbb{Q}}(G_2, G_3) - a\beta}{\gamma}, \quad c = \sqrt{\text{Var}^{\mathbb{Q}}(G_2) - a^2 - b^2}. \quad (2.7)$$

And we find $\lambda_1, \dots, \lambda_4$ by solving

$$\begin{cases} \alpha \lambda_1 & = \text{Cov}^{\mathbb{Q}}(G_1, G_4) \\ \beta \lambda_1 + \gamma \lambda_2 & = \text{Cov}^{\mathbb{Q}}(G_3, G_4) \\ a \lambda_1 + b \lambda_2 + c \lambda_3 & = \text{Cov}^{\mathbb{Q}}(G_2, G_4) \\ \lambda_1^2 + \lambda_2^2 + \lambda_3^2 + \lambda_4^2 & = \text{Var}^{\mathbb{Q}}(G_4). \end{cases} \quad (2.8)$$

We observe that $\alpha, \beta, \gamma, a, b, c$ can be written explicitly in terms of the parameters of the problem.

2.4 Integral computation

The last part of (2.4) is equal to

$$\begin{aligned}\mathbb{E}^{\mathbb{Q}} \left(\exp \left(\left(\frac{\theta^2}{3} + \frac{\theta}{2} \right) G_1^2 - \frac{\theta^2}{2} G_1 G_3 - h^2 \tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) G_1 \right. \right. \\ \left. \left. + h^2 \tau^{3/2} x G_2 + \frac{h^2 \tau^{3/2}}{2} (z - x) G_3 - h \sqrt{\tau} G_4 - \frac{\theta}{2} \right) \right) = \\ \int_{u_1, \dots, u_4 \in \mathbb{R}} \exp \left(\left(\frac{\theta^2}{3} + \frac{\theta}{2} \right) \alpha^2 u_1^2 - \frac{\theta^2}{2} \alpha u_1 (\beta u_1 + \gamma u_2) - h^2 \tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) \alpha u_1 \right. \\ \left. + h^2 \tau^{3/2} x (a u_1 + b u_2 + c u_3) + \frac{h^2 \tau^{3/2}}{2} (z - x) (\beta u_1 + \gamma u_2) - h \sqrt{\tau} (\lambda_1 u_1 + \dots + \lambda_4 u_4) - \frac{\theta}{2} \right) \\ \frac{\exp \left(-\frac{(u_1^2 + \dots + u_4^2)}{2} \right)}{(2\pi)^2} du_1 \dots du_4 = \\ \int_{u_1, \dots, u_4 \in \mathbb{R}} \exp \left\{ -\frac{1}{2\sigma_1^2} \left[u_1 - \sigma_1^2 \left(-\frac{\theta^2 \alpha \gamma}{2} u_2 - h^2 \tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) \alpha + h^2 \tau^{3/2} x a \right. \right. \right. \\ \left. \left. \left. + \frac{h^2 \tau^{3/2}}{2} (z - x) \beta - h \sqrt{\tau} \lambda_1 \right) \right]^2 \right. \\ \left. + \frac{\sigma_1^2}{2} \left[-\frac{\theta^2 \alpha \gamma}{2} u_2 - h^2 \tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) \alpha + h^2 \tau^{3/2} x a + \frac{h^2 \tau^{3/2}}{2} (z - x) \beta - h \sqrt{\tau} \lambda_1 \right]^2 \right\} \end{aligned}$$

$$+ h^2 \tau^{3/2} x (b u_2 + c u_3) + \frac{h^2 \tau^{3/2}}{2} (z-x) \gamma u_2 - h \sqrt{\tau} (\lambda_2 u_2 + \dots + \lambda_4 u_4) - \frac{\theta}{2} - \frac{(u_2^2 + \dots + u_4^2)}{2} \left. \right\} \frac{1}{(2\pi)^2} du_1 \dots du_4, \quad (2.9)$$

where

$$\sigma_1^2 = \left(2 \left(- \left(\frac{\theta^2}{3} + \frac{\theta}{2} \right) \alpha^2 + \frac{\theta^2 \alpha \beta}{2} + \frac{1}{2} \right) \right)^{-1}. \quad (2.10)$$

As the above expectation is finite then σ_1^2 is well defined. We set

$$m_1 = \sigma_1^2 \left(-h^2 \tau^{3/2} \left(\frac{x}{3} + \frac{z}{6} \right) \alpha + h^2 \tau^{3/2} x a + \frac{h^2 \tau^{3/2}}{2} (z-x) \beta - h \sqrt{\tau} \lambda_1 \right). \quad (2.11)$$

The above expectation (2.9) is equal to:

$$\begin{aligned} & \int_{u_1, \dots, u_4 \in \mathbb{R}} \exp \left(-\frac{1}{2\sigma_1^2} \left[u_1 + \sigma_1^2 \frac{\theta^2 \alpha \gamma}{2} u_2 - m_1 \right]^2 \right. \\ & \quad + \left(\frac{\sigma_1^2 \theta^2 \alpha \gamma}{2} \right)^2 u_2^2 \frac{1}{2\sigma_1^2} + \frac{m_1^2}{2\sigma_1^2} - \frac{1}{2\sigma_1^2} \times 2 \left(\frac{\sigma_1^2 \theta^2 \alpha \gamma}{2} \right) m_1 u_2 \\ & \quad + h^2 \tau^{3/2} x (b u_2 + c u_3) + \frac{h^2 \tau^{3/2}}{2} (z-x) \gamma u_2 \\ & \quad \left. - h \sqrt{\tau} (\lambda_2 u_2 + \lambda_3 u_3 + \lambda_4 u_4) - \frac{(u_2^2 + u_3^2 + u_4^2)}{2} - \frac{\theta}{2} \right) \frac{1}{(2\pi)^2} du_1 \dots du_4 = \\ & \int_{u_1, \dots, u_4 \in \mathbb{R}} \exp \left(-\frac{1}{2\sigma_1^2} \left[u_1 + \sigma_1^2 \frac{\theta^2 \alpha \gamma}{2} u_2 - m_1 \right]^2 \right. \\ & \quad \left. - \frac{1}{2\sigma_2^2} \left[u_2 - \sigma_2^2 \left(h^2 \tau^{3/2} x b + \frac{h^2 \tau^{3/2}}{2} (z-x) \gamma - h \sqrt{\tau} \lambda_2 - \frac{\theta^2 \alpha \gamma m_1}{2} \right) \right]^2 \right. \\ & \quad \left. + \frac{m_2^2}{2\sigma_2^2} + \frac{m_1^2}{2\sigma_1^2} + h^2 \tau^{3/2} x c u_3 - h \sqrt{\tau} (\lambda_3 u_3 + \lambda_4 u_4) - \frac{(u_3^2 + u_4^2)}{2} - \frac{\theta}{2} \right) \frac{1}{(2\pi)^2}, \quad (2.12) \end{aligned}$$

where

$$\sigma_2^2 = \left(2 \left(-\frac{\sigma_1^2 \theta^4 \alpha^2 \gamma^2}{8} + \frac{1}{2} \right) \right)^{-1}, \quad (2.13)$$

and

$$m_2 = \sigma_2^2 \left(h^2 \tau^{3/2} x b + \frac{h^2 \tau^{3/2}}{2} (z-x) \gamma - h \sqrt{\tau} \lambda_2 - \frac{\theta^2 \alpha \gamma}{2} m_1 \right). \quad (2.14)$$

Then (2.12) is equal to:

$$\begin{aligned} & \int_{u_1, \dots, u_4 \in \mathbb{R}} \exp \left(-\frac{1}{2\sigma_1^2} \left[u_1 + \sigma_1^2 \frac{\theta^2 \alpha \gamma}{2} u_2 - m_1 \right]^2 - \frac{1}{2\sigma_2^2} [u_2 - m_2]^2 + \frac{m_2^2}{2\sigma_2^2} + \frac{m_1^2}{2\sigma_1^2} \right. \\ & \quad \left. - \frac{1}{2} [u_3 - h^2 \tau^{3/2} c x + h \sqrt{\tau} \lambda_3]^2 - \frac{1}{2} [u_4 + h \sqrt{\tau} \lambda_4]^2 \right. \\ & \quad \left. + \frac{1}{2} (-h^2 \tau^{3/2} c x + h \sqrt{\tau} \lambda_3)^2 + \frac{1}{2} (-h \sqrt{\tau} \lambda_4)^2 - \frac{\theta}{2} \right) \frac{1}{(2\pi)^2} du_1 \dots du_4 = \\ & \sigma_1 \sigma_2 \exp \left(\frac{m_1^2}{2\sigma_1^2} + \frac{m_2^2}{2\sigma_2^2} + \frac{1}{2} (-h^2 \tau^{3/2} c x + h \sqrt{\tau} \lambda_3)^2 + \frac{1}{2} (h \sqrt{\tau} \lambda_4)^2 - \frac{\theta}{2} \right) \quad (2.15) \end{aligned}$$

2.5 Asymptotic $\tau \rightarrow +\infty$

From (2.3), (2.4), (2.9), (2.12), (2.15), we see that $\widehat{\psi}(y_{0:\tau}, x, z) \propto \exp(P(x, z))$ with P a polynomial of degree 2 in x, z . Let us write $-A_2(\theta)$ for the coefficient of x^2 in P , $-B_2(\theta)$ for the coefficient of z^2 in P , $C_1(\theta)$ for the coefficient of xz in P , $A_1(\theta)$ for the coefficient of x in P , $B_1(\theta)$ for the coefficient of z in P and $C_0(\theta)$ for the ‘‘constant’’ coefficient. We will write $A_1^{y_{0:\tau}}(\theta) = A_1(y_{0:\tau}, \theta)$ (or simply $A_1^{y_{0:\tau}}$), etc, when in want of stressing the dependency in y . When there will be no ambiguity, we will drop the y , superscript. The coefficients $A_2^{y_{0:\tau}}, B_2^{y_{0:\tau}}, C_1^{y_{0:\tau}}$ do not depend on y as it will be seen below. We have

$$\widehat{\psi}(y_{0:\tau}, x, z) = \sigma_1 \sigma_2 \exp(-A_2 x^2 - B_2 z^2 + A_1^{y_{0:\tau}} x + B_1^{y_{0:\tau}} z + C_1^{y_{0:\tau}} xz + C_0^{y_{0:\tau}}). \quad (2.16)$$

We are interested in the limit $\tau \rightarrow +\infty$, with h being fixed (or equivalently $\theta \rightarrow +\infty$ with h being fixed).

Lemma 2.4. *We have*

$$A_2(\theta) \xrightarrow{\theta \rightarrow +\infty} \frac{h}{2}, \quad B_2(\theta) \xrightarrow{\theta \rightarrow +\infty} \frac{h}{2}, \quad C_1(\theta) = -\frac{3h}{2\theta} + o\left(\frac{1}{\theta}\right).$$

The proof relies partly on a symbolic computation program (see Section 7.2). Let us set, for all $s \leq t$,

$$\mathcal{W}_{s,t} = \sup_{(s_1, s_2) \in [s,t]} |W_{s_1} - W_{s_2}|, \quad \mathcal{V}_{s,t} = \sup_{(s_1, s_2) \in [s,t]} |V_{s_1} - V_{s_2}|.$$

Definition 2.5. Suppose we have functions f_1, f_2 going from some set F to \mathbb{R} . We write

$$f_1 \preceq f_2$$

if there exists a constant B in \mathbb{R}_+ , which does not depend on the parameters of our problem, such that $f_1(z) \leq B f_2(z)$, for all z in F .

In the particular case when we are dealing with functions of a parameter $\Delta \in \mathbb{R}$, we write

$$f_1 \underset{\Delta}{\preceq} f_2 \text{ or } f_1(\Delta) \underset{\Delta}{\preceq} f_2(\Delta)$$

if there exists a constant B_1 in \mathbb{R}_+ , which does not depend on the parameters of our problem, and a constant Δ_0 , which may depend on the parameters of our problem, such that

$$\Delta \geq \Delta_0 \Rightarrow f_1(\Delta) \leq B_1 f_2(\Delta).$$

If, in addition, Δ_0 depends continuously on the parameter τ , we write

$$f_1 \underset{\Delta, c}{\preceq} f_2.$$

The notation $\underset{\Delta}{\preceq}$ is akin to the notation $O(\dots)$. It has the advantage that one can single out which asymptotic we are studying.

We state here (without proof) useful properties concerning the above Definition.

Lemma 2.6. *Suppose we have functions f, f_1, f_2, h_1, h_2 .*

1. *If $f \leq f_1 + f_2$ and $f_1 \preceq f_2$ then $f \preceq f_2$.*
2. *If $f \leq f_1 + f_2$ and $\log(f_1) \underset{\Delta}{\preceq} h_1$ and $\log(f_2) \underset{\Delta}{\preceq} h_2$, with $h_1(\Delta), h_2(\Delta) \xrightarrow{\Delta \rightarrow \infty} -\infty$, then $\log(f) \underset{\Delta}{\preceq} \sup(h_1, h_2)$.*

3. If we have $f_1 \preceq_{\Delta, c} f_2$, say for $\tau \geq \tau_0$ ($\tau_0 > 0$), then there exists a constant B_1 and a continuous function Δ_0 such that, for all $\tau \geq \tau_0$ and $\Delta \geq \Delta_0(\tau)$, $f_1(\Delta) \leq B_1 f_2(\Delta)$. In particular, for any $\tau_1 > \tau_0$, if $\tau \in [\tau_0, \tau_1]$ and $\Delta \geq \sup_{t \in [\tau_0, \tau_1]} \Delta_0(t)$ then $f_1(\Delta) \leq B_1 f_2(\Delta)$.

We have the following ‘‘variational’’ bounds on B_1 (see the proof in Section 7.2).

Lemma 2.7. For all $k \in \mathbb{N}$,

$$|B_1(Y_{k\tau:(k+1)\tau}, \theta) - B_1(Y_{(k+1)\tau:(k+2)\tau}, \theta)| \leq Mh\tau + h\mathcal{V}_{k\tau, (k+2)\tau} + (h + \frac{1}{\tau})\mathcal{W}_{k\tau, (k+2)\tau},$$

and

$$|B_1(Y_{0:\tau}, \theta)| \leq Mh\tau + h\mathcal{V}_{0, 2\tau} + (h + \frac{1}{\tau})\mathcal{W}_{0, 2\tau}.$$

3 Definitions and useful notions

We follow here the ideas of [OR05].

3.1 Notations

We state here notations and definitions that will be useful throughout the paper.

- The set \mathbb{R}, \mathbb{R}^2 are endowed, respectively, with $\mathcal{B}(\mathbb{R}), \mathcal{B}(\mathbb{R}^2)$, their Borel tribes.
- The elements of \mathbb{R}^2 are treated like line vectors. If $(x, z) \in \mathbb{R}^2$ and $\delta > 0$, $B((x, z), \epsilon) = \{(x', z') \in \mathbb{R}^2 : \sqrt{(x - x')^2 + (z - z')^2} < \epsilon\}$ (the ball of center (x, z) and radius ϵ). The superscript T denotes the transposition. For example, if $(x, z) \in \mathbb{R}^2$, then $(x, z)^T$ is a column vector.
- The set of probability distributions on a measurable space (E, \mathcal{F}) and the set of nonnegative measures on (E, \mathcal{F}) are denoted by $\mathcal{P}(E)$ and $\mathcal{M}^+(E)$ respectively. We write $\mathcal{C}(E)$ for the set of continuous function on a topological space E and $\mathcal{C}_b^+(E)$ for the set of bounded, continuous, nonnegative functions on E .
- When applied to measures, $\|\dots\|$ stands for the total variation norm (for μ, ν probabilities on a measurable space (F, \mathcal{F}) , $\|\mu - \nu\| = \sup_{A \in \mathcal{F}} |\mu(A) - \nu(A)|$).
- For any nonnegative kernel K on a measurable space E and any $\mu \in \mathcal{M}^+(E)$, we set

$$K\mu(dv') = \int_E \mu(dv) K(v, dv').$$

- If we have a sequence of nonnegative kernels K_1, K_2, \dots on some measured spaces E_1, E_2, \dots (meaning that for all $i \geq 1$, $x \in E_{i-1}$, $K_i(x, \cdot)$ is a nonnegative measure on E_i , then for all $i < j$, we define the kernel

$$K_{i+1:j}(x_i, dx_j) = \int_{x_{i+1} \in E_{i+1}} \dots \int_{x_{j-1} \in E_{j-1}} K_{i+1}(x_i, dx_{i+1}) K_{i+2}(x_{i+1}, dx_{i+2}) \dots K_j(x_{j-1}, dx_j).$$

- For any measurable space E and any nonzero $\mu \in \mathcal{M}^+(E)$, we define the normalized nonnegative measure,

$$\bar{\mu} = \frac{1}{\mu(E)} \mu.$$

- For any measurable space E and any nonnegative kernel K defined on E , we define the normalized nonnegative nonlinear operator \overline{K} on $\mathcal{M}^+(E)$, taking values in $\mathcal{P}(E)$, and defined by

$$\overline{K}(\mu) = \frac{K\mu}{(K\mu)(E)} = \frac{K\overline{\mu}}{(K\overline{\mu})(E)} = \overline{K}(\overline{\mu}), \quad (3.1)$$

for any $\mu \in \mathcal{M}^+(E)$ such that $K\mu(E) \neq 0$, and defined by $\overline{K}(\mu) = 0$ otherwise.

- A kernel K from a measurable space E_1 into another measurable space E_2 is said to be ϵ -mixing ($\epsilon \in (0, 1)$) if there exists λ in $\mathcal{M}^+(E_2)$ and $\epsilon_1, \epsilon_2 > 0$ such that, for all x_1 in E_1 ,

$$\epsilon_1 \lambda(\cdot) \leq K(x_1, \cdot) \leq \frac{1}{\epsilon_2} \lambda(\cdot) \text{ with } \epsilon_1 \epsilon_2 = \epsilon^2.$$

This property implies that, for all A , μ , $\overline{K}(\mu)(A) \geq \epsilon^2 \overline{\lambda}(A)$. If \overline{K} is Markov, this last inequality implies that \overline{K} is $(1 - \epsilon^2)$ -contracting in total variation (see [DG01] p. 161-162 for more details):

$$\forall \mu, \nu \in \mathcal{P}(E), \|\overline{K}(\mu) - \overline{K}(\nu)\| \leq (1 - \epsilon^2) \|\mu - \nu\|.$$

- For any measurable space E and any $\psi : E \rightarrow \mathbb{R}^+$ (measurable) and $\mu \in \mathcal{M}^+(E)$, we set

$$\langle \mu, \psi \rangle = \int_E \psi(x) \mu(dx).$$

If in addition, $\langle \mu, \psi \rangle > 0$, we set

$$\psi \bullet \mu(dv) = \frac{1}{\langle \mu, \psi \rangle} \times \psi(v) \mu(dv).$$

- For μ and μ' in $\mathcal{M}^+(E)$ ((E, \mathcal{F}) being a measurable space), we say that μ and μ' are comparable if there exist positive constants a and b such that, for all $A \in \mathcal{F}$,

$$a\mu'(A) \leq \mu(A) \leq b\mu'(A).$$

We then define the Hilbert metric between μ and μ' by

$$h(\mu, \mu') = \log \left(\frac{\sup_{A \in \mathcal{F}: \mu'(A) > 0} \frac{\mu(A)}{\mu'(A)}}{\inf_{A \in \mathcal{F}: \mu'(A) > 0} \frac{\mu(A)}{\mu'(A)}} \right).$$

It is easily seen (see for instance [Oud00], Chapter 2) that, for any nonnegative kernel K and any A in \mathcal{F} ,

$$h(K\mu, K\mu') \leq h(\mu, \mu'), \quad (3.2)$$

$$h(\overline{\mu}, \overline{\mu}') \leq h(\mu, \mu'), \quad (3.3)$$

$$\exp(-h(\mu, \mu')) \leq \frac{\mu(A)}{\mu'(A)} \leq \exp(h(\mu, \mu')), \text{ if } \mu'(A) > 0. \quad (3.4)$$

In addition, we have the following relation with the total variation norm:

$$\|\overline{\mu} - \overline{\mu}'\| \leq \frac{2}{\log(3)} h(\mu, \mu'). \quad (3.5)$$

- We set \tilde{Q} to be the transition of the chain $(X_{k\tau}, X_{(k+1)\tau})_{k \geq 0}$.
- We write \propto between two quantities if they are equal up to a multiplicative constant.

- For $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}$, we write $\psi(0, \cdot)$ for the function such that, for all x in \mathbb{R} , $\psi(0, \cdot)(x) = \psi(0, x)$.
- We write λ_W for the Wiener measure on $\mathcal{C}([0, \tau])$.

We suppose here that the observation $(Y_t)_{t \geq 0}$ is fixed. For $k \in \mathbb{N}^*$ and $x, z \in \mathbb{R}$, we define

$$\psi_k(x, z) = \psi(Y_{(k-1)\tau:k\tau}, x, z) \quad (3.6)$$

(the density ψ is defined in Lemma 2.2). For $x_1 \in \mathbb{R}$, $x_2 \in \mathbb{R}$ and $n \in \mathbb{N}^*$, we introduce the nonnegative kernel

$$R_n(x_1, dx_2) = \psi_n(x_1, x_2)Q(x_1, dx_2). \quad (3.7)$$

Using the above notations, we now have, for all $n \in \mathbb{N}^*$, and for all probability law π'_0 (with $(\pi'_t)_{t \geq 0}$ defined in Equation (1.3))

$$\pi_{n\tau} = \overline{R}_n(\pi_{(n-1)\tau}), \pi'_{n\tau} = \overline{R}_n(\pi'_{(n-1)\tau})$$

and for $0 < m < n$,

$$\pi_{n\tau} = \overline{R}_n \overline{R}_{n-1} \dots \overline{R}_m(\pi_{(m-1)\tau}), \pi'_{n\tau} = \overline{R}_n \overline{R}_{n-1} \dots \overline{R}_m(\pi'_{(m-1)\tau}). \quad (3.8)$$

3.2 Representation of the optimal filter as the law of a Markov chain

Regardless of the notations of the other sections, we suppose we have a Markov chain $(\mathfrak{X}_n)_{n \geq 0}$ taking values in measured spaces E_0, E_1, \dots , with nonnegative kernels $\Omega_1, \Omega_2, \dots$ (it might be a non-homogeneous Markov chain). Suppose we have potentials $\Psi_1 : E_1 \rightarrow \mathbb{R}_+, \Psi_2 : E_2 \rightarrow \mathbb{R}_+, \dots$ (measurable functions with values in \mathbb{R}_+) and a law η_0 on E_0 . We are interested in the sequence of probability measures $(\eta_k)_{k \geq 1}$, respectively on E_1, E_2, \dots , defined by

$$\forall k \geq 1, \forall \varphi \in \mathcal{C}_b^+(E_k), \eta_k(f) = \frac{\mathbb{E}_{\eta_0}(\varphi(\mathfrak{X}_k) \prod_{1 \leq i \leq k} \Psi_i(\mathfrak{X}_i))}{\mathbb{E}_{\eta_0}(\prod_{1 \leq i \leq k} \Psi_i(\mathfrak{X}_i))}, \quad (3.9)$$

where $\eta_0 \in \mathcal{P}(E_0)$ and the index η_0 means we start with \mathfrak{X}_0 of law η_0 . We will say that $(\eta_k)_{k \geq 0}$ is a Feynman-Kac sequence on $(E_k)_{k \geq 0}$ based on the transitions $(\Omega_k)_{k \geq 1}$, the potentials $(\Psi_k)_{k \geq 1}$ and the initial law η_0 . Suppose we have another law η'_0 , we then set

$$\forall k \geq 1, \forall \varphi \in \mathcal{C}_b^+(E_k), \eta'_k(f) = \frac{\mathbb{E}_{\eta'_0}(\varphi(\mathfrak{X}_k) \prod_{1 \leq i \leq k} \Psi_i(\mathfrak{X}_i))}{\mathbb{E}_{\eta'_0}(\prod_{1 \leq i \leq k} \Psi_i(\mathfrak{X}_i))}.$$

If the functions Ψ_k 's are likelihood associated to observations of a Markov chain with transitions $\Omega_1, \Omega_2, \dots$ and initial law η_0 , then the measures η_k 's are optimal filters. We fix $n \geq 1$. We would like to express η_n as the marginal law of some Markov process. We will do so using ideas from [DG01]. We set, for all $k \in \{1, \dots, n\}$,

$$\mathfrak{R}_k(x, dx') = \Psi_k(x')\Omega_k(x, dx'). \quad (3.10)$$

We suppose that, for all k , \mathfrak{R}_k is ϵ_k -mixing (notice that Ω_k being ϵ_k -mixing implies that \mathfrak{R}_k is ϵ_k -mixing). By a simple recursion, we have, for all n ,

$$\overline{\mathfrak{R}}_{1:n} = \overline{\mathfrak{R}}_n \overline{\mathfrak{R}}_{n-1} \dots \overline{\mathfrak{R}}_1. \quad (3.11)$$

We set, for all $k \in \{0, 1, \dots, n-1\}$,

$$\Psi_{n|k}(x) = \int_{x_{k+1} \in E_{k+1}} \dots \int_{x_n \in E_n} \mathfrak{R}_{k+1}(x, dx_{k+1}) \prod_{k+2 \leq i \leq n} \mathfrak{R}_i(x_{i-1}, dx_i).$$

If $k = n$, we set $\Psi_{n|n}$ to be constant equal to 1. For $k \in \{1, 2, \dots, n\}$, we set

$$\mathfrak{S}_{n|k}(x, dx') = \frac{\Psi_{n|k+1}(x')}{\Psi_{n|k}(x)} \mathfrak{R}_{k+1}(x, dx').$$

From [DG01], we get the following result (a simple proof can also be found in [OR05], Proposition 3.1).

Proposition 3.1. *The operators $(\mathfrak{S}_{n|k})_{0 \leq k \leq n-1}$ are Markov kernels. For all $k \in \{0, \dots, n-1\}$, $\mathfrak{S}_{n|k}$ is ϵ_{k+1} -mixing. We have*

$$\eta_n = \mathfrak{S}_{n|n-1} \mathfrak{S}_{n|n-2} \dots \mathfrak{S}_{n|0} (\Psi_{n|0} \bullet \eta_0),$$

$$\eta'_n = \mathfrak{S}_{n|n-1} \mathfrak{S}_{n|n-2} \dots \mathfrak{S}_{n|0} (\Psi_{n|0} \bullet \eta'_0),$$

and

$$\|\eta_n - \eta'_n\| \leq \prod_{1 \leq k \leq n} (1 - \epsilon_k^2) \times \|\Psi_{n|0} \bullet \eta_0 - \Psi_{n|0} \bullet \eta'_0\|.$$

Following the computations of [OR05], p. 434 (or [Oud00], p.66), we have, for all measurable $\Psi : \mathbb{R}^2 \rightarrow \mathbb{R}^+$,

$$\|\Psi \bullet \eta_0 - \Psi \bullet \eta'_0\| \leq 2 \int_{x \in E_0} \frac{\Psi(x)}{\langle \eta_0, \Psi \rangle} |\eta_0 - \eta'_0|(dx) \leq 2 \inf \left(1, \frac{\|\Psi\|_\infty}{\langle \eta_0, \Psi \rangle} \|\eta_0 - \eta'_0\| \right). \quad (3.12)$$

For all x in E_0 , as \mathfrak{R}_1 is ϵ_1 -mixing,

$$\begin{aligned} \frac{\Psi_{n|0}(x)}{\langle \eta_0, \Psi_{n|0} \rangle} &= \frac{\int_{z \in E_1} \mathfrak{R}_{2:n}(z, E_n) \mathfrak{R}_1(x, dz)}{\int_{y \in E_0} \int_{z \in E_1} \mathfrak{R}_{2:n}(z, E_n) \mathfrak{R}_1(y, dz) \eta_0(dy)} \\ (\text{for some } \epsilon'_1, \epsilon''_1, \lambda_1 \text{ with } \epsilon'_1 \epsilon''_1 &= \epsilon_1^2) &\leq \frac{\int_{z \in E_1} \mathfrak{R}_{2:n}(z, E_n) \frac{1}{\epsilon'_1} \lambda_1(dz)}{\int_{z \in E_1} \mathfrak{R}_{2:n}(z, E_n) \epsilon''_1 \lambda_1(dz)} = \frac{1}{\epsilon_1^2}. \end{aligned} \quad (3.13)$$

4 Truncated filter

We introduce in this section a filter built with truncated likelihoods. We will call it truncated filter or robust filter, the use of the adjective ‘‘robust’’ refers to the fact that it has stability properties (it appears in the proof of Proposition 1.5 below).

4.1 Integrals of the potential

We are look here at $\widehat{\psi}(y_{0:\tau}, x, z)$ for some x, z in \mathbb{R} and a fixed observation $y_{0:\tau}$ between 0 and τ . All what will be said is also true for observations between $k\tau$ and $(k+1)\tau$ (for any k in \mathbb{N}). From Equations (7.21), (7.23), (7.24), we see that $A_1^{y_{0:\tau}}$, $B_1^{y_{0:\tau}}$ are polynomials of degree 1 in $\lambda_1, \dots, \lambda_3$ and that C_1 does not depend on $y_{0:\tau}$. We fix x and z in \mathbb{R} . Recall that, by Equation (2.8), Lemmas 7.1 and 7.2, $\lambda_1, \lambda_2, \lambda_3$ are functions of $y_{0:\tau}$ and that they can be expressed as integrals of deterministic functions against $dy_{\tau s}$ (this requires some integrations by parts, see Lemma 7.4). Under the law $\widetilde{\mathbb{P}}$ (defined in Equation (7.1)), conditioned to $X_0 = x, X_\tau = z$, we can write

$$X_t = \left(1 - \frac{s}{\tau}\right) x + \frac{s}{\tau} z + \widetilde{B}_s - \frac{s}{\tau} \widetilde{B}_\tau, \quad (4.1)$$

where $(\widetilde{B}_s)_{s \geq 0}$ is a Brownian motion, independent of W . And we can write, using integration by parts and Equation (7.24) (see Lemma 7.5),

$$A_1^{Y_{0:\tau}} = \int_0^\tau (f_1(s) dW_s + f_2(s) dX_s), \quad (4.2)$$

for some deterministic functions f_1, f_2 (and the same goes for $B_1^{Y_{0:\tau}}$, see Equation (7.23)).

We set

$$p_{1,1} = \left(1 - \frac{C_1^2}{4A_2B_2}\right)_+^{1/2}, \quad p_{2,1} = -\frac{C_1}{2B_2}, \quad p_{2,2} = 1. \quad (4.3)$$

Fact 4.1. *We now fix a parameter $\iota \in (1/2, 1)$. From now on, we suppose that the parameter τ is chosen such that*

$$\theta \geq 1, \quad A_2 \geq \frac{h}{4}, \quad B_2 \geq \frac{h}{4}, \quad C_1 \leq \frac{h}{8}, \quad \frac{1}{1+p_{2,1}} - \left(2 + \frac{4}{h}\right)B_2p_{2,1}\theta^{1-\iota} > 0, \quad p_{1,1} > \frac{1}{2}, \quad |p_{2,1}| \leq \frac{1}{2}. \quad (4.4)$$

This is possible because of Lemma 2.4 and because this Lemma implies: $p_{2,1} = O(\theta^{-1})$, $p_{1,1} \xrightarrow{\theta \rightarrow +\infty} 1$.

Let us set

$$\kappa = \begin{bmatrix} A_2 & -\frac{C_1}{2} \\ -\frac{C_1}{2} & B_2 \end{bmatrix}. \quad (4.5)$$

If we take

$$P = \begin{bmatrix} p_{1,1} & 0 \\ p_{2,1} & p_{2,2} \end{bmatrix} = \begin{bmatrix} \left(1 - \frac{C_1^2}{4A_2B_2}\right)^{1/2} & 0 \\ -\frac{C_1}{2B_2} & 1 \end{bmatrix}, \quad (4.6)$$

then

$$\kappa = P^T \begin{bmatrix} A_2 & 0 \\ 0 & B_2 \end{bmatrix} P. \quad (4.7)$$

We have

$$P^{-1} = \begin{bmatrix} \frac{1}{p_{1,1}} & 0 \\ -\frac{p_{2,1}}{p_{1,1}} & 1 \end{bmatrix}.$$

First, we have to rule out the case where $A_1^{y_{0:\tau}}$ and $B_1^{y_{0:\tau}}$ are colinear.

Lemma 4.2. *The quantities $A_1^{y_{0:\tau}}$ and $B_1^{y_{0:\tau}}$ are not colinear (as functions of $y_{0:\tau}$).*

Proof. Suppose there exists $\lambda \in \mathbb{R}$ such that $B_1^{y_{0:\tau}} = \lambda A_1^{y_{0:\tau}}$ for λ_W -almost all $y_{0:\tau}$. We have, for all φ in $\mathcal{C}_b^+(\mathbb{R})$, using Lemma 2.2 (remember Equations (2.3), (2.4), (2.9), (2.12), (2.15))

$$\begin{aligned} & \int_{\mathcal{C}([0;\tau])} \varphi(A_1^{y_{0:\tau}}) \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \\ & \leq e^{2M|x-z|+\tau(M+\frac{M^2}{2})} \int_{\mathcal{C}([0;\tau])} \varphi(A_1^{y_{0:\tau}}) \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \\ & = \sigma_1 \sigma_2 e^{2M|x-z|+\tau(M+\frac{M^2}{2})} \int_{\mathbb{R}} \varphi(t) \exp(-A_2x^2 - B_2z^2 + C_1xz + tx + \lambda tz) \Psi'(t) dt, \quad (4.8) \end{aligned}$$

where

$$\Psi'(t) = \mathbb{E}^{\mathbb{P}}(\exp(C_0^{W_{0:\tau}}) | A_1^{W_{0:\tau}} = t).$$

We know the last integral is finite (because $\int_{\mathcal{C}([0,1])} \psi(y_{0:\tau}, x, z) = 1$ and because of Lemma 2.2).

We introduce Ψ'_1 such that

$$\Psi'(t) = \exp\left(-\frac{1}{4}(t, \lambda t) \kappa^{-1}(t, \lambda t)^T\right) \Psi'_1(t),$$

and

$$\forall (t_1, t_2) \in \mathbb{R}^2, \quad \mathcal{Q}(t_1, t_2) = \exp\left(-\frac{1}{4}(t_1, t_2) \kappa^{-1}(t_1, t_2)^T\right).$$

We have, for all t , (remember that $\tilde{\mathbb{P}}$ is defined in Equation (7.1))

$$e^{-M|X_0 - X_t| - \frac{Mt}{2} - \frac{M^2 t}{2}} \leq \frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} \Big|_{\mathcal{F}_t} \leq e^{M|X_t - X_0| + \frac{Mt}{2}}$$

(this can be deduced from the computations in the proof of Lemma 2.1). Thus, we have, for all φ (the first equality being a consequence of Lemma 2.2)

$$\begin{aligned} \int_{\mathcal{C}([0;1])} \varphi(A_1^{y_{0:\tau}}) \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) &= \mathbb{E}^{\mathbb{P}}(\varphi(A_1^{Y_{0:\tau}}) | X_0 = x, X_\tau = z) \\ &= \frac{\mathbb{E}^{\tilde{\mathbb{P}}}\left(\varphi(A_1^{Y_{0:\tau}}) \frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} \Big|_{\mathcal{F}_\tau} \Big| X_0 = x, X_\tau = z\right)}{\mathbb{E}^{\tilde{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\tilde{\mathbb{P}}} \Big|_{\mathcal{F}_\tau} \Big| X_0 = x, X_\tau = z\right)} \\ &\stackrel{\text{(by Equation (4.2))}}{\geq} e^{-2M|x-z|-\tau\left(M+\frac{M^2}{2}\right)} \int_{\mathbb{R}} \varphi(t) \mathcal{Q}''_{x,z}(t) dt \end{aligned} \quad (4.9)$$

for some Gaussian density $\mathcal{Q}''_{x,z}$ (the density of $A_1^{Y_{0:\tau}}$ knowing $X_0 = x, X_\tau = z$, under $\tilde{\mathbb{P}}$). From Equations (4.8), (4.9), we deduce that, for (x, z) fixed, we have for almost all t ,

$$e^{-2M|x-z|-\tau\left(M+\frac{M^2}{2}\right)} \mathcal{Q}''_{x,z}(t) \leq e^{2M|x-z|+\tau\left(M+\frac{M^2}{2}\right)} \sigma_1 \sigma_2 \mathcal{Q}((t, \lambda t) - 2(x, z)\kappa) \Psi'_1(t). \quad (4.10)$$

In the same way, for (x, z) fixed, we have for almost all t ,

$$e^{2M|x-z|+\tau\left(M+\frac{M^2}{2}\right)} \mathcal{Q}''_{x,z}(t) \geq e^{-2M|x-z|-\tau\left(M+\frac{M^2}{2}\right)} \sigma_1 \sigma_2 \mathcal{Q}((t, \lambda t) - 2(x, z)\kappa) \Psi'_1(t). \quad (4.11)$$

Looking at Equations (4.1), (4.2), we can say that the density $\mathcal{Q}''_{x,z}(t)$ is of the form

$$\mathcal{Q}''_{x,z}(t) = \frac{1}{\sqrt{2\pi\sigma_0^2}} \exp\left(-\frac{1}{2\sigma_0^2}(t - (a_0x + b_0z))^2\right),$$

with σ_0, a_0, b_0 independent of (x, z) . So, looking at the above inequalities in $(x, z) = (0, 0)$, we see there exists $\epsilon > 0$ and a constant C_ϵ , such that, for almost all t in $(-\epsilon, \epsilon)$,

$$(C_\epsilon \sigma_1 \sigma_2)^{-1} e^{-\tau(2M+M^2)} \leq \Psi'_1(t) \leq C_\epsilon (\sigma_1 \sigma_2)^{-1} e^{\tau(2M+M^2)}. \quad (4.12)$$

For any t , the quantities $\log(\mathcal{Q}''_{x,z}(t))$, $\log(\mathcal{Q}((t, \lambda t) - 2(x, z)\kappa))$ are polynomials in x, z , of degree less than 2. Using the above remarks and studying adequate sequences $(x_n, z_n)_{n \geq 0}$ (for example, with $x_n \xrightarrow{n \rightarrow +\infty} +\infty$, z_n remaining in a neighborhood of 0), one can show that the coefficients in x^2, z^2 and xz in these two polynomials the same. We then have

$$\frac{a_0^2}{2\sigma_0^2} = A_2, \quad \frac{b_0^2}{2\sigma_0^2} = B_2, \quad \frac{a_0 b_0}{\sigma_0^2} = C_1.$$

By Equation (4.4), we have

$$\frac{a_0 b_0}{\sigma_0^2} = 2\sqrt{A_2 B_2} \geq \frac{h}{2} > \frac{h}{8}$$

and $C_1 \leq h/8$, which is not possible, hence the result. \square

We can now write for any test function φ in $\mathcal{C}_b^+([0, \tau])$ (remember Equation (2.16))

$$\int_{\mathcal{C}([0, \tau])} \varphi(A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) =$$

$$\sigma_1 \sigma_2 \int_{\mathbb{R}^k} \varphi(t_1, t_2) \exp[-A_2 x^2 - B_2 z^2 + C_1 xz + t_1 x + t_2 z] \times \Psi(t_1, t_2) dt_1 dt_2, \quad (4.13)$$

where

$$\Psi(t_1, t_2) = \mathbb{E}^{\mathbb{P}}(\exp(C_0^{W_0:\tau}) | A_1^{W_0:\tau} = t_1, B_1^{W_0:\tau} = t_2).$$

We know the integral is finite (because $\int_{\mathcal{C}([0,1])} \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) = 1$ and because of Lemma 2.2). Let us define Ψ_1 by the formula

$$\Psi(t_1, t_2) = \exp\left(-\frac{1}{4}(t_1, t_2) \kappa^{-1}(t_1, t_2)^T\right) \Psi_1(t_1, t_2).$$

The next result tells us that, somehow, $\log(\Psi_1(t_1, t_2))$ is negligible before $t_1^2 + t_2^2$ (when $(t_1, t_2) \rightarrow +\infty$).

Lemma 4.3. *There exists a constant $C'_1(h, \tau)$ (continuous in (h, τ)) and $\epsilon > 0$ such that for all (x, z) and for almost all (t_1, t_2) in $B(2(x, z)\kappa, \epsilon)$,*

$$\begin{aligned} \frac{1}{C'_1(h, \tau)} \exp(-4M|x-z| - \tau(2M + M^2)) &\leq \Psi_1(t_1, t_2) \\ &\leq C'_1(h, \tau) \exp(4M|x-z| + \tau(2M + M^2)). \end{aligned} \quad (4.14)$$

Proof. We fix (x, z) in \mathbb{R}^2 . Similarly as in Equation (4.8), we get, using Lemma 2.2, for all $\varphi \in \mathcal{C}_b^+(\mathbb{R}^2)$,

$$\begin{aligned} &\int_{\mathcal{C}([0,\tau])} \varphi(A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \\ &\leq \int_{\mathcal{C}([0,\tau])} \varphi(A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) e^{2M|x-z| + \tau(M + \frac{M^2}{2})} \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \\ &= \sigma_1 \sigma_2 e^{2M|x-z| + \tau(M + \frac{M^2}{2})} \int_{\mathbb{R}^2} \varphi(t_1, t_2) e^{-A_2 x^2 - B_2 z^2 + C_1 xz + t_1 x + t_2 z} \Psi(t_1, t_2) dt_1 dt_2 \\ &= \sigma_1 \sigma_2 e^{2M|x-z| + \tau(M + \frac{M^2}{2})} \int_{\mathbb{R}^2} \varphi(t_1, t_2) \exp\left(-\frac{1}{4}((t_1, t_2) - 2(x, z)\kappa)^{-1}((t_1, t_2)^T - 2\kappa(x, z)^T)\right) \\ &\quad \times \Psi_1(t_1, t_2) dt_1 dt_2. \end{aligned} \quad (4.15)$$

Similarly as in Equation (4.9), we get, for all φ ,

$$\begin{aligned} &\int_{\mathcal{C}([0,\tau])} \varphi(A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \leq e^{2M|z-x| + \tau(M + \frac{M^2}{2})} \int_{\mathbb{R}^2} \varphi(t_1, t_2) \mathcal{Q}'_{x,z}(t_1, t_2) dt_1 dt_2, \\ &e^{-2M|z-x| - \tau(M + \frac{M^2}{2})} \int_{\mathbb{R}^2} \varphi(t_1, t_2) \mathcal{Q}'_{x,z}(t_1, t_2) dt_1 dt_2 \leq \int_{\mathcal{C}([0,\tau])} \varphi(A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) \psi(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}), \end{aligned}$$

for some Gaussian density $\mathcal{Q}'_{x,z}$ with covariance matrix which does not depend on x, z (see Equation (4.2)). This is the density of $(A_1^{Y_{0:\tau}}, B_1^{Y_{0:\tau}})$ knowing $X_0 = x$ and $X_\tau = z$, under $\widetilde{\mathbb{P}}$. We then have, a.s. in (t_1, t_2) (for the Lebesgue measure),

$$\mathcal{Q}'_{x,z}(t_1, t_2) \leq \sigma_1 \sigma_2 e^{4M|x-z| + \tau(2M + M^2)} \mathcal{Q}((t_1, t_2) - 2(x, z)\kappa) \Psi_1(t_1, t_2). \quad (4.16)$$

Using the lower bound in the inequality in Lemma 2.2, we get in the same way, a.s. in (t_1, t_2) ,

$$\mathcal{Q}'_{x,z}(t_1, t_2) \geq \sigma_1 \sigma_2 e^{-4M|x-z| - \tau(2M + M^2)} \mathcal{Q}((t_1, t_2) - 2(x, z)\kappa) \Psi_1(t_1, t_2). \quad (4.17)$$

We deduce from Equation (4.17), that there exists $\epsilon_1 > 0$ such that, for all (x, z) and for almost all (t_1, t_2) in $B(2(x, z)\kappa, \epsilon_1)$

$$\Psi_1(t_1, t_2) \leq C'_1(\tau, h) e^{4M|x-z| + \tau(2M + M^2)},$$

for some function $C'_1(\tau, h)$ of the parameters τ, h (continuous in (h, τ)). Looking at Equations (4.1), (4.2), one can also see that $\mathcal{Q}'_{x,z}$ reaches its maximum at $(x, z)\kappa'$, where κ' is fixed in $\mathcal{M}_{2,2}(\mathbb{R})$ (the set of 2×2 matrices with coefficients in \mathbb{R}). From Equation (4.16), we get that there exists $\epsilon_2 > 0$ such that, for all x, z and for almost all (t_1, t_2) in $B((x, z)\kappa', \epsilon'_2)$,

$$\begin{aligned} \mathcal{Q}((t_1, t_2) - 2(x, z)\kappa) &\geq \frac{1}{2} \mathcal{Q}'_{x,z}((x, z)\kappa') (\sigma_1 \sigma_2)^{-1} e^{-4M|x-z|-\tau(4M+2M^2)} (C'_1(h, \tau))^{-1} \\ &\quad \times \exp\left(-4M \left| \frac{1}{2}(x, z)\kappa'\kappa^{-1}(1, -1)^T \right| \right), \end{aligned}$$

and so, by continuity,

$$\begin{aligned} \mathcal{Q}((x, z)\kappa' - 2(x, z)\kappa) &\geq \frac{1}{2} \mathcal{Q}'_{x,z}((x, z)\kappa') (\sigma_1 \sigma_2)^{-1} e^{-4M|x-z|-\tau(4M+2M^2)} (C'_1(h, \tau))^{-1} \\ &\quad \times \exp\left(-4M \left| \frac{1}{2}(x, z)\kappa'\kappa(1, -1)^T \right| \right), \end{aligned}$$

If $\kappa' \neq 2\kappa$, we can find a sequence (x_n, z_n) such that $x_n^2 + z_n^2 \xrightarrow{n \rightarrow +\infty} +\infty$ and

$$\log(\mathcal{Q}((x_n, z_n)\kappa' - 2(x_n, z_n)\kappa)) \preceq -(x_n^2 + z_n^2),$$

whereas

$$\begin{aligned} \log\left(\mathcal{Q}'_{x_n, z_n}((x_n, z_n)\kappa') (\sigma_1 \sigma_2)^{-1} e^{-4M|x_n-z_n|-\tau(4M+2M^2)} \exp\left(-4M \left| \frac{1}{2}(x, z)\kappa'\kappa^{-1}(1, -1)^T \right| \right)\right) \\ \succeq -|x_n| - |z_n|, \end{aligned}$$

which is not possible. So $\kappa' = 2\kappa$.

So, we get from Equations (4.16), (4.17) that there exists $\epsilon_3 > 0$ such that for all x, z , and for almost all (t_1, t_2) in $B(2(x, z)\kappa, \epsilon_3)$,

$$C'_1(h, \tau) e^{4M|x-z|+\tau(2M+M^2)} \geq \Psi_1(t_1, t_2) \geq \frac{e^{-4M|x-z|-\tau(2M+M^2)}}{C'_1(h, \tau)}$$

(with, possibly, a new $C'_1(h, \tau)$). □

Lemma 4.4. *If we have a set $\mathcal{A} = \{y_{0:\tau} \in \mathcal{C}([0; \tau]) : (A_1^{y_{0:\tau}}, B_1^{y_{0:\tau}}) \in \mathcal{B}\}$ for some measurable subset \mathcal{B} of \mathbb{R}^2 , then*

$$\begin{aligned} \int_{\mathcal{A}} \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) &\leq \sigma_1 \sigma_2 C'_1(h, \tau) \\ &\quad \times \int_{(t_2, t_2) \in \mathcal{B}} \exp\left\{-\frac{A_2^{-1}}{4} \left(\left| \frac{t_1}{p_{1,1}} - \frac{p_{2,1}t_2}{p_{1,1}} - 2A_2 p_{1,1}x \right| - 4M \left| \frac{1}{p_{1,1}} + \frac{p_{2,1}}{p_{1,1}} \right| \right)_+^2 \right. \\ &\quad \left. - \frac{B_2^{-1}}{4} (|t_2 - 2B_2(p_{2,1}x + z)| - 4M)_+^2 \right\} \\ &\quad \times \exp(4M^2(1, -1)\kappa^{-1}(1, -1)^T + \tau(2M + M^2) + 4M|x - z|) dt_1 dt_2, \end{aligned}$$

and

$$\begin{aligned} \int_{\mathcal{A}} \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) &\geq \sigma_1 \sigma_2 C'_1(h, \tau)^{-1} \\ &\quad \times \int_{(t_2, t_2) \in \mathcal{B}} \exp\left\{-\frac{A_2^{-1}}{4} \left(\left| \frac{t_1}{p_{1,1}} - \frac{p_{2,1}t_2}{p_{1,1}} - 2A_2 p_{1,1}x \right| + 4M \left| \frac{1}{p_{1,1}} + \frac{p_{2,1}}{p_{1,1}} \right| \right)_+^2 \right. \\ &\quad \left. - \frac{B_2^{-1}}{4} (|t_2 - 2B_2(p_{2,1}x + z)| + 4M)_+^2 \right\} \\ &\quad \times \exp(4M^2(1, -1)\kappa^{-1}(1, -1)^T - \tau(2M + M^2) - 4M|x - z|) dt_1 dt_2. \end{aligned}$$

Proof. We have (computing as in Equation (4.15))

$$\begin{aligned}
& \int_{\mathcal{A}} \widehat{\psi}(y_{0:\tau}, x, z) \lambda_W(dy_{0:\tau}) \\
&= \sigma_1 \sigma_2 \int_{\mathcal{B}} \exp(-(x, z) \kappa(x, z)^T + t_1 x + t_2 z) \exp\left(-\frac{1}{4}(t_1, t_2) \kappa^{-1}(t_1, t_2)^T\right) \Psi_1(t_1, t_2) dt_1 dt_2 \\
&\quad \text{(by Lemma 4.3, for a complete proof see Lemma 7.6)} \\
&\leq \sigma_1 \sigma_2 \int_{\mathcal{B}} \exp\left(-\frac{1}{4}[(t_1, t_2)^T - 2\kappa(x, z)^T]^T \kappa^{-1}[(t_1, t_2)^T - 2\kappa(x, z)^T]\right) \\
&\quad \times C'_1(h, \tau) \exp(2M \times |(1, -1) \kappa^{-1}(t_1, t_2)^T| + \tau(2M + M^2)) dt_1 dt_2,
\end{aligned}$$

and we can bound by below by

$$\begin{aligned}
& \sigma_1 \sigma_2 \int_{\mathcal{B}} \exp\left(-\frac{1}{4}[(t_1, t_2)^T - 2\kappa(x, z)^T]^T \kappa^{-1}[(t_1, t_2)^T - 2\kappa(x, z)^T]\right) \\
&\quad \times \frac{1}{C'_1(h, \tau)} \exp(-2M \times |(1, -1) \kappa^{-1}(t_1, t_2)^T| - \tau(2M + M^2)) dt_1 dt_2,
\end{aligned}$$

For $(t_1, t_2) \in \mathbb{R}^2$, we have, for any $\delta \in \{-1, 1\}$

$$\begin{aligned}
& \exp\left(-\frac{1}{4}[(t_1, t_2)^T - 2\kappa(x, z)^T]^T \kappa^{-1}[(t_1, t_2)^T - 2\kappa(x, z)^T] + 2\delta M \times (1, -1) \kappa^{-1}(t_1, t_2)^T\right) \\
&= \exp\left(-\frac{1}{4}[(t_1, t_2)^T - 2\kappa(x, z)^T - 4\delta M(1, -1)^T]^T \kappa^{-1}[(t_1, t_2)^T - 2\kappa(x, z)^T - 4\delta M(1, -1)^T]\right. \\
&\quad \left.+ 4M^2(1, -1) \kappa^{-1}(1, -1)^T + 4\delta M(x, z)(1, -1)^T\right) \\
&= \exp\left(-\frac{1}{4}[(P^{-1})^T(t_1, t_2)^T - 2 \begin{bmatrix} A_2 & 0 \\ 0 & B_2 \end{bmatrix} P(x, z)^T - 4\delta M(P^{-1})^T(1, -1)^T]^T\right. \\
&\quad \times \left. \begin{bmatrix} A_2^{-1} & 0 \\ 0 & B_2^{-1} \end{bmatrix} [(P^{-1})^T(t_1, t_2)^T - 2 \begin{bmatrix} A_2 & 0 \\ 0 & B_2 \end{bmatrix} P(x, z)^T - 4\delta M(P^{-1})^T(1, -1)^T]\right) \\
&\quad \times \exp(4M^2(1, -1) \kappa^{-1}(1, -1)^T + 4\delta M(x, z)(1, -1)^T).
\end{aligned}$$

From there, we get the result. \square

4.2 Truncation

In the following, the parameter $\Delta > 0$ is to be understood as a truncation level. For $k \geq 0$ and $\Delta > 0$, we set (for all b)

$$C_{k+1}(\Delta, b) = \{z : |2B_2(1 + p_{2,1})z - b| \leq \Delta\} \quad (4.18)$$

(which indeed does not depend on k),

$$C_{k+1}(\Delta) = C_{k+1}(\Delta, B_1^{y_{k\tau:(k+1)\tau}}) \quad (4.19)$$

and

$$m_{k+1}(b) = \frac{b}{2B_2(1 + p_{2,1})}, \quad (4.20)$$

(which indeed does not depend on k) and

$$m_{k+1} = m_{k+1}(B_1^{Y_{k\tau:(k+1)\tau}}).$$

We suppose that m_0 is a point in the support of π_0 (the law of X_0) and we set

$$C_0(\Delta) = \left[m_0 - \frac{\Delta}{2B_2(1+p_{2,1})}, m_0 + \frac{\Delta}{2B_2(1+p_{2,1})} \right].$$

Under Hypothesis 2, there exists constants C_0 and C'_0 such that

$$\pi_0(C_0(\Delta)^{\mathfrak{G}}) \leq C'_0 e^{-C_0 \Delta^2}, \quad \forall \Delta > 0.$$

From Equation (4.4) and Lemma 2.7, we see that there exists a universal constant C such that

$$|m_k - m_{k-1}| \leq \begin{cases} C(M\tau + \mathcal{V}_{(k-2)\tau, k\tau} + (1 + \frac{1}{\theta})\mathcal{W}_{(k-2)\tau, k\tau}) & \text{if } k \geq 2, \\ |m_0| + C(M\tau + \mathcal{V}_{0, 2\tau} + (1 + \frac{1}{\theta})\mathcal{W}_{0, 2\tau}) & \text{if } k = 1. \end{cases} \quad (4.21)$$

We set

$$d(\Delta) = \frac{\Delta}{1+p_{2,1}} - (2 + \frac{16}{h})B_2 p_{2,1} \theta^{1-\iota} \Delta - 4M, \quad (4.22)$$

(because of Equation (4.4), we have $d(\Delta) \xrightarrow{\Delta \rightarrow +\infty} +\infty$) and

$$\begin{aligned} T(\Delta) &= \frac{C\sqrt{\tau} \exp\left(-\frac{1}{2} \left(\frac{\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right)^2\right)}{\theta^{1-\iota}\Delta} \\ &+ \left(M \frac{(1+p_{2,1})}{p_{1,1}} + \sqrt{A_2}\right) C'_1(h, \tau) \sigma_1 \sigma_2 p_{1,1} \frac{B_2 e^{26M\tau + \frac{7\tau M}{2} + 40M^2/h}}{d(\Delta)} \exp\left(-\frac{1}{4B_2} d(\Delta)^2\right) + e^{-C_0 \Delta^2}. \end{aligned} \quad (4.23)$$

We define, for all $k \geq 1$, x and x' in \mathbb{R} (recall that ψ_k is defined in Equation 3.6)

$$\psi_k^\Delta(x, x') = \psi_k(x, x') \mathbb{1}_{C_k(\Delta)}(x'), \quad (4.24)$$

$$D_k = |m_k - m_{k-1}|, \quad (4.25)$$

and for $D \geq 0$,

$$\xi_1(D, \Delta) = \frac{1}{\sqrt{2\pi\tau}} \exp\left(-\frac{\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right)^2}{2\tau}\right) \exp\left(-M\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right) - \left(\frac{\tau}{2} + \frac{\tau^2}{2}\right)M\right), \quad (4.26)$$

$$\xi_2(D, \Delta) = \frac{1}{\sqrt{2\pi\tau}} \exp\left(-\frac{\left(\left(D - \frac{\Delta}{B_2(1+p_{2,1})}\right)_+\right)^2}{2\tau}\right) \exp\left(M\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right) + \frac{\tau}{2}M\right), \quad (4.27)$$

and,

$$R_k^\Delta(x, dx') = \begin{cases} \psi_k^\Delta(x, x') Q(x, dx') & \text{if } x \in C_{k-1}(\Delta), \\ \psi_k^\Delta(x, x') \xi_1(D_k, \Delta) dx' & \text{if } x \notin C_{k-1}(\Delta). \end{cases} \quad (4.28)$$

We define $(\pi_n^\Delta)_{n \geq 0}$ by the following

$$\begin{cases} \pi_0^\Delta = \pi_0 \\ \pi_k^\Delta = \overline{R}_k^\Delta \overline{R}_{k-1}^\Delta \dots \overline{R}_1^\Delta(\pi_0) \quad \text{for all } k \geq 1. \end{cases} \quad (4.29)$$

The next lemma tells us that the measures π_k are concentrated on the compacts $C_k(\Delta)$.

Lemma 4.5. *If*

$$\theta^{1-\iota}\Delta > 3|m_0| + 3CM\tau \text{ and } d(\Delta) > 0, \quad (4.30)$$

then we have, for all $k \geq 0$,

$$\mathbb{E}(\pi_{k\tau}(C_k(\Delta))^{\mathfrak{G}}) \leq T(\Delta).$$

Proof. We suppose $k \geq 1$ (the proof is similar for $k = 0$). For a measure μ in $\mathcal{M}^+(\mathbb{R})$, we define

$$\tilde{Q}\mu(dx, dx') = \mu(dx)Q(x, dx'), \quad \forall x, x' \in \mathbb{R}, \quad (4.31)$$

(recall \tilde{Q} has been defined as a Markov kernel on \mathbb{R}^2 , so the above is an extension of the definition of \tilde{Q}). We have

$$\begin{aligned} & \pi_k(C_k(\Delta))^{\mathfrak{G}} \times \mathbb{1}_{|m_k - m_{k-1}| \leq \Delta\theta^{1-\iota}} = \\ & \mathbb{1}_{|m_k - m_{k-1}| \leq \Delta\theta^{1-\iota}} \int_{(x, x') \in \mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q}\pi_{k-1}, \psi_k \rangle} \mathbb{1}_{C_k(\Delta)^{\mathfrak{G}}}(x) (\mathbb{1}_{C_{k-1}(2\Delta)}(x) + \mathbb{1}_{C_{k-1}(2\Delta)^{\mathfrak{G}}}(x)) \tilde{Q}\pi_{k-1}(dx, dx'), \end{aligned}$$

and (using the same computations as in [LO03], proof of Proposition 5.3, [Oud00], p. 66)

$$\begin{aligned} & \mathbb{E} \left(\mathbb{1}_{|m_k - m_{k-1}| \leq \Delta\theta^{1-\iota}} \int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q}\pi_{k-1}, \psi_k \rangle} \mathbb{1}_{C_k(\Delta)^{\mathfrak{G}}}(x') \mathbb{1}_{C_{k-1}(2\Delta)}(x) \tilde{Q}\pi_{k-1}(dx, dx') \Big| Y_{0:(k-1)\tau} \right) \\ & = \int_{y \in \mathcal{C}([0, \tau])} \mathbb{1}_{|m_k(B_1^y) - m_{k-1}| \leq \Delta\theta^{1-\iota}} \\ & \quad \times \left(\int_{\mathbb{R}^2} \frac{\psi(y, x, x')}{\langle \tilde{Q}\pi_{k-1}, \psi(y, \cdot, \cdot) \rangle} \mathbb{1}_{C_k(\Delta, B_1^y)^{\mathfrak{G}}}(x') \mathbb{1}_{C_{k-1}(2\Delta)}(x) \tilde{Q}\pi_{k-1}(dx, dx') \right) \\ & \quad \times \left(\int_{(u, u') \in \mathbb{R}^2} \tilde{Q}\pi_{k-1}(u, du') \psi(y, u, u') \right) \lambda_W(dy) \\ & \quad \text{(by Fubini's theorem)} \\ & = \int_{y \in \mathcal{C}([0, 1])} \mathbb{1}_{|m_k(B_1^y) - m_{k-1}| \leq \Delta\theta^{1-\iota}} \int_{\mathbb{R}^2} \psi(y, x, x') \mathbb{1}_{C_k(\Delta, B_1^y)^{\mathfrak{G}}}(x') \mathbb{1}_{C_{k-1}(2\Delta)}(x) \tilde{Q}\pi_{k-1}(dx, dx') \lambda_W(dy) \\ & \quad \text{(using Lemma 2.2 and Lemma 4.4)} \\ & \leq \sigma_1 \sigma_2 C'_1(h, \tau) \int_{(x, x') \in \mathbb{R}^2} \int_{(t_1, t_2) \in \mathbb{R}^2 : |m_k(t_2) - m_{k-1}| \leq \Delta\theta^{1-\iota}} e^{6M|x-x'| + \tau(3M+3M^2/2) + 4M^2|(1, -1)\kappa^{-1}(1, -1)^T|} \\ & \quad \times e^{-\frac{1}{4A_2} \left(\left| \frac{t_1}{p_{1,1}} - \frac{p_{2,1}t_2}{p_{1,1}} - 2A_2 p_{1,1}x \right| - 4M \left| \frac{1+p_{2,1}}{p_{1,1}} \right| \right)_+^2} \times e^{-\frac{1}{4B_2} \left(|t_2 - 2B_2(p_{2,1}x + x')| - 4M \right)_+^2} \\ & \quad \times \mathbb{1}_{C_k(\Delta, t_2)^{\mathfrak{G}}}(x') \mathbb{1}_{C_{k-1}(2\Delta)}(x) dt_1 dt_2 \tilde{Q}\pi_{k-1}(dx, dx'). \quad (4.32) \end{aligned}$$

By a similar computation, we get

$$\begin{aligned} & \mathbb{E} \left(\int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q}\pi_{k-1}, \psi_k \rangle} \mathbb{1}_{C_k(\Delta)^{\mathfrak{G}}}(x) \mathbb{1}_{C_{k-1}(2\Delta)^{\mathfrak{G}}}(x) \tilde{Q}\pi_{k-1}(dx, dx') \Big| Y_{0:(k-1)\tau} \right) \\ & = \int_{y \in \mathcal{C}([0, 1])} \int_{\mathbb{R}^2} \psi(y, x, x') \mathbb{1}_{C_k(\Delta, B_1^y)^{\mathfrak{G}}}(x') \mathbb{1}_{C_{k-1}(2\Delta)^{\mathfrak{G}}}(x) \tilde{Q}\pi_{k-1}(dx, dx') \lambda_W(dy) \\ & \leq \int_{\mathbb{R}^2} \mathbb{1}_{C_{k-1}(2\Delta)^{\mathfrak{G}}}(x) \tilde{Q}\pi_{k-1}(dx, dx') \leq \pi_{k-1}(C_{k-1}(2\Delta))^{\mathfrak{G}}. \end{aligned}$$

For all t_2 such that $|m_k(t_2) - m_{k-1}| \leq \theta^{1-\iota}\Delta$, for $x \in C_{k-1}(2\Delta)$, $x' \in C_k(\Delta, t_2)^{\mathfrak{G}}$,

$$|t_2 - 2B_2(p_{2,1}x + x')|$$

$$\begin{aligned}
&= \left| -2B_2x' + \frac{t_2}{p_{2,1}+1} + \frac{p_{2,1}}{p_{2,1}+1}t_2 - 2B_2p_{2,1}x \right| \\
&\geq \frac{\Delta}{1+p_{2,1}} - 2B_2p_{2,1}|m_k(t_2) - m_{k-1}| - 2B_2p_{2,1}|m_{k-1} - x| \\
&\quad (\text{using Equation (4.4)}) \geq \frac{\Delta}{1+p_{2,1}} - \left(2 + \frac{16}{h}\right)B_2p_{2,1}\theta^{1-\iota}\Delta. \quad (4.33)
\end{aligned}$$

So

$$\begin{aligned}
\mathbb{E}(\pi_k(C_k(\Delta)^{\mathfrak{L}})) &\leq \mathbb{P}(|m_k - m_{k-1}| \geq \theta^{1-\iota}\Delta) + \mathbb{E}(\pi_{k-1}(C_{k-1}(2\Delta)^{\mathfrak{L}})) \\
&\quad + \sigma_1\sigma_2C'_1(h, \tau) \int_{(x, x') \in \mathbb{R}^2} \int_{(t_1, t_2) \in \mathbb{R}^2 : |t_2 - 2B_2(p_{2,1}x + x')| > \frac{\Delta}{1+p_{2,1}} - (2 + \frac{16}{h})\Delta\theta^{1-\iota}B_2p_{2,1}} \\
&\quad e^{6M|x-x'| + \tau(3M+3M^2/2) + 4M^2(1, -1)\kappa^{-1}(1, -1)^T} \\
&\quad \times e^{-\frac{1}{4A_2} \left(\left| \frac{t_1}{p_{1,1}} - \frac{p_{2,1}t_2}{p_{1,1}} - 2A_2p_{1,1}x \right| - 4M \left| \frac{1+p_{2,1}}{p_{1,1}} \right| \right)_+^2} \\
&\quad \times e^{-\frac{1}{4B_2} (|t_2 - 2B_2(p_{2,1}x + x')| - 4M)_+^2} dt_1 dt_2 \tilde{Q}\pi_{k-1}(dx, dx').
\end{aligned}$$

We have, for all $x \geq 0$,

$$\begin{aligned}
\mathbb{P}(\mathcal{V}_{0,2\tau} \geq x) &= \mathbb{P}\left(\sup_{s \in [0, 2\tau]} \bar{V}_s - \inf_{s \in [0, 2\tau]} V_s \geq x\right) \\
&\leq \mathbb{P}\left(\sup_{s \in [0, 2\tau]} V_s \geq x/2\right) + \mathbb{P}\left(-\inf_{s \in [0, 2\tau]} V_s \geq x/2\right) \\
&= 4\mathbb{P}(V_{2\tau} \geq x/2) \\
&= 2\mathbb{P}(2|V_{2\tau}| \geq x).
\end{aligned}$$

And so, we can bound (for all x)

$$\mathbb{P}(\mathcal{V}_{(k-2)_+, k\tau} \geq x) \leq 2\mathbb{P}(2|W_{2\tau}| \geq x), \quad (4.34)$$

$$\mathbb{P}(\mathcal{W}_{(k-2)_+, k\tau} \geq x) \leq 2\mathbb{P}(2|W_{2\tau}| \geq x). \quad (4.35)$$

So (with the constant C defined in Equation (4.21)), using the inequality

$$\forall z > 0, \int_z^{+\infty} \frac{\exp\left(-\frac{t^2}{2\sigma^2}\right)}{\sqrt{2\pi\sigma^2}} dt \leq \frac{\sigma \exp\left(-\frac{z^2}{2\sigma^2}\right)}{z\sqrt{2\pi}}, \quad (4.36)$$

and using Equation (4.21), as $\theta^{1-\iota}\Delta > 3|m_0| + 3CM\tau$, we get

$$\begin{aligned}
\mathbb{P}(|m_k - m_{k-1}| \geq \theta^{1-\iota}\Delta) &\leq 4\mathbb{P}\left(2 \times |W_{2\tau}| \geq \frac{\theta^{1-\iota}\Delta}{6C}\right) \\
&= 4\mathbb{P}\left(|W_1| \geq \frac{\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right) \\
&\leq \frac{96C\sqrt{\tau}}{\theta^{1-\iota}\Delta\sqrt{\pi}} \exp\left(-\frac{1}{2} \left(\frac{\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right)^2\right). \quad (4.37)
\end{aligned}$$

For all x, x' , we have

$$\begin{aligned}
&\int_{(t_1, t_2) \in \mathbb{R}^2 : |t_2 - 2B_2(p_{2,1}x + x')| > \frac{\Delta}{1+p_{2,1}} - (2 + \frac{16}{h})\Delta\theta^{1-\iota}B_2p_{2,1}} \\
&\quad e^{-\frac{1}{4A_2} \left(\left| \frac{t_1}{p_{1,1}} - \frac{p_{2,1}t_2}{p_{1,1}} - 2A_2p_{1,1}x \right| - 4M \left| \frac{1+p_{2,1}}{p_{1,1}} \right| \right)_+^2}
\end{aligned}$$

$$\begin{aligned}
& \times e^{-\frac{1}{4B_2}(|t_2 - 2B_2(p_{2,1}x + x')| - 4M)_+^2} dt_1 dt_2 \\
& \text{(change of variables : } \begin{pmatrix} t'_1 \\ t'_2 \end{pmatrix} = (P^{-1})^T \begin{pmatrix} t_1 \\ t_2 \end{pmatrix} \text{)} \\
& = \int_{(t'_1, t'_2) \in \mathbb{R}^2 : |t'_2 - 2B_2(p_{2,1}x + x')| > \frac{\Delta}{1+p_{2,1}} - (2 + \frac{16}{h})\Delta\theta^{1-\iota} B_2 p_{2,1}} e^{-\frac{1}{4A_2}(|t'_1 - 2A_2 p_{1,1}x| - 4M|\frac{1+p_{2,1}}{p_{1,1}}|)_+^2} \\
& \quad \times e^{-\frac{1}{4B_2}(|t'_2 - 2B_2(p_{2,1}x + x')| - 4M)_+^2} p_{1,1} dt'_1 dt'_2 \\
& \quad \text{(by (4.36))} \leq \left(8M \frac{|1 + p_{2,1}|}{p_{1,1}} + 2\sqrt{\pi A_2} \right) \\
& \quad \times \frac{4B_2}{d(\Delta)} \exp\left(-\frac{1}{4B_2}d(\Delta)^2\right) p_{1,1}. \quad (4.38)
\end{aligned}$$

We have, by Lemma 2.1 and Equation (4.4),

$$\begin{aligned}
& \int_{(x, x') \in \mathbb{R}^2} e^{6M|x-x'| + \tau(3M+3M^2/2) + 4M^2(1, -1)\kappa^{-1}(1, -1)^T} \tilde{Q}\pi_{k-1}(dx, dx') \\
& \leq \int_{(x, x') \in \mathbb{R}^2} \frac{1}{\sqrt{2\pi\tau}} \exp\left(-\frac{(x-x')^2}{2\tau} + 7M|x-x'| + \tau\left(\frac{7}{2}M + \frac{3}{2}M^2\right) + 40M^2\right) dx' \pi_{k-1}(dx) \\
& \leq 2 \exp\left(\frac{49}{2}M^2\tau + \tau\left(\frac{7}{2}M + \frac{3}{2}M^2\right) + \frac{40M^2}{h}\right). \quad (4.39)
\end{aligned}$$

So we have

$$\begin{aligned}
\mathbb{E}(\pi_k(C_k(\Delta)^{\mathfrak{G}})) & \leq \mathbb{E}(\pi_{k-1}(C_{k-1}(2\Delta)^{\mathfrak{G}})) + \frac{96C\sqrt{\tau}}{\theta^{1-\iota}\Delta\sqrt{\pi}} \exp\left(-\frac{1}{2}\left(\frac{\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right)^2\right) \\
& \quad + \left(8M \frac{|1 + p_{2,1}|}{p_{1,1}} + 2\sqrt{\pi A_2}\right) C'_1(h, \tau)\sigma_1\sigma_2 p_{1,1} \frac{8B_2}{d(\Delta)} \\
& \quad \times \exp\left(-\frac{1}{4B_2}d(\Delta)^2 + 26M^2\tau + \frac{7}{2}\tau M + \frac{40M^2}{h}\right).
\end{aligned}$$

Then, by recurrence,

$$\begin{aligned}
\mathbb{E}(\pi_k(C_k(\Delta)^{\mathfrak{G}})) & \leq \mathbb{E}(\pi_0(C_0(2^k\Delta)^{\mathfrak{G}})) + \sum_{i=0}^{k-1} \left[\frac{96C\sqrt{\tau}}{2^i\theta^{1-\iota}\Delta\sqrt{\pi}} \exp\left(-\frac{1}{2}\left(\frac{2^i\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right)^2\right) \right. \\
& \quad \left. + \left(8M \frac{|1 + p_{2,1}|}{p_{1,1}} + 2\sqrt{\pi A_2}\right) C'_1(h, \tau)\sigma_1\sigma_2 p_{1,1} \frac{8B_2 e^{26M^2\tau + \frac{7}{2}\tau M + 40M^2/h}}{d(2^i\Delta)} \exp\left(-\frac{1}{4B_2}d(2^i\Delta)^2\right) \right].
\end{aligned}$$

We have

$$\begin{aligned}
\frac{1}{d(\Delta)} + \frac{1}{d(2\Delta)} + \frac{1}{d(4\Delta)} + \dots & \leq \frac{1}{d(\Delta)} + \frac{1}{d(\Delta)} + \frac{1}{d(\Delta) + 2\Delta \left(\frac{1}{1+p_{2,1}} - (2 + \frac{16}{h})B_2 p_{2,1}\theta^{1-\iota}\right)} \\
& \quad + \frac{1}{d(\Delta) + 4\Delta \left(\frac{1}{1+p_{2,1}} - (2 + \frac{16}{h})B_2 p_{2,1}\theta^{1-\iota}\right)} + \dots \\
& \leq \frac{2}{d(\Delta)} + \frac{1}{\Delta \left(\frac{1}{1+p_{2,1}} - (2 + \frac{16}{h})B_2 p_{2,1}\theta^{1-\iota}\right)} \leq \frac{3}{d(\Delta)}.
\end{aligned}$$

So

$$\begin{aligned} \mathbb{E}(\pi_k(C_k(\Delta)^{\mathfrak{G}})) &\leq \mathbb{E}(\pi_0(C_0(2^k \Delta)^{\mathfrak{G}})) + \frac{192C\sqrt{\tau}}{\theta^{1-\iota}\Delta\sqrt{\pi}} \exp\left(-\frac{1}{2}\left(\frac{\theta^{1-\iota}\Delta}{12C\sqrt{2\tau}}\right)^2\right) \\ &\quad + \left(8M\frac{|1+p_{2,1}|}{p_{1,1}} + 2\sqrt{\pi A_2}\right) C'_1(h, \tau)\sigma_1\sigma_2 p_{1,1} 8B_2 e^{26M^2\tau + \frac{7}{2}\tau M + 40M^2/h} \\ &\quad \times \frac{3}{d(\Delta)} \exp\left(-\frac{1}{4B_2}d(\Delta)^2\right). \end{aligned}$$

And we can conclude because of Hypothesis 2. \square

Corollary 4.6. *We suppose that $\pi'_0 \in \mathcal{P}(\mathbb{R})$ is such that π_0 and π'_0 are comparable. We suppose that $(\pi'_t)_{t \geq 0}$ is defined by Equation (1.3). Under the assumption of the previous Lemma, we have, for all $k \geq 0$,*

$$\mathbb{E}(\pi'_{k\tau}(C_k(\Delta)^{\mathfrak{G}})) \leq T(\Delta)e^{h(\pi_0, \pi'_0)}.$$

Proof. By Equations (3.2), (3.3), (3.8), we have, for all k ,

$$h(\pi'_{k\tau}, \pi_{k\tau}) \leq h(\pi'_0, \pi_0). \quad (4.40)$$

So, by Equation (3.4),

$$\begin{aligned} \mathbb{E}(\pi'_{k\tau}(C_k(\Delta)^{\mathfrak{G}})) &\leq \mathbb{E}(e^{h(\pi'_0, \pi_0)}\pi_{k\tau}(C_k(\Delta)^{\mathfrak{G}})) \\ &\leq T(\Delta)e^{h(\pi_0, \pi'_0)}. \end{aligned}$$

\square

The next result tells us that \bar{R}_k and \bar{R}_k^Δ ($k \geq 1$) are close in some sense (recall that $\pi'_{k\tau} = \bar{R}_k(\pi'_{(k-1)\tau})$)

Proposition 4.7. *We suppose that Δ satisfies the assumption of the previous Lemma (Equation (4.30)). We suppose that $(\pi'_{n\tau})_{n \geq 0}$ satisfies the assumptions of the above Corollary. For all $k \geq 1$, we have*

$$\mathbb{E}(\|\pi'_{k\tau} - \bar{R}_k^\Delta(\pi'_{(k-1)\tau})\|) \leq T(\Delta)e^{2h(\pi_0, \pi'_0)}.$$

Proof. We define measures on \mathbb{R}^2 (remember Equation (4.31)):

$$\begin{aligned} \mu &= \tilde{Q}\pi'_{(k-1)\tau} \\ &= \tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)^{\mathfrak{G}}}\pi'_{(k-1)\tau})(dx, dx') + (\mathbb{1}_{C_k(\Delta)}(x') + \mathbb{1}_{C_k(\Delta)^{\mathfrak{G}}}(x'))\tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)}\pi'_{(k-1)\tau})(dx, dx'), \end{aligned}$$

$$\begin{aligned} \mu'(dx, dx') &= \mathbb{1}_{C_k(\Delta)}(x')\tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)}\pi'_{(k-1)\tau})(dx, dx') \\ &\quad + \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}})\xi_1(D_k, \Delta)\mathbb{1}_{C_k(\Delta)}(x')dx dx', \end{aligned}$$

where (by a slight abuse of notation)

$$\tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)}\pi'_{(k-1)\tau})(dx, dx') = \mathbb{1}_{C_{k-1}(\Delta)}(x)\pi'_{(k-1)\tau}(dx)Q(x, dx'),$$

$$\tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)^{\mathfrak{G}}}\pi'_{(k-1)\tau})(dx, dx') = \mathbb{1}_{C_{k-1}(\Delta)^{\mathfrak{G}}}(x)\pi'_{(k-1)\tau}(dx)Q(x, dx'),$$

By the definition of R^Δ (Equation (4.28)) and computing as in [OR05], p. 433 (or as in [Oud00], p.66), we get

$$\begin{aligned} \|\pi'_{k\tau} - \bar{R}_k^\Delta(\pi'_{(k-1)\tau})\| &= \|\psi_k \bullet \mu - \psi_k \bullet \mu'\| \\ \text{(using Equation (3.12))} &\leq 2 \int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q}\pi'_{k-1}, \psi_k \rangle} \times [\mathbb{1}_{C_k(\Delta)^{\mathfrak{G}}}(x')\tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)}\pi'_{(k-1)\tau})(dx, dx')] \end{aligned}$$

$$\begin{aligned}
& + \tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)^c} \pi'_{(k-1)\tau})(dx, dx') \\
& + \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}}) \xi_1(D_k, \Delta) \mathbb{1}_{C_k(\Delta)}(x') dx dx'.
\end{aligned}$$

We have, by Lemma 2.2,

$$\begin{aligned}
& \mathbb{E}(\mathbb{1}_{[0, \theta^{1-\iota} \Delta]} (|m_k - m_{k-1}|) \\
& \quad \times \int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q} \pi'_{k-1}, \psi_k \rangle} \times \mathbb{1}_{C_k(\Delta)^c}(x') \tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)} \pi'_{(k-1)\tau})(dx, dx') | Y_{0:(k-1)\tau}) \\
& \quad = \int_{y_{0:\tau} \in \mathcal{C}([0, \tau])} \mathbb{1}_{[0, \theta^{1-\iota} \Delta]} (|m_k(B_1^{y_{0:\tau}}) - m_{k-1}(B_1^{Y_{(k-2)\tau:(k-1)\tau}})|) \\
& \quad \times \int_{\mathbb{R}^2} \frac{\psi(y_{0:\tau}, x, x')}{\langle \tilde{Q} \pi'_{k-1}, \psi(y_{0:\tau}, \cdot, \cdot) \rangle} \mathbb{1}_{C_k(\Delta, B_1^{y_{0:\tau}})^c}(x') \tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)} \pi'_{(k-1)\tau})(dx, dx') \\
& \quad \quad \times \left(\int_{\mathbb{R}^2} \tilde{Q} \pi_{(k-1)\tau}(du, du') \psi(y_{0:\tau}, u, u') \right) \lambda_W(dy_{0:\tau}) \\
& \quad \quad \quad \text{(using Equations (3.4), (4.40))} \\
& \leq e^{2h(\pi_0, \pi'_0)} \int_{y_{0:\tau} \in \mathcal{C}([0, \tau])} \mathbb{1}_{[0, \theta^{1-\iota} \Delta]} (|m_k(B_1^{y_{0:\tau}}) - m_{k-1}(B_1^{Y_{(k-2)\tau:(k-1)\tau}})|) \\
& \quad \times \int_{\mathbb{R}^2} \psi(y_{0:\tau}, x, x') \mathbb{1}_{C_k(\Delta, B_1^{y_{0:\tau}})^c}(x') \mathbb{1}_{C_{k-1}(\Delta)}(x) \tilde{Q} \pi_{k-1}(dx, dx') \lambda_W(dy_{0:\tau}) \\
& \quad \text{(using (4.32), (4.33), (4.38), (4.39) and the fact that } C_{k-1}(\Delta) \subset C_{k-1}(2\Delta)) \\
& \leq T(\Delta) e^{2h(\pi_0, \pi'_0)}.
\end{aligned}$$

We have, in the same way,

$$\begin{aligned}
& \mathbb{E} \left(\int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q} \pi'_{k-1}, \psi_k \rangle} \times \tilde{Q}(\mathbb{1}_{C_{k-1}(\Delta)^c} \pi'_{(k-1)\tau})(dx, dx') \Big| Y_{0:(k-1)\tau} \right) \\
& \leq e^{h(\pi_0, \pi'_0)} \int_{y_{0:\tau} \in \mathcal{C}([0, \tau])} \int_{\mathbb{R}^2} \psi(y_{0:\tau}, x, x') \mathbb{1}_{C_{k-1}(\Delta)^c}(x) \tilde{Q} \pi'_{(k-1)\tau}(dx, dx') \lambda_W(dy_{0:\tau}) \\
& \quad = e^{h(\pi_0, \pi'_0)} \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}})
\end{aligned}$$

and

$$\begin{aligned}
& \mathbb{E} \left(\int_{\mathbb{R}^2} \frac{\psi_k(x, x')}{\langle \tilde{Q} \pi'_{k-1}, \psi_k \rangle} \times \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}}) \xi_1(D_k, \Delta) \mathbb{1}_{C_k(\Delta, B_1^y)}(x') dx dx' \Big| Y_{0:(k-1)\tau} \right) \\
& \leq e^{h(\pi_0, \pi'_0)} \int_{y_{0:\tau} \in \mathcal{C}([0, \tau])} \int_{\mathbb{R}^2} \psi(y_{0:\tau}, x, x') \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}}) \xi_1(m_k(B_1^{y_{0:\tau}}) - m_{k-1}, \Delta) \\
& \quad \quad \times \mathbb{1}_{C_k(\Delta, B_1^y)}(x') dx dx' \lambda_W(dy_{0:\tau}) \\
& \leq e^{h(\pi_0, \pi'_0)} \pi'_{(k-1)\tau}(C_{k-1}(\Delta)^{\mathfrak{G}}).
\end{aligned}$$

So, using Equation (4.37) and Corollary 4.6, we get the result. \square

5 New formula for the truncated filter

We have reduced the problem to a discrete-time problem. For all n , $\pi_{n\tau}$ is the marginal of a Feynman-Kac sequence based on the transition \tilde{Q} and the potentials $(\psi_k)_{k \geq 1}$ (see Equations (3.6), (3.7), (3.8), Section 3.1 for the definition of \tilde{Q} , Section 3.2 for the definition of a Feynman-Kac sequence). As in [OR05], we restrict the state space to the compacts $(C_k(\Delta))_{k \geq 0}$. If \tilde{Q} restricted

to compacts was mixing, then, due to Proposition 3.1 (and to the remark below Equation (3.10)), $\pi_{n\tau}^\Delta$ could be viewed as the law at time n of a Markov chain with contracting Markov kernels ; and so Lemma 1.4 would be relatively easy to prove. By construction, \tilde{Q} restricted to some compacts cannot be mixing. This is an effect of the fact that the observations are continuous in time.

The purpose of this Section is to find another representation of the sequence $(\pi_{n\tau})_{n \geq 0}$ as a Feynman-Kac sequence, in such a way that the underlying Markov operators would be mixing, when restricted to compacts. Looking at Equation (3.9), we see that a Feynman-Kac sequence is a result of the deformation of a measure on trajectories (we weight the trajectories with potentials $(\psi_k)_{k \geq 1}$). The main idea of the following is to incorporate the deformations delicately (in two steps), in order to retain something of the mixing property of the operator Q (which is mixing when restricted to compacts).

In this Section, we work with a fixed observation $(Y_s)_{s \geq 0} = (y_s)_{s \geq 0}$.

5.1 Filter based on partial information

We define, for all $x = (x_1, x_2)$, $x' = (x'_1, x'_2)$ in \mathbb{R}^2 , k in \mathbb{N}^* , n in \mathbb{N}^* , $n \geq k$, (recall that ψ_k^Δ is defined in Equation (4.24) and that ξ_1, ξ_2 are defined in (4.26), (4.27))

$$\text{for } k \geq 2, \tilde{R}_k^\Delta(x, dx') = \begin{cases} \mathbb{1}_{C_{k-1}(\Delta)}(x'_1) \psi_k^\Delta(x') \tilde{Q}^2(x, dx') & \text{if } x_2 \in C_{k-2}(\Delta), \\ \mathbb{1}_{C_{k-1}(\Delta)}(x'_1) \psi_k^\Delta(x') \xi_1(D_{k-1}, \Delta) dx' & \text{otherwise,} \end{cases} \quad (5.1)$$

$$\psi_{2n|2k}^\Delta(x) = \begin{cases} \tilde{R}_{2n}^\Delta \tilde{R}_{2n-2}^\Delta \dots \tilde{R}_{2k+2}^\Delta(x, \mathbb{R}^2) & \text{if } k \leq n-1, \\ 1 & \text{if } k = n, \end{cases} \quad (5.2)$$

(so $\psi_{2n|2k}^\Delta(x)$ does not depend on x_1),

$$S_{2n|2k}^\Delta(x, dx') = \begin{cases} \frac{\psi_{2n|2k+2}^\Delta(x')}{\psi_{2n|2k}^\Delta(x)} \tilde{R}_{2k+2}^\Delta(x, dx') & \text{if } k \leq n-1, \\ dx' & \text{if } k = n-1. \end{cases}$$

These notations come from [DG01]. As Q has a density with respect to the Lebesgue measure on \mathbb{R} , so has $S_{2n|2k}^\Delta$ (with respect to the Lebesgue measure on \mathbb{R}^2). We write $(x, x') \in E^2 \mapsto S_{2n|2k}^\Delta(x, x')$ for this density. We fix n in \mathbb{N}^* in the rest of this subsection and in the following subsection. We define $S_{2n|2k}^{\Delta, (p)}$, $\psi_{2n|2k}^{\Delta, (p)}$, for $0 \leq k \leq n$, with the same formulas used above to define $S_{2n|2k}^\Delta$, $\psi_{2n|2k}^\Delta$, except we replace ψ_{2n}^Δ by 1. For all $D > 0$, we set

$$\epsilon(D, \Delta) = \frac{\xi_1(D, \Delta)}{\xi_2(D, \Delta)}, \quad (5.3)$$

and, for all k ,

$$\epsilon_k = \epsilon(D_k, \Delta).$$

Lemma 5.1. *For $k \leq n-1$, $S_{2n|2k}^\Delta$ is a Markov operator and $S_{2n|2k}^\Delta$ is $(1 - \epsilon_{2k+1}^2)$ -contracting for the total variation norm, $S_{2n|2k}^{\Delta, (p)}$ is a Markov operator and $S_{2n|2k}^{\Delta, (p)}$ is $(1 - \epsilon_{2k+1}^2)$ -contracting for the total variation norm*

Proof. We write the proof only for the kernels S_{\dots}^Δ , it would be very similar for the kernels $S_{\dots}^{\Delta, (p)}$. By Proposition 3.1, $S_{2n|2k}^\Delta$ is a Markov operator. We set, for all $k \geq 1$, x_1, x_2 in \mathbb{R} ,

$$\lambda_k(dx_1, dx_2) = \mathbb{1}_{C_{k-1}(\Delta)}(x_1) \mathbb{1}_{C_k(\Delta)}(x_2) \psi_k(x_1, x_2) Q(x_1, x_2) dx_1 dx_2.$$

By Lemma 2.1, we have, for all x_1, x_2, z_1, z_2 in \mathbb{R} , $k \geq 2$ (we use here the second line of Equation (5.1))

$$\xi_1(D_{k-1}, \Delta) \lambda_k(dz_1, dz_2) \leq \tilde{R}_k^\Delta(x_1, x_2, dz_1, dz_2) \leq \xi_2(D_{k-1}, \Delta) \lambda_k(dz_1, dz_2). \quad (5.4)$$

So \tilde{R}_k^Δ is $\sqrt{\epsilon_{k-1}}$ -mixing. So, for all x in \mathbb{R}^2 , all k such that $0 \leq k \leq n-1$ (the convention being that, if $k = n-1$, $(\tilde{R}_{2n}^\Delta \dots \tilde{R}_{2k+4}^\Delta)(y, dz) = \delta_y(dz)$)

$$\begin{aligned} \psi_{2n|2k}^\Delta(x) &= \int_{\mathbb{R}^2 \times \mathbb{R}^2} (\tilde{R}_{2n}^\Delta \dots \tilde{R}_{2k+4}^\Delta)(y, dz) \tilde{R}_{2k+2}^\Delta(x, dy) \\ &\leq \int_{\mathbb{R}^2 \times \mathbb{R}^2} (\tilde{R}_{2n}^\Delta \dots \tilde{R}_{2k+4}^\Delta)(y, dz) \xi_2(D_{2k+1}, \Delta) \lambda_{2k+2}(dy), \end{aligned}$$

and, for x' in \mathbb{R}^2 ,

$$\tilde{R}_{2k+2}^\Delta(x, dx') \geq \xi_1(D_{2k+1}, \Delta) \lambda_{2k+2}(dx'),$$

so

$$S_{2n|2k}^\Delta(x, dx') \geq \frac{\xi_1(D_{2k+1}, \Delta)}{\xi_2(D_{2k+1}, \Delta)} \times \frac{\tilde{R}_{2k+4:2n}^\Delta(x', \mathbb{R}^2) \lambda_{2k+2}(dx')}{\int_{\mathbb{R}^2} \tilde{R}_{2k+4:2n}^\Delta(y, \mathbb{R}^2) \lambda_{2k+2}(dy)}. \quad (5.5)$$

In the same way as above, we can also obtain

$$S_{2n|2k}^\Delta(x, dx') \leq \frac{\xi_2(D_{2k+1}, \Delta)}{\xi_1(D_{2k+1}, \Delta)} \times \frac{\tilde{R}_{2k+4:2n}^\Delta(x', \mathbb{R}^2) \lambda_{2k+2}(dx')}{\int_{\mathbb{R}^2} \tilde{R}_{2k+4:2n}^\Delta(y, \mathbb{R}^2) \lambda_{2k+2}(dy)}. \quad (5.6)$$

This implies that $S_{2n|2k}^\Delta$ is $(1 - \epsilon_{2k+1}^2)$ -contracting for the total variation norm (see Subsection 3.1). One can also use Proposition 3.1 to prove this result. We did it this way because we will re-use Equations (5.5), (5.6). \square

We set Z_0 to be of the form $Z_0 = (0, Z_0^{(2)})$, with $Z_0^{(2)}$ a random variable. We set $(Z_{2k})_{0 \leq k \leq n}$ to be a non-homogeneous Markov chain with kernels $S_{2n|0}^\Delta, S_{2n|2}^\Delta, \dots, S_{2n|2n-2}^\Delta$ (for k in $\{1, 2, \dots, n\}$), the law of Z_{2k} knowing Z_{2k-2} is $S_{2n|2k-2}^\Delta(Z_{2k-2}, \cdot)$. For Z_{2k} being a element of this chain, we denote by $Z_{2k}^{(1)}$ and $Z_{2k}^{(2)}$ its first and second component respectively. Recalling Proposition 3.1 (or Proposition 3.1, p. 428 in [OR05], or similar results in [DG01]), if the law of Z_0 is chosen properly, then $Z_{2n}^{(2)}$ has the same law as X_{2n} knowing $Y_{\tau:2\tau}, \dots, Y_{(2n-1)\tau:2n\tau}$, hence the title of this Subsection.

Remark 5.2. We have that, for all $k \geq 1$, $Z_{2k}^{(2)}$ takes values in $C_{2k}(\Delta)$ and $Z_{2k}^{(1)}$ takes values in $C_{2k-1}(\Delta)$.

We set $(Z_{2k}^{(p)})_{0 \leq k \leq n}$ to be a non-homogeneous Markov chain with $Z_0^{(p)} = Z_0$ and with kernel $S_{2n|0}^{\Delta, (p)}, S_{2n|2}^{\Delta, (p)}, \dots, S_{2n|2n-2}^{\Delta, (p)}$.

We set $U_{2k+1} = (Z_{2k}^{(2)}, Z_{2k+2}^{(1)})$ for $k \in \{0, 1, \dots, n-1\}$ and $U_{2n+1} = Z_{2n}^{(2)}$. We set $U_{2k+1}^{(p)} = (Z_{2k}^{(p)(2)}, Z_{2k+2}^{(p)(1)})$ for $k \in \{0, 1, 2, \dots, n-1\}$.

5.2 New Markov chain

Lemma 5.3. *The sequence $(U_1, U_3, \dots, U_{2n-1}, U_{2n+1}^{(1)})$ is a non-homogeneous Markov chain. The sequence $(U_1^{(p)}, U_3^{(p)}, \dots, U_{2n-3}^{(p)}, U_{2n-1}^{(p)})$ is a non-homogeneous Markov chain.*

If $Z_0^{(2)}$ is of law μ , then the law of U_1 is given by, for all (z, z') in \mathbb{R}^2 ,

$$\mathbb{P}(U_1 \in (dz, dz')) = \int_{x \in \mathbb{R}} S_{2n|0}^\Delta((0, z), (dz', dx)) \mu(dz). \quad (5.7)$$

We write $S_{2n|2k+1}^U$ for the transition kernel between U_{2k-1} and U_{2k+1} (for $k = 1, 2, \dots, n-1$) and $S_{2n|2n+1}^U$ for the transition between U_{2n-1} and $U_{2n+1}^{(1)}$. We write $S_{2n|2k+1}^{(p)U}$ for the transition kernel between $U_{2n|2k-1}^{(p)}$ and $U_{2n|2k+1}^{(p)}$ (for $k = 1, 2, \dots, n-1$)

Proof. We write the proof only for $(U_1, U_3, \dots, U_{2n-1}, U_{2n+1}^{(1)})$, it would be very similar for the sequence $(U_1^{(p)}, U_3^{(p)}, \dots, U_{2n-1}^{(p)})$. Let φ be a test function (in $\mathcal{C}_b^+(\mathbb{R}^2)$). For $k \in \{1, \dots, n-1\}$, we have (for $z_0^{(2)}, z_2^{(1)}, \dots, z_{2k}^{(1)}$ in \mathbb{R})

$$\begin{aligned} \mathbb{E}(\varphi(U_{2k+1})|U_1 = (z_0^{(2)}, z_2^{(1)}), \dots, U_{2k-1} = (z_{2k-2}^{(2)}, z_{2k}^{(1)})) &= \\ \mathbb{E}(\varphi(Z_{2k}^{(2)}, Z_{2k+2}^{(1)})|Z_0^{(2)} = z_0^{(2)}, \dots, Z_{2k-2}^{(2)} = z_{2k-2}^{(2)}, Z_{2k}^{(1)} = z_{2k}^{(1)}) &= \\ \mathbb{E}(\varphi(Z_{2k}^{(2)}, Z_{2k+2}^{(1)})|Z_{2k-2}^{(2)} = z_{2k-2}^{(2)}, Z_{2k}^{(1)} = z_{2k}^{(1)}), & \end{aligned}$$

as $S_{2n|2k-2}^\Delta(z_{2k-2}^{(1)}, z_{2k-2}^{(2)}, \dots)$ does not depend on $z_{2k-2}^{(1)}$. So the quantity above is equal to, for any $z \in C_{2k-1}(\Delta)$,

$$\begin{aligned} \int_{\mathbb{R}^2} \varphi(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \frac{S_{2n|2k-2}^\Delta((z, z_{2k-2}^{(2)}), (z_{2k}^{(1)}, z_{2k}^{(2)}))}{\int_{\mathbb{R}} S_{2n|2k-2}^\Delta((z, z_{2k-2}^{(2)}), (z_{2k}^{(1)}, z')) dz'} \\ \times \left(\int_{\mathbb{R}} S_{2n|2k}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), (z_{2k+2}^{(1)}, z_{2k+2}^{(2)})) dz_{2k+2}^{(2)} \right) dz_{2k}^{(2)} dz_{2k+2}^{(1)}. \end{aligned}$$

A similar computation can be made for $\mathbb{E}(\varphi(U_{2n+1}^{(1)})|U_1, \dots, U_{2n-1})$. \square

We set, for all k ,

$$\begin{aligned} \epsilon'(D, \Delta) &= \exp \left[- \left(\frac{B_2 p_{2,1}^2}{2} + \frac{1}{4\tau} \right) \left(\frac{\Delta}{B_2(1+p_{2,1})} + D \right)^2 \right. \\ &\quad \left. - (\Delta p_{2,1} + 3M) \left(\frac{\Delta}{B_2(1+p_{2,1})} + D \right) - 3\tau \left(\frac{M}{2} + \frac{M^2}{4} \right) \right], \quad (5.8) \\ \epsilon'_k &= \epsilon'(|m_k - m_{k-1}|, \Delta). \end{aligned}$$

Proposition 5.4. *For any $k = 1, 2, \dots, n$, the Markov kernel $S_{2n|2k+1}^U$ is $(\epsilon_{2k-1}^2(\epsilon'_{2k})^2)$ -contracting. For any $k = 1, 2, \dots, n-1$, the Markov kernel $S_{2n|2k+1}^{(p)U}$ is $(\epsilon_{2k-1}^2(\epsilon'_{2k})^2)$ -contracting.*

Before going into the proof of the above Proposition, we need the following technical result. We are interested in the bounds appearing in Lemma 4.4. We suppose that t_1, t_2, x, z in \mathbb{R} are fixed. To simplify the computations, we introduce the following notations:

$$\begin{aligned} \begin{pmatrix} t'_1 \\ t'_2 \end{pmatrix} &= (P^{-1})^T \begin{pmatrix} t_1 \\ t_2 \end{pmatrix}, \\ M_1 &= \frac{2M(|1+p_{2,1}|)}{p_{1,1}}. \end{aligned}$$

Lemma 5.5. *Suppose that, for some $D \geq 0$,*

$$|x - z| \leq D,$$

$$|2B_2(p_{1,2} + 1)z - t'_2| \leq \Delta$$

then

$$\begin{aligned} \exp \left(- \frac{1}{4B_2} (t'_2 - 2B_2(p_{2,1} + 1)z)^2 - B_2 p_{2,1}^2 D^2 - p_{2,1} D \Delta \right) \\ \leq \exp \left(- \frac{1}{4B_2} (t'_2 - 2B_2(p_{2,1}x + z))^2 \right) \\ \leq \exp \left(- \frac{1}{4B_2} (t'_2 - 2B_2(p_{2,1} + 1)z)^2 + p_{2,1} D \Delta \right). \end{aligned}$$

Proof. We have

$$\begin{aligned}
& \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1}x + z))^2\right) \\
&= \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1} + 1)z) + 2B_2p_{2,1}(z - x))^2\right) \\
&\leq \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1} + 1)z)^2 - B_2p_{2,1}^2(z - x)^2 + |p_{2,1}(z - x)| \times |t'_2 - 2B_2(p_{2,1} + 1)z|\right) \\
&\leq \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1} + 1)z)^2 + p_{2,1}D\Delta\right)
\end{aligned}$$

and

$$\begin{aligned}
& \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1}x + z))^2\right) \\
&\geq \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1} + 1)z)^2 - B_2p_{2,1}^2(z - x)^2 - |p_{2,1}(z - x)| \times |t'_2 - 2B_2(p_{2,1} + 1)z|\right) \\
&\geq \exp\left(-\frac{1}{4B_2}(t'_2 - 2B_2(p_{2,1} + 1)z)^2 - B_2p_{2,1}^2D^2 - p_{2,1}D\Delta\right)
\end{aligned}$$

□

Proof of Proposition 5.4. We write the proof in the case $k \in \{1, 2, \dots, n-1\}$ and for $S_{2n|2k+1}^U$ (the other cases being very similar). Let φ be a test function (in $C_b^+(\mathbb{R})$). By Remark 5.2, we have that $U_{2k-1}^{(2)}$ takes its values in $C_{2k-1}(\Delta)$. We write, for any $z_{2k-2}^{(2)} \in \mathbb{R}$, $z_{2k}^{(1)} \in C_{2k-1}(\Delta)$, $z \in \mathbb{R}$ (like in the proof of Lemma 5.3)

$$\begin{aligned}
\mathbb{E}(\varphi(U_{2k+1})|U_{2k-1} = (z_{2k-2}^{(2)}, z_{2k}^{(1)})) &= \\
& \int_{\mathbb{R}^2} \varphi(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \frac{S_{2n|2k-2}^\Delta((z, z_{2k-2}^{(2)}), (z_{2k}^{(1)}, z_{2k}^{(2)}))}{\int_{\mathbb{R}} S_{2n|2k-2}^\Delta((z, z_{2k-2}^{(2)}), (z_{2k}^{(1)}, z')) dz'} \\
& \times \left(\int_{\mathbb{R}} S_{2n|2k}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), (z_{2k+2}^{(1)}, z_{2k+2}^{(2)})) dz_{2k+2}^{(2)} \right) dz_{2k}^{(2)} dz_{2k+2}^{(1)} \geq \\
& \quad \text{(by Equations (5.5), (5.6))} \\
& \int_{\mathbb{R}^2} \varphi(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \epsilon_{2k-1}^2 \\
& \times \frac{\tilde{R}_{2n:2k+2}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), \mathbb{R}^2) \mathbb{1}_{C_{2k-1}(\Delta)}(z_{2k}^{(1)}) \mathbb{1}_{C_{2k}(\Delta)}(z_{2k}^{(2)}) \psi_{2k}(z_{2k}^{(1)}, z_{2k}^{(2)}) Q(z_{2k}^{(1)}, z_{2k}^{(2)})}{\int_{\mathbb{R}} \tilde{R}_{2n:2k+2}^\Delta((z_{2k}^{(1)}, z'), \mathbb{R}^2) \mathbb{1}_{C_{2k-1}(\Delta)}(z_{2k}^{(1)}) \mathbb{1}_{C_{2k}(\Delta)}(z') \psi_{2k}(z_{2k}^{(1)}, z') Q(z_{2k}^{(1)}, z') dz'} \\
& \times \left(\int_{\mathbb{R}} S_{2n|2k}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), (z_{2k+2}^{(1)}, z_{2k+2}^{(2)})) dz_{2k+2}^{(2)} \right) dz_{2k}^{(2)} dz_{2k+2}^{(1)} \quad (5.9)
\end{aligned}$$

From Lemma 2.2 and using the same kind of computation as in the proof of Lemma 4.4 and Equation (4.15), we get, for all z_{2k} such that $z_{2k}^{(2)} \in C_{2k}(\Delta)$,

$$\begin{aligned}
& \psi_{2k}(z_{2k}^{(1)}, z_{2k}^{(2)}) \geq \sigma_1 \sigma_2 e^{-2M|z_{2k}^{(1)} - z_{2k}^{(2)}| - \tau \left(M + \frac{M^2}{2}\right)} \\
& \times \exp\left(-A_2(z_{2k}^{(1)})^2 - B_2(z_{2k}^{(2)})^2 + C_1 z_{2k}^{(1)} z_{2k}^{(2)} + A_1^{y(2k-1)\tau:(2k)\tau} z_{2k}^{(1)} + B_1^{y(2k-1)\tau:(2k)\tau} z_{2k}^{(2)} + C_0^{y(2k-1)\tau:(2k)\tau}\right) \\
& = \sigma_1 \sigma_2 e^{-2M|z_{2k}^{(1)} - z_{2k}^{(2)}| - \tau \left(M + \frac{M^2}{2}\right)} \times \exp\left[-\frac{1}{4A_2} \left(\frac{A_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - \frac{p_{2,1} B_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - 2A_2 p_{1,1} z_{2k}^{(1)}\right)^2\right]
\end{aligned}$$

$$\begin{aligned}
& - \frac{1}{4B_2} \left(B_1^{y(2k-1)\tau:(2k)\tau} - 2B_2(p_{2,1}z_{2k}^{(1)} + z_{2k}^{(2)}) \right)^2 + C_0^{y(2k-1)\tau:(2k)\tau} \\
& + \frac{1}{4} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right) \kappa^{-1} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right)^T \Big] \\
& \geq (\text{by Lemma 5.5}) \sigma_1 \sigma_2 e^{-2M|z_{2k}^{(1)} - z_{2k}^{(2)}| + C_0^{y(2k-1)\tau:(2k)\tau}} \times e^{-\tau \left(M + \frac{M^2}{2} \right)} \\
& \times \exp \left[- \frac{1}{4A_2} \left(\frac{A_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - \frac{p_{2,1}B_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - 2A_2 p_{1,1} z_{2k}^{(1)} \right)^2 \right. \\
& - \frac{1}{4B_2} \left(B_1^{y(2k-1)\tau:(2k)\tau} - 2B_2(p_{2,1} + 1)z_{2k}^{(2)} \right)^2 - B_2 p_{2,1}^2 (z_{2k}^{(1)} - z_{2k}^{(2)})^2 - \Delta p_{2,1} |z_{2k}^{(1)} - z_{2k}^{(2)}| \\
& \left. + \frac{1}{4} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right) \kappa^{-1} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right)^T \right].
\end{aligned}$$

We set

$$\begin{aligned}
\psi_{2k}^{(1)}(x) &= \exp \left(- \frac{1}{4A_2} \left(\frac{A_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - \frac{p_{2,1}B_1^{y(2k-1)\tau:(2k)\tau}}{p_{1,1}} - 2A_2 p_{1,1} x \right)^2 \right), \\
\psi_{2k}^{(2)}(x) &= \exp \left(- \frac{1}{4B_2} \left(B_1^{y(2k-1)\tau:(2k)\tau} - 2B_2(p_{2,1} + 1)x \right)^2 \right)
\end{aligned}$$

In the same way as above:

$$\begin{aligned}
\psi_{2k}(z_{2k}^{(1)}, z_{2k}^{(2)}) &\leq \sigma_1 \sigma_2 e^{2M|z_{2k}^{(1)} - z_{2k}^{(2)}| + \tau \left(M + \frac{M^2}{2} \right) + C_0^{y(2k-1)\tau:(2k)\tau}} \psi_{2k}^{(1)}(z_{2k}^{(1)}) \psi_{2k}^{(2)}(z_{2k}^{(2)}) \\
&\times \exp \left(\Delta p_{2,1} |z_{2k}^{(1)} - z_{2k}^{(2)}| + \frac{1}{4} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right) \kappa^{-1} \left(A_1^{y(2k-1)\tau:(2k)\tau}, B_1^{y(2k-1)\tau:(2k)\tau} \right)^T \right).
\end{aligned}$$

From Lemma 2.1, we get for all $z_{2k}^{(1)} \in C_{2k-1}(\Delta)$, $z_{2k}^{(2)} \in C_{2k}(\Delta)$,

$$\begin{aligned}
\frac{1}{\sqrt{2\pi\tau}} \exp \left(- \frac{1}{2\tau} \left(\frac{\Delta}{B_2(1+p_{2,1})} + D_{2k} \right)^2 - M \left(\frac{\Delta}{B_2(1+p_{2,1})} + D_{2k} \right) - \tau \left(\frac{M}{2} + \frac{M^2}{2} \right) \right) \\
\leq Q(z_{2k}^{(1)}, z_{2k}^{(2)}) \leq \frac{1}{\sqrt{2\pi\tau}} \exp \left(M \left(\frac{\Delta}{B_2(1+p_{2,1})} + D_{2k} \right) + \frac{M\tau}{2} \right)
\end{aligned}$$

Looking back at (5.9), we get

$$\begin{aligned}
\mathbb{E}(\varphi(U_{2k+1}) | U_{2k-1} = (z_{2k-2}^{(2)}, z_{2k}^{(1)})) &\geq \\
\int_{\mathbb{R}^2} \varphi(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \epsilon_{2k-1}^2 (\epsilon'_{2k})^2 &\frac{\tilde{R}_{2n:2k+2}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), \mathbb{R}^2) \mathbb{1}_{C_{2k-1}(\Delta)}(z_{2k}^{(1)}) \mathbb{1}_{C_{2k}(\Delta)}(z_{2k}^{(2)}) \psi_{2k}^{(2)}(z_{2k}^{(2)})}{\int_{\mathbb{R}} \tilde{R}_{2n:2k+2}^\Delta((z_{2k}^{(1)}, z'), \mathbb{R}^2) \mathbb{1}_{C_{2k-1}(\Delta)}(z_{2k}^{(1)}) \mathbb{1}_{C_{2k}(\Delta)}(z_{2k}^{(2)}) \psi_{2k}^{(2)}(z') dz'} \\
&\times \left(\int_{\mathbb{R}} S_{2n|2k}^\Delta((z_{2k}^{(1)}, z_{2k}^{(2)}), (z_{2k+2}^{(1)}, z_{2k+2}^{(2)})) dz_{2k+2}^{(2)} \right) dz_{2k}^{(2)} dz_{2k+2}^{(1)}.
\end{aligned}$$

As $R_{2n:2k+2}((z_{2k}^{(1)}, z'), \cdot)$ and $S_{2n|2k}^\Delta((z_{2k}^{(1)}, z'), \cdot)$ do not depend on $z_{2k}^{(1)}$ for any z' , we get that $S_{2n|2k+1}^U$ is $(1 - \epsilon_{2k-1}^2 (\epsilon'_{2k})^2)$ -contracting (remember Section 3.1). \square

5.3 New representation

Proposition 5.6. *Let $n \geq 1$. If we suppose that $Z_0^{(2)}$ is of law $\psi_{2n|0}^\Delta(0, \cdot) \bullet \mu$, then, for all test function φ (in $C_b^+(\mathbb{R})$),*

$$\frac{\mathbb{E}(\varphi(U_{2n+1}^{(1)}) \prod_{1 \leq i \leq n} \psi_{2i-1}^\Delta(U_{2i-1}))}{\mathbb{E}(\prod_{1 \leq i \leq n} \psi_{2i-1}^\Delta(U_{2i-1}))} = \left(\bar{R}_{2n}^\Delta \bar{R}_{2n-1}^\Delta \dots \bar{R}_1^\Delta(\mu) \right) (\varphi). \quad (5.10)$$

If we suppose that $Z_0^{(2)}$ is of law $\psi_{2n|0}^{\Delta,(p)}(0, \cdot) \bullet \mu$, then, for all test function φ (in $\mathcal{C}_b^+(\mathbb{R})$),

$$\frac{\mathbb{E}(\varphi(U_{2n-1}^{(p)(2)}) \prod_{1 \leq i \leq n} \psi_{2i-1}^{\Delta}(U_{2i-1}^{(p)}))}{\mathbb{E}(\prod_{1 \leq i \leq n} \psi_{2i-1}^{\Delta}(U_{2i-1}^{(p)}))} = \left(\overline{R}_{2n-1}^{\Delta} \overline{R}_{2n-2}^{\Delta} \dots \overline{R}_1^{\Delta}(\mu) \right) (\varphi). \quad (5.11)$$

Remark 5.7. Recall that we are working with a fixed observation $(Y_s)_{s \geq 0} = (y_s)_{s \geq 0}$. The above Proposition tells that, for all n , $\overline{R}_n^{\Delta} \overline{R}_{n-1}^{\Delta} \dots \overline{R}_1^{\Delta}(\mu)$ can be written as the n -th term of a Feynman-Kac sequence based on mixing kernels (by Proposition 5.4). We can apply Proposition 3.1 to this Feynman-Kac sequence. This representation and this result are also true for a measure $\overline{R}_n^{\Delta} \overline{R}_{n-1}^{\Delta} \dots \overline{R}_k^{\Delta}(\eta)$ for any $k \leq n$, η probability measure on \mathbb{R} .

Proof. We write the proof only for Equation (5.10). The computation leading to Equation (5.11) would be very similar. It would simplify nicely because we replace ψ_{2n}^{Δ} by 1 in the definition of the $S_{2n|\dots}^{\Delta(p)}$, $\psi_{2n|\dots}^{\Delta(p)}$.

We have, for any test function φ (in $\mathcal{C}_b^+(\mathbb{R})$),

$$\begin{aligned} \mathbb{E}(\varphi(U_{2n+1}^{(1)}) \prod_{1 \leq i \leq n} \psi_{2i-1}^{\Delta}(U_{2i-1})) &= \\ &\int_{\mathbb{R} \times (\mathbb{R}^2)^n} \varphi(z_{2n}^{(2)}) \prod_{0 \leq k \leq n-1} \left[S_{2n|2k}^{\Delta}(z_{2k}, z_{2k+2}) \psi_{2k+1}^{\Delta}(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \right] \\ &\quad \delta_0(dz_0^{(1)}) (\psi_{2n|0}^{\Delta}(0, \cdot) \bullet \mu)(dz_0^{(2)}) dz_2 \dots dz_{2n} = \\ &\int_{\mathbb{R} \times (\mathbb{R}^2)^n} \varphi(z_{2n}^{(2)}) \prod_{0 \leq k \leq n-1} \left[\frac{\psi_{2n|2k+2}^{\Delta}(z_{2k+2})}{\psi_{2n|2k}^{\Delta}(z_{2k})} \widetilde{R}_{2k+2}^{\Delta}(z_{2k}, dz_{2k+2}) \psi_{2k+1}^{\Delta}(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \right] \\ &\quad \times \delta_0(dz_0^{(1)}) (\psi_{2n|0}^{\Delta}(0, \cdot) \bullet \mu)(dz_0^{(2)}) dz_2 \dots dz_{2n} = \\ &\int_{\mathbb{R} \times (\mathbb{R}^2)^n} \varphi(z_{2n}^{(2)}) \prod_{0 \leq k \leq n-1} \left[\widetilde{R}_{2k+2}^{\Delta}(z_{2k}, dz_{2k+2}) \psi_{2k+1}^{\Delta}(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \right] \\ &\quad \times \frac{1}{\mu(\psi_{2n|0}^{\Delta}(0, \cdot))} \delta_0(dz_0^{(1)}) \mu(dz_0^{(2)}) dz_2 \dots dz_{2n} = \\ &\int_{\mathbb{R} \times (\mathbb{R}^2)^n} \varphi(z_{2n}^{(2)}) \prod_{0 \leq k \leq n-1} \left[\psi_{2k+2}^{\Delta}(z_{2k+2}^{(1)}, z_{2k+2}^{(2)}) \psi_{2k+1}^{\Delta}(z_{2k}^{(2)}, z_{2k+2}^{(1)}) \widetilde{Q}^2(z_{2k}, dz_{2k+2}) \right] \\ &\quad \times \frac{1}{\mu(\psi_{2n|0}^{\Delta}(0, \cdot))} \delta_0(dz_0^{(1)}) \mu(dz_0^{(2)}) dz_2 \dots dz_{2n}, \end{aligned}$$

which proves the desired result (recall Equation (3.11)). \square

6 Stability results

In this section, the observations are non longer fixed.

6.1 Stability of the truncated filter

We show here that a product of coefficients τ decays geometrically in expectation (see the Lemma below). These coefficients are the contraction coefficients of the operators S_t^U , $S_t^{U,(p)}$, which are related to the truncated filter through Proposition 5.6. This is why we say that the result below means the stability of the truncated filter.

We set, for all t in \mathbb{R} , $k \geq 1$,

$$\tau(t, \Delta) = 1 - (\epsilon'(t, \Delta) \epsilon(t, \Delta))^2,$$

$$\tau_k = 1 - (\epsilon'_k \epsilon_{k-1})^2.$$

We set, for $L > 0$,

$$\tilde{\alpha}(L) = \frac{192C\sqrt{\tau}}{L\sqrt{\pi}} \exp\left(-\frac{1}{2}\left(\frac{L}{12C\sqrt{2\tau}}\right)^2\right).$$

We fix $L > 0$ such that

$$L > 3|m_0| + 3CM\tau \text{ and } \tilde{\alpha}(L) \leq \frac{1}{4}. \quad (6.1)$$

We set

$$\rho = \frac{\tau(L, \Delta) + \sqrt{\tau(L, \Delta)^2 + 4\tilde{\alpha}(L)(1 - \tau(L, \Delta))}}{2}.$$

Lemma 6.1. *For $0 \leq k \leq n-1$, we have*

$$\begin{aligned} \mathbb{E}(\tau_{2n+1}\tau_{2n-1}\dots\tau_{2k+3}|\mathcal{F}_{0:(2k+1)\tau}) &\leq \left(1 - \frac{\epsilon(L, \Delta)^2\epsilon'(L, \Delta)^2}{2}\right)^{\lceil \frac{(n-k-2)_+}{2} \rceil}, \\ \mathbb{E}(\tau_{2n}\tau_{2n-2}\dots\tau_{2k+2}|\mathcal{F}_{0:2k\tau}) &\leq \left(1 - \frac{\epsilon(L, \Delta)^2\epsilon'(L, \Delta)^2}{2}\right)^{\lceil \frac{(n-k-2)_+}{2} \rceil}. \end{aligned}$$

Proof. We only write the proof of the second Equation above (the proof of the other equation is very similar). We take $L > 0$ and we set

$$\theta_{2k} = \begin{cases} \tau(L, \Delta) & \text{if } |m_{2k} - m_{2k-1}| < L \text{ and } |m_{2k-1} - m_{2k-2}| < L \\ 1 & \text{otherwise.} \end{cases}$$

For all k , we have $\tau_{2k} \leq \theta_{2k}$. For any $k \geq 1$, $|m_k - m_{k-1}|$ is a function of $Y_{(k-2)_+\tau:k\tau}$. So, for all k , θ_{2k} is a function of $Y_{(2k-3)_+\tau:2k\tau}$. We fix $k \geq 0$ and we define, for $n \geq 0$,

$$e_{2n|2k+2} = \begin{cases} \mathbb{E}(\theta_{2n}\theta_{2n-2}\dots\theta_{2k+2}|\mathcal{F}_{2k\tau}) & \text{if } k \leq n-1, \\ 1 & \text{otherwise.} \end{cases}$$

We suppose now that $n \geq k+2$. We then have

$$e_{2n|2k+2} = \mathbb{E}(\mathbb{E}(\theta_{2n}\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau})\theta_{2n-4}\dots\theta_{2k+2}|\mathcal{F}_{2k\tau})$$

and

$$\begin{aligned} &\mathbb{E}(\theta_{2n}\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau}) \\ &= \mathbb{E}(\theta_{2n-2}(1 - \mathbb{1}_{[0,L)}(D_{2n})\mathbb{1}_{[0,L)}(D_{2n-1})) + \tau(L, \Delta)\theta_{2n-2}\mathbb{1}_{[0,L)}(D_{2n})\mathbb{1}_{[0,L)}(D_{2n-1})|\mathcal{F}_{(2n-3)\tau}) \\ &= \mathbb{E}(\theta_{2n-2}\tau(L, \Delta) + (1 - \tau(L, \Delta))\theta_{2n-2}(1 - \mathbb{1}_{[0,L)}(D_{2n})\mathbb{1}_{[0,L)}(D_{2n-1}))|\mathcal{F}_{(2n-3)\tau}) \\ &\leq \tau(L, \Delta)\mathbb{E}(\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau}) + (1 - \tau(L, \Delta))[\mathbb{P}(|m_{2n} - m_{2n-1}| \geq L|\mathcal{F}_{(2n-3)\tau}) \\ &\quad + \mathbb{P}(|m_{2n-1} - m_{2n-2}| \geq L|\mathcal{F}_{(2n-3)\tau})]. \end{aligned}$$

Using Equation (4.21), we get

$$\begin{aligned} &\mathbb{E}(\theta_{2n}\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau}) \leq \tau(L, \Delta)\mathbb{E}(\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau}) \\ &+ (1 - \tau(L, \Delta))\left(\mathbb{P}\left(C\mathcal{V}_{(2n-2)\tau, 2n\tau} \geq \frac{L}{3}|\mathcal{F}_{(2n-3)\tau}\right) + \mathbb{P}\left(C\mathcal{W}_{(2n-2)\tau, 2n\tau}\left(1 + \frac{1}{\theta}\right) \geq \frac{L}{3}|\mathcal{F}_{(2n-3)\tau}\right)\right) \\ &+ (1 - \tau(L, \Delta))\left(\mathbb{P}\left(C\mathcal{V}_{(2n-3)\tau, (2n-1)\tau} \geq \frac{L}{3}|\mathcal{F}_{(2n-3)\tau}\right) + \mathbb{P}\left(C\mathcal{W}_{(2n-3)\tau, (2n-1)\tau}\left(1 + \frac{1}{\theta}\right) \geq \frac{L}{3}|\mathcal{F}_{(2n-3)\tau}\right)\right) \\ &\leq \tau(L, \Delta)\mathbb{E}(\theta_{2n-2}|\mathcal{F}_{(2n-3)\tau}) + 4(1 - \tau(L, \Delta))\mathbb{P}\left(C\mathcal{V}_{0, 2\tau} \geq \frac{L}{6}\right) \end{aligned}$$

(like in Equations (4.34), (4.35))

$$\begin{aligned} &\leq \tau(L, \Delta) \mathbb{E}(\theta_{2n-2} | \mathcal{F}_{(2n-3)\tau}) + 8(1 - \tau(L, \Delta)) \mathbb{P}\left(2C|W_{2\tau}| \geq \frac{L}{6}\right) \\ &\quad \text{(using Equation (4.36))} \\ &\leq \tau(L, \Delta) \mathbb{E}(\theta_{2n-2} | \mathcal{F}_{(2n-3)\tau}) + (1 - \tau(L, \Delta)) \tilde{\alpha}(L). \end{aligned}$$

The constant ρ is the positive root of the polynomial $X^2 - \tau(L, \Delta)X - (1 - \tau(L, \Delta))\tilde{\alpha}(L)$. So we have

$$1 > \rho = \tau(L, \Delta) + \frac{1}{\rho}(1 - \tau(L, \Delta))\tilde{\alpha}(L) \geq \tau(L, \Delta) + (1 - \tau(L, \Delta))\tilde{\alpha}(L).$$

So, we have

$$e_{2n|2k+2} \leq \tau(L, \Delta)e_{2n-2|2k+2} + (1 - \tau(L, \Delta))\tilde{\alpha}(L)e_{2n-4|2k+2} \leq \rho \times \sup(e_{2n-2|2k+2}, e_{2n-4|2k+2}).$$

Suppose now that k is fixed. We have

$$e_{2k+2|2k+2} \leq 1, \quad e_{2k+4|2k+2} \leq 1.$$

So, by recurrence,

$$e_{2n|2k+2} \leq \rho^{\lceil \frac{(n-k-2)_+}{2} \rceil}.$$

As $\tilde{\alpha}(L) \leq 1/4$, we have

$$\rho \leq \frac{1}{2}(\tau(L, \Delta) + \sqrt{\tau(L, \Delta)^2 + 1 - \tau(L, \Delta)}) \leq \frac{\tau(L, \Delta) + 1}{2} = 1 - \frac{(\epsilon(L, \Delta)^2 \epsilon'(L, \Delta))^2}{2}.$$

□

Proof of Lemma 1.4. We write the proof in the case where n and k are even. If k was even and n was odd, we would have to use the operators $S_{\dots}^{(p)U}$. If k was odd, the proof would be very similar but would require to introduce new and heavy notations.

By Proposition 5.6, Remark 5.7 and Equation (5.7), we have, for all μ in $\mathcal{P}(\mathbb{R})$ and all test function φ in $\mathcal{C}_b^+(\mathbb{R})$,

$$\begin{aligned} &(\overline{R}_n^\Delta \overline{R}_{n-1}^\Delta \dots \overline{R}_{k+1}^\Delta(\mu))(\varphi) \propto \int \varphi(u_{n+1}^{(1)}) \left(\prod_{i=1}^{(n-k)/2} \psi_{k+2i-1}^\Delta(u_{k+2i-1}) \right) \\ &\quad \times \left(\prod_{i=1}^{(n-k-2)/2} S_{n|k+2i+1}^U(u_{k+2i-1}, du_{k+2i+1}) \right) \\ &\quad \times S_{n|n+1}^U(u_{n-1}, du_{n+1}^{(1)}) \left(\int_{z' \in \mathbb{R}} S_{n|k}^\Delta((0, u_{k+1}^{(1)}), (du_{k+1}^{(2)}, dz')) (\psi_{n|k}^\Delta(0, \cdot) \bullet \mu)(du_{k+1}^{(1)}), \right) \end{aligned}$$

where we integrate over $u_{k+1}^{(1)} \in \mathbb{R}$, $u_{k+1}, u_{k+3}, \dots, u_{n-1} \in \mathbb{R}^2$, $u_{n+1}^{(1)} \in \mathbb{R}$.

By Proposition 5.4, we know that $S_{n|k+2i+1}^U$ is $(\epsilon_{k+2i-1}^2 (\epsilon'_{k+2i})^2)$ -mixing for all i in $\{1, 2, \dots, 1 + (n-k)/2\}$. We now apply Proposition 3.1 with the $S_{n|k+2i+1}^U$ playing the roles of the Ω_{\dots} and the ψ_{k+2i-1}^Δ playing the roles of the Ψ_{\dots} . By Equations (3.12), (3.13), we then have, for all μ and μ' in $\mathcal{P}(\mathbb{R})$,

$$\begin{aligned} &\left\| \overline{R}_n^\Delta \overline{R}_{n-1}^\Delta \dots \overline{R}_{k+1}^\Delta(\mu) - \overline{R}_n^\Delta \overline{R}_{n-1}^\Delta \dots \overline{R}_{k+1}^\Delta(\mu') \right\| \\ &\leq \prod_{i=1}^{(n-k)/2} (1 - \epsilon_{k+2i-1}^2 (\epsilon'_{k+2i})^2) \times 2 \inf \left(1, \frac{\|\psi_{n|k}^\Delta(0, \cdot) \bullet \mu - \psi_{n|k}^\Delta(0, \cdot) \bullet \mu'\|}{\epsilon_{k+1}^2 (\epsilon'_{k+2})^2} \right). \end{aligned}$$

By Equations (5.4), (3.12), (3.13), we have

$$\begin{aligned} \|\psi_{n|k}^\Delta(0, \cdot) \bullet \mu - \psi_{n|k}^\Delta(0, \cdot) \bullet \mu'\| &\leq 2 \inf \left(1, \frac{\|\psi_{n|k}^\Delta(0, \cdot)\|_\infty}{\langle \mu, \psi_{n|k}^\Delta(0, \cdot) \rangle} \|\mu - \mu'\| \right) \\ &\leq 2 \inf \left(1, \frac{\|\mu - \mu'\|}{\epsilon_{k+1}^2} \right). \end{aligned}$$

From which we get the result. \square

6.2 Approximation of the optimal filter by the truncated filter

We recall that “ \prec_{Δ} ” is defined in Definition 2.5.

Proof of Proposition 1.5. We write the proof only for Equation (1.4), the proof for Equation (1.5) being very similar. We have

$$\|\pi_{n\tau} - \pi_{n\tau}^\Delta\| \leq \|\pi_{n\tau} - \bar{R}_n^\Delta(\pi_{(n-1)\tau})\| + \sum_{1 \leq k \leq n-1} \|\bar{R}_{n:k+1}^\Delta(\pi_{k\tau}) - \bar{R}_{n:k+1}^\Delta(\bar{R}_k^\Delta(\pi_{(k-1)\tau}))\|. \quad (6.2)$$

Let us fix $k \in \{1, 2, \dots, n-1\}$. From Lemma 1.4, we get

$$\begin{aligned} &\mathbb{E}(\|\bar{R}_{n:k+1}^\Delta(\pi_{k\tau}) - \bar{R}_{n:k+1}^\Delta(\bar{R}_k^\Delta(\pi_{(k-1)\tau}))\|) \\ &\leq \mathbb{E} \left(\mathbb{E} \left(\prod_{3 \leq i \leq \lfloor \frac{n-k}{2} \rfloor} (1 - (\epsilon_{k+2i-1}'^2 (\epsilon_{k+2i}')^2)) \middle| \mathcal{F}_{(k+2)\tau} \right) \times 2 \inf \left(1, \frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|}{(\epsilon_{k+2}')^2 \epsilon_{k+1}^4} \right) \right), \end{aligned} \quad (6.3)$$

with the convention that a product over indexes in the null set is equal to one. From Lemma 6.1, we get

$$\begin{aligned} \mathbb{E}(\|\bar{R}_{n:k+1}^\Delta(\pi_{k\tau}) - \bar{R}_{n:k+1}^\Delta(\bar{R}_k^\Delta(\pi_{(k-1)\tau}))\|) &\leq \left(1 - \frac{(\epsilon(L, \Delta)^2 \epsilon'(L, \Delta))^2}{2} \right)^{\lfloor \frac{1}{2} (\lfloor \frac{n-k}{2} \rfloor - 4)_+ \rfloor} \\ &\quad \times 2 \mathbb{E} \left(\inf \left(1, \frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|}{(\epsilon_{k+2}')^2 \epsilon_{k+1}^4} \right) \right). \end{aligned} \quad (6.4)$$

As in [OR05], p. 434, we can bound

$$\inf \left(1, \frac{\|\pi_{n\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|}{(\epsilon_{k+2}')^2 \epsilon_{k+1}^4} \right) \leq \inf \left(1, \frac{T(\Delta)}{(\epsilon_{k+2}')^4 \epsilon_{k+1}^8} \right) + \inf \left(1, \frac{\|\pi_{n\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)} \right). \quad (6.5)$$

We have, if Δ satisfies the assumption of Proposition 4.7,

$$\begin{aligned} &\mathbb{E} \left(\inf \left(1, \frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)} \right) \right) \\ &= \mathbb{E} \left(\frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)} \mathbb{1}_{[0,1]} \left(\frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)} \right) \right) \\ &\quad + \mathbb{P} \left(\frac{\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)} > 1 \right) \\ &\leq \frac{2}{\sqrt{T(\Delta)}} \mathbb{E}(\|\pi_{k\tau} - \bar{R}_k^\Delta(\pi_{(k-1)\tau})\|) \end{aligned}$$

$$\text{(using Prop. 4.7)} \preceq \sqrt{T(\Delta)}. \quad (6.6)$$

We look now at the term $\inf(1, T(\Delta)(\epsilon'_{k+2})^{-4}\epsilon_{k+1}^{-8})$. Using Equations (4.4), (4.22), (4.23), (7.12), (7.17), (7.18), (7.20) and the remarks below Equation (4.4), we have, for all k ,

$$\begin{aligned} T(\Delta) \underset{\Delta, c}{\preceq} & \frac{h^{-1+\iota}\tau^{\iota-\frac{1}{2}}}{\Delta} \exp\left(-\frac{1}{2}\left(\frac{\tau^{\frac{1}{2}-\iota}h^{1-\iota}\Delta}{12C\sqrt{2}}\right)^2\right) \\ & + \left(M + \sqrt{h}\right) \frac{he^{26M^2\tau + \frac{\tau M}{2} + 40M^2/h}}{\Delta\theta^{3/2}} \exp\left(-\frac{1}{4B_2}d(\Delta)^2\right) C'_1(h, \tau) + e^{-C_0\Delta^2}, \end{aligned} \quad (6.7)$$

and using Equations (4.26), (4.27), (5.3),

$$\begin{aligned} \epsilon(D, \Delta)^{-1} &= \exp\left(\frac{\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right)^2}{2\tau} - \frac{\left(\left(D - \frac{\Delta}{B_2(1+p_{2,1})}\right)_+\right)^2}{2\tau}\right) e^{2M\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right) + \left(\tau + \frac{\tau^2}{2}\right)M} \\ &= \begin{cases} \exp\left(\frac{2D\Delta}{\tau B_2(1+p_{2,1})} + 2M\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right) + \left(\tau + \frac{\tau^2}{2}\right)M\right) & \text{if } D \geq \frac{\Delta}{B_2(1+p_{2,1})}, \\ \exp\left(\frac{\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right)^2}{2\tau} + 2M\left(D + \frac{\Delta}{B_2(1+p_{2,1})}\right) + \left(\tau + \frac{\tau^2}{2}\right)M\right) & \text{otherwise.} \end{cases} \end{aligned} \quad (6.8)$$

and using Equation (5.8),

$$\begin{aligned} (\epsilon'(D, \Delta))^{-2} &= \exp\left[\left(B_2p_{2,1}^2 + \frac{1}{2\tau}\right)\left(\frac{\Delta}{B_2(1+p_{2,1})} + D\right)^2\right. \\ &\quad \left.+ (2\Delta p_{2,1} + 6M)\left(\frac{\Delta}{B_2(1+p_{2,1})} + D\right) + 3\tau\left(M + \frac{M^2}{2}\right)\right]. \end{aligned} \quad (6.9)$$

We note that the above expressions are nondecreasing functions of D . From Equation (4.21), we get, for $j = k + 1, k + 2$.

$$D_j \leq C(\tau M + \mathcal{V}_{(k-1)\tau, (k+2)\tau} + 2\mathcal{W}_{(k-1)\tau, (k+2)\tau}). \quad (6.10)$$

The variables $\mathcal{V}_{(k-1)\tau, (k+2)\tau}$ and $\mathcal{W}_{(k-1)\tau, (k+2)\tau}$ are independent and can be controlled as in Equations (4.34), (4.35). So we can bound

$$\forall x \in \mathbb{R}, \mathbb{P}(\mathcal{V}_{(k-1)\tau, (k+2)\tau} + 2\mathcal{W}_{(k-1)\tau, (k+2)\tau} \geq x) \leq 2\mathbb{P}(2\mathcal{V}_{0,3\tau} \geq \frac{x}{2}) \leq 4\mathbb{P}(8|W_{3\tau}| \geq x).$$

So, by Lemma 7.7,

$$\begin{aligned} \mathbb{E}\left(\inf\left(1, \frac{T(\Delta)}{(\epsilon'_{k+2})^4\epsilon_{k+1}^8}\right)\right) &\leq \int_0^{+\infty} \inf\{1, T(\Delta)\epsilon(C\tau M + Cz, \Delta)^{-8} \\ &\quad \times \epsilon'(C\tau M + Cz, \Delta)^{-4}\} \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi \times 192\tau}} dz. \end{aligned} \quad (6.11)$$

We have

$$\int_0^{\left(\frac{\Delta}{CB_2(1+p_{2,1})}\right)^+} \epsilon(C\tau M + Cz, \Delta)^{-8} \epsilon'(C\tau M + Cz, \Delta)^{-4} \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi \times 192\tau}} dz$$

$$\begin{aligned}
&\leq 8\epsilon \left(\frac{\Delta}{B_2(1+p_{2,1})}, \Delta \right)^{-8} \epsilon' \left(\frac{\Delta}{B_2(1+p_{2,1})}, \Delta \right)^{-4} \\
&= 8 \exp \left[\frac{20}{\tau} \left(\frac{\Delta}{B_2(1+p_{2,1})} \right)^2 + 32 \left(\frac{M\Delta}{B_2(1+p_{2,1})} \right) + 8 \left(\tau + \frac{\tau^2}{2} \right) M + 8B_2p_{2,1}^2 \left(\frac{\Delta}{B_2(1+p_{2,1})} \right)^2 \right. \\
&\quad \left. + 8\Delta p_{2,1} \left(\frac{\Delta}{B_2(1+p_{2,1})} \right) + 24M \left(\frac{\Delta}{B_2(1+p_{2,1})} \right) + 6\tau \left(M + \frac{M^2}{2} \right) \right]. \quad (6.12)
\end{aligned}$$

From Subsection 2.5 (remember also Equation (4.3)), we get

$$p_{2,1} = O\left(\frac{1}{\theta}\right), \quad B_2 \xrightarrow{\theta \rightarrow +\infty} \frac{h}{2}. \quad (6.13)$$

So there exists τ_0 , such that, for $\tau \geq \tau_0$,

$$\begin{aligned}
&\log \left(\int_0^{\left(\frac{\Delta}{CB_2(1+p_{2,1})} - \tau M\right)_+} \inf\{1, T(\Delta)\epsilon(C\tau M + Cz, \Delta)^{-8} \times \epsilon'(C\tau M + Cz, \Delta)^{-4}\} \right. \\
&\quad \left. \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi} \times 192\tau} dz \right) \underset{\Delta, c}{\leq} -\inf\left(\frac{1}{h}, C_0\right) \Delta^2. \quad (6.14)
\end{aligned}$$

We now want to bound

$$\int_{\left(\frac{\Delta}{CB_2(1+p_{2,1})} - \tau M\right)_+}^{+\infty} \inf\{1, T(\Delta)\epsilon(C\tau M + Cz, \Delta)^{-8} \times \epsilon'(C\tau M + Cz, \Delta)^{-4}\} \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi} \times 192\tau} dz.$$

Let us set, for $z \geq \left(\frac{\Delta}{CB_2(1+p_{2,1})} - \tau M\right)_+$,

$$\begin{aligned}
\Phi(\Delta, z) &= T(\Delta)\epsilon(C\tau M + Cz, \Delta)^{-8} \times \epsilon'(C\tau M + Cz, \Delta)^{-4} \\
&= T(\Delta) \times \exp\left(\frac{16(C\tau M + Cz)\Delta}{\tau B_2(1+p_{2,1})} + 16M \left(C\tau M + Cz + \frac{\Delta}{B_2(1+p_{2,1})}\right) + 8 \left(\tau + \frac{\tau^2}{2}\right) M\right) \\
&\quad \times \exp\left[\left(2B_2p_{2,1}^2 + \frac{1}{\tau}\right) \left(\frac{\Delta}{B_2(1+p_{2,1})} + C\tau M + Cz\right)^2\right. \\
&\quad \left.+ (4\Delta p_{2,1} + 12M) \left(\frac{\Delta}{B_2(1+p_{2,1})} + C\tau M + Cz\right) + 6\tau \left(M + \frac{M^2}{2}\right)\right].
\end{aligned}$$

For $z \leq \left(\frac{\Delta}{CB_2(1+p_{2,1})} - \tau M\right)_+$, we define $\Phi(\Delta, z)$ by

$$\begin{aligned}
\Phi(\Delta, z) &= \\
&T(\Delta) \times \exp\left(\frac{16(C\tau M + Cz)\Delta}{\tau B_2(1+p_{2,1})} + 16M \left(C\tau M + Cz + \frac{\Delta}{B_2(1+p_{2,1})}\right) + 8 \left(\tau + \frac{\tau^2}{2}\right) M\right) \\
&\quad \times \exp\left[\left(2B_2p_{2,1}^2 + \frac{1}{\tau}\right) \left(\frac{\Delta}{B_2(1+p_{2,1})} + C\tau M + Cz\right)^2\right. \\
&\quad \left.+ (4\Delta p_{2,1} + 12M) \left(\frac{\Delta}{B_2(1+p_{2,1})} + C\tau M + Cz\right) + 6\tau \left(M + \frac{M^2}{2}\right)\right].
\end{aligned}$$

For a constant \bar{C} bigger than C , we define $\bar{\Phi}(\Delta, z)$ to have the same expression as Φ , except that we replace C by \bar{C} . We choose a

$$\bar{C} = \sup\left(C, \frac{1}{\sqrt{2 \times 192}}\right), \quad (6.15)$$

so that $z \mapsto \bar{\Phi}(\Delta, z) \times \exp(-z^2/(2 \times 192\tau))$ is nondecreasing in z . We have $\bar{\Phi}(\Delta, z) \rightarrow \infty$ when $z \rightarrow +\infty$ and Δ is fixed. Let us set, for a fixed Δ ,

$$z_0 = \inf\{z : \bar{\Phi}(\Delta, z) \geq 1\}.$$

There exists $\lambda_1 > 0$ and $\tau_1 > 0$ such that for all $\tau \geq \tau_1$,

$$T(\Delta)\epsilon(\bar{C}\tau M + \bar{C}\lambda_1\sqrt{\tau}\Delta, \Delta)^{-8}\epsilon'(\bar{C}\tau M + \bar{C}\lambda_1\sqrt{\tau}\Delta, \Delta)^{-4} \xrightarrow{\Delta \rightarrow +\infty} 0.$$

Looking at the definition of $p_{2,1}$ (Equation (4.3)) and at Equations (7.22), (7.20), we see that λ_1 can be chosen as a function of h , which we denote by $\lambda_1(h)$. And there exists $\lambda_2 > 0$ and $\tau_2 > 0$ such that for all $\tau \geq \tau_2$,

$$T(\Delta)\epsilon(\bar{C}\tau M + \bar{C}\lambda_2\sqrt{\tau}\Delta, \Delta)^{-8}\epsilon'(\bar{C}\tau M + \bar{C}\lambda_2\sqrt{\tau}\Delta, \Delta)^{-4} \xrightarrow{\Delta \rightarrow +\infty} +\infty.$$

So there exists Δ_1 such that, for Δ bigger than Δ_1 and τ bigger than $\sup(\tau_1, \tau_2)$,

$$\lambda_1\sqrt{\tau}\Delta \leq z_0 \leq \lambda_2\sqrt{\tau}\Delta.$$

We can then bound, if $\tau \geq \sup(\tau_1, \tau_2)$,

$$\begin{aligned} \int_0^{z_0} \inf(1, \Phi(\Delta, z)) \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi(192\tau)}} dz &\leq \int_0^{z_0} \inf(1, \bar{\Phi}(\Delta, z)) \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi(192\tau)}} dz \\ &\leq z_0 \bar{\Phi}(\Delta, z_0) \frac{8 \exp\left(-\frac{z_0^2}{2(192\tau)}\right)}{\sqrt{2\pi(192\tau)}} \\ &= z_0 \frac{8 \exp\left(-\frac{z_0^2}{2(192\tau)}\right)}{\sqrt{2\pi(192\tau)}} \\ &\leq \lambda_2\sqrt{\tau}\Delta \frac{8 \exp\left(-\frac{\lambda_1^2\Delta^2}{2 \times 192}\right)}{\sqrt{2\pi(192\tau)}}. \end{aligned}$$

So, if $\tau \geq \sup(\tau_0, \tau_1, \tau_2)$, we get, using again Equation (4.36)

$$\begin{aligned} &\int_{\left(\frac{\Delta}{C_{B_2(1+p_2,1)}-\tau^2 M}\right)_+}^{+\infty} \inf\{1, T(\Delta)\epsilon(C\tau^2 M + Cz, \Delta)^{-8} \times \epsilon'(C\tau^2 M + Cz, \Delta)^{-4}\} \\ &\quad \times \frac{8 \exp\left(-\frac{z^2}{2(192\tau)}\right)}{\sqrt{2\pi \times 192\tau}} dz \\ &\leq \int_0^{z_0} \Phi(\Delta, z) \frac{e^{-\frac{z^2}{2(192\tau)}}}{\sqrt{2\pi \times 192\tau}} dz + \int_{z_0}^{+\infty} \frac{e^{-\frac{z^2}{2(192\tau)}}}{\sqrt{2\pi \times 192\tau}} dz \\ &\leq \lambda_2\sqrt{\tau}\Delta \frac{8 \exp\left(-\frac{\lambda_1^2\Delta^2}{384}\right)}{\sqrt{2\pi(192\tau)}} + \frac{e^{-\frac{z_0^2}{2(192\tau)}}}{z_0\sqrt{2\pi}} \times \sqrt{192\tau}, \quad (6.16) \end{aligned}$$

and so

$$\begin{aligned} &\log \left(\int_{-\infty}^{+\infty} \inf(1, T(\Delta)\epsilon(C\tau M + Cz, \Delta)^{-8} \times \epsilon'(C\tau M + Cz, \Delta)^{-4}) \frac{e^{-\frac{z^2}{2(192\tau)}}}{\sqrt{2\pi \times 192\tau}} dz \right) \\ &\leq_{\Delta, c} -\Delta^2 \inf\left(\frac{1}{h}, C_0, \lambda_1^2\right). \quad (6.17) \end{aligned}$$

In the remaining of the proof, we will suppose $\tau \geq \sup(\tau_0, \tau_1, \tau_2)$. Looking at Equation (6.2), we see that we can now bound all the terms on its right-hand side. We have

$$\mathbb{E}(\|\pi_{n\tau} - \overline{R}_n^\Delta(\pi_{(n-1)\tau})\|) \preceq T(\Delta),$$

by Proposition 4.7. Recall, that, from Equations (6.7), (4.22), we get

$$\log(T(\Delta)) \underset{\Delta, c}{\preceq} -\Delta^2 \inf\left(\frac{1}{h}, C_0\right). \quad (6.18)$$

For k in $\{1, \dots, n-1\}$, we have bounded

$$\begin{aligned} & \mathbb{E}(\|\overline{R}_{n:k+1}^\Delta(\pi_{k\Delta}) - \overline{R}_{n:k+1}^\Delta(\overline{R}_k^\Delta(\pi_{(k-1)\Delta}))\|) \\ & \leq \left(1 - \frac{\epsilon(L, \Delta)^2 \epsilon'(L, \Delta)^2}{2}\right)^{\left[\frac{1}{2}(\lfloor \frac{n-k}{2} \rfloor - 4)_+\right]} \times 2\mathbb{E}\left(\inf\left(1, \frac{\|\pi_{k\tau} - \overline{R}_k^\Delta(\pi_{(k-1)\tau})\|}{(\epsilon'_{k+2})^2 \epsilon_{k+1}^4}\right)\right). \end{aligned}$$

And the last expectation can be bounded by the sum of the following expectations :

$$\begin{aligned} & \mathbb{E}\left(\inf\left(1, \frac{\|\pi_{k\tau} - \overline{R}_k^\Delta(\pi_{(k-1)\tau})\|^2}{T(\Delta)}\right)\right) \preceq \sqrt{T(\Delta)}, \\ & \mathbb{E}\left(\inf\left(1, \frac{T(\Delta)}{(\epsilon'_{k+2})^4 \epsilon_{k+1}^8}\right)\right) \leq \exp\left(-\widehat{B}_1 \Delta^2 \inf\left(\frac{1}{h}, C_0, \lambda_1^2\right)\right) \text{ for } \Delta \geq \Delta_0(\tau), \end{aligned}$$

for some constant \widehat{B}_1 and some function Δ_0 , where the bounds come from Equations (6.6), (6.11), (6.17) (we also use Lemma 2.6). The constant \widehat{B}_1 above is universal and Δ_0 is continuous in τ . So we get, for all $\Delta \geq \Delta_0(\tau)$, using Equation (6.18),

$$\mathbb{E}(\|\pi_{n\tau} - \pi_{n\tau}^\Delta\|) \leq \exp\left(-\widehat{C}_1 \Delta^2 \inf\left(\frac{1}{h}, C_0, \lambda_1^2\right)\right) \sum_{k \geq 0} \left(1 - \frac{\epsilon(L, \Delta)^4 \epsilon'(L, \Delta)^2}{2}\right)^{\left[\frac{1}{2}(\lfloor \frac{n-k}{2} \rfloor - 4)_+\right]},$$

(for some universal constant \widehat{C}_1) from which we get

$$\sup_{n \geq 0} \log \mathbb{E}(\|\pi_{n\tau} - \pi_{n\tau}^\Delta\|) \underset{\Delta, c}{\preceq} -\log(\epsilon(L, \Delta) \epsilon'(L, \Delta)) - \Delta^2 \inf\left(\frac{1}{h}, C_0, \lambda_1^2\right).$$

Looking at Equations (6.8), (6.9), we see there exists τ_3 , such that, for $\tau > \tau_3$,

$$\sup_{n \geq 0} \log \mathbb{E}(\|\pi_{n\tau} - \pi_{n\tau}^\Delta\|) \underset{\Delta, c}{\preceq} -\Delta^2 \inf\left(\frac{1}{h}, C_0, \lambda_1^2\right).$$

□

6.3 Stability of the optimal filter

Proof of Theorem 1.2. We decompose, for all n ,

$$\begin{aligned} \|\pi_{n\tau} - \pi'_{n\tau}\| & \leq \|\pi_{n\tau} - \overline{R}_n^\Delta \dots \overline{R}_1^\Delta(\pi_0)\| + \|\overline{R}_n^\Delta \dots \overline{R}_1^\Delta(\pi_0) - \overline{R}_n^\Delta \dots \overline{R}_1^\Delta(\pi'_0)\| \\ & \quad + \|\overline{R}_n^\Delta \dots \overline{R}_1^\Delta(\pi_0) - \pi'_{n\tau}\|. \end{aligned}$$

Let τ_∞ be the parameter defined in Proposition 1.5. Recall that the operators $(R_n)_{n \geq 0}$, $(R_n^\Delta)_{n \geq 0}$ depend on τ . Suppose that L is such that (as in Equation (6.1))

$$L > 3|m_0| + 3CM\tau_\infty, \quad \tilde{\alpha}(L) \leq \frac{1}{4}.$$

Then, as in Equation (6.4), we have, for all $\tau \in [\tau_\infty, 2\tau_\infty]$, for all $n \geq 0$,

$$\begin{aligned} & \mathbb{E}(\|\bar{R}_n^\Delta \dots \bar{R}_1^\Delta(\pi_0) - \bar{R}_n^\Delta \dots \bar{R}_1^\Delta(\pi'_0)\|) \\ & \leq \left(1 - \frac{\epsilon(L, \Delta)^2 \epsilon'(L, \Delta)^2}{2}\right)^{\left[\frac{1}{2}(\lfloor \frac{n}{2} \rfloor - 4)_+\right]} \times 2\mathbb{E}\left(\inf\left(1, \frac{\|\pi_0 - \pi'_0\|}{(\epsilon'_2)^2 \epsilon_1^4}\right)\right) \\ & \leq 2\left(1 - \frac{\epsilon(L, \Delta)^2 \epsilon'(L, \Delta)^2}{2}\right)^{\left[\frac{1}{2}(\lfloor \frac{n}{2} \rfloor - 4)_+\right]}. \end{aligned}$$

We have (using Equations (6.8), (6.9))

$$\log(\epsilon(L, \Delta)\epsilon'(L, \Delta)) \underset{\Delta, c}{\geq} - \left(\frac{\Delta^2}{\tau B_2^2(1+p_{2,1})^2} + \frac{(B_2 p_{2,1}^2 + \frac{1}{\tau})\Delta^2}{B_2^2(1+p_{2,1})^2} + \frac{p_{2,1}\Delta^2}{B_2(1+p_{2,1})}\right).$$

We now take a sequence $\Delta_n = \sqrt{\nu \log(n)}$, for some $\nu > 0$. By Lemma 2.6, 3, there exist a constants b_1 and an integer n_0 such that, for all $\tau \in [\tau_\infty, 2\tau_\infty]$, for $n > n_0$,

$$\begin{aligned} & \left(1 - \frac{(\epsilon(L, \Delta_n)\epsilon'(L, \Delta_n))^2}{2}\right)^{\left[\frac{1}{2}(\lfloor \frac{n}{2} \rfloor - 4)_+\right]} \\ & \leq \exp\left[-\frac{1}{2}\left[\frac{1}{2}\left(\lfloor \frac{n}{2} \rfloor - 4\right)_+\right]\right] \\ & \times \exp\left(-b_1 \Delta_n^2 \left(\frac{1}{\tau B_2^2(1+p_{2,1})^2} + \frac{B_2 p_{2,1}^2 + \frac{1}{\tau}}{B_2^2(1+p_{2,1})^2} + \frac{p_{2,1}}{B_2(1+p_{2,1})}\right)\right) \\ & = \exp\left[-\frac{1}{2}\left[\frac{1}{2}\left(\lfloor \frac{n}{2} \rfloor - 4\right)_+\right] n^{-\nu'}\right], \quad (6.19) \end{aligned}$$

with

$$\nu' = b_1 \nu \left(\frac{1}{\tau B_2^2(1+p_{2,1})^2} + \frac{B_2 p_{2,1}^2 + \frac{1}{\tau}}{B_2^2(1+p_{2,1})^2} + \frac{p_{2,1}}{B_2(1+p_{2,1})}\right).$$

By Proposition 1.5, we know there exists a constants b'_1 and a integer n'_0 such that, for all $\tau \in [\tau_\infty, 2\tau_\infty]$ and $n \geq n'_0$,

$$\begin{aligned} & \sup(\mathbb{E}(\|\pi_{n\tau} - \bar{R}_n^{\Delta_n} \dots \bar{R}_1^{\Delta_n}(\pi_0)\|), \mathbb{E}(\|\pi'_{n\tau} - \bar{R}_n^{\Delta_n} \dots \bar{R}_1^{\Delta_n}(\pi'_0)\|)) \\ & \leq \exp(-b'_1 \Delta_n^2 \lambda'_1(h)) \leq n^{-\nu''}, \end{aligned}$$

with $\nu'' = b'_1 \nu \lambda'_1(h)$. Let us set $\epsilon \in (0, 1)$. We choose

$$\nu = \frac{(1-\epsilon)}{b_1} \left(\frac{1}{\tau B_2^2(1+p_{2,1})^2} + \frac{B_2 p_{2,1}^2 + \frac{1}{\tau}}{B_2^2(1+p_{2,1})^2} + \frac{p_{2,1}}{B_2(1+p_{2,1})}\right)^{-1},$$

which leads to $\nu' = 1 - \epsilon$. We set $\nu_0 = \nu''$. For any $t \geq \tau_\infty$, if we set $n = \lfloor t/\tau_\infty \rfloor$, then $t = n\tau$ with $\tau \in [\tau_\infty, 2\tau_\infty]$, and so :

$$\begin{aligned} & \mathbb{E}(\|\pi_t - \pi'_t\|) \leq 2n^{-\nu_0} + 2\exp\left(-\frac{1}{2}\left[\frac{1}{2}\left(\lfloor \frac{n-k}{2} \rfloor - 4\right)_+\right] n^{-\nu'}\right), \\ & \mathbb{E}(\|\pi_t - \pi'_t\|) = O(t^{-\nu_0}). \end{aligned}$$

□

Remark 6.2. One could seek to obtain a sharper bound in the above Theorem by choosing another sequence $(\Delta_n)_{n \geq 0}$. Up to some logarithmic terms, the bound would still be a power of t .

7 Appendix

7.1 Proofs of Section 1

Proof of Lemma 2.1. Following [BC09] (Chapter 6, Section 6.1), we introduce the process

$$\widehat{V}_t = V_t + \int_0^t f(X_s) ds, \quad t \geq 0.$$

We introduce a new probability $\widetilde{\mathbb{P}}$ defined by

$$\left. \frac{d\mathbb{P}}{d\widetilde{\mathbb{P}}} \right|_{\mathcal{F}_t} = \exp \left(\int_0^t f(X_s) d\widehat{V}_s - \frac{1}{2} \int_0^t f(X_s)^2 ds \right). \quad (7.1)$$

By Girsanov's theorem, \widehat{V} is a standard Brownian motion under $\widetilde{\mathbb{P}}$. We set F to be a primitive of f . We have, for all $t \geq 0$,

$$\begin{aligned} \int_0^t f(X_s) d\widehat{V}_s - \frac{1}{2} \int_0^t f(X_s)^2 ds &= \int_0^t f(X_s) dX_s - \frac{1}{2} \int_0^t f(X_s)^2 ds \\ &= F(X_t) - F(X_0) - \frac{1}{2} \int_0^t f'(X_s) ds - \frac{1}{2} \int_0^t f(X_s)^2 ds \\ &\geq -M|X_t - X_0| - \frac{Mt}{2} - \frac{M^2 t}{2}. \end{aligned}$$

So, for any test function φ in $\mathcal{C}_b^+(\mathbb{R})$ (the set of bounded continuous functions on \mathbb{R}), $t \geq 0$

$$\begin{aligned} \mathbb{E}(\varphi(X_t)) &= \mathbb{E}^{\mathbb{P}}(\varphi(X_t)) \\ &= \mathbb{E}^{\widetilde{\mathbb{P}}} \left(\varphi(X_t) \left. \frac{d\mathbb{P}}{d\widetilde{\mathbb{P}}} \right|_{\mathcal{F}_t} \right) \\ &\geq \mathbb{E}^{\widetilde{\mathbb{P}}} \left(\varphi(X_t) \exp \left(-M|X_t - X_0| - \frac{Mt}{2} - \frac{M^2 t}{2} \right) \right). \end{aligned}$$

Similarly:

$$\mathbb{E}(\varphi(X_t)) \leq \mathbb{E}^{\widetilde{\mathbb{P}}} \left(\varphi(X_t) \exp \left(M|X_t - X_0| + \frac{Mt}{2} \right) \right)$$

So we have the result. \square

Proof of Lemma 2.2. For any test function φ in $\mathcal{C}_b^+([0, t])$ and any $t \geq 0$,

$$\begin{aligned} \mathbb{E}^{\mathbb{P}}(\varphi(Y_{0:t}) | X_0, X_t) &= \frac{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\varphi(Y_{0:t}) \left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t \right)}{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t \right)} \\ &= \frac{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\varphi(Y_{0:t}) \mathbb{E}^{\widehat{\mathbb{P}}} \left(\left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t, Y_{0:t} \right) \middle| X_0, X_t \right)}{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t \right)}. \end{aligned}$$

By Girsanov's Theorem, (\widehat{V}, Y) is a standard two-dimensional Brownian motion under $\widehat{\mathbb{P}}$. So, conditionally on X_0, X_t , the law of $Y_{0:t}$ under \mathbb{P} has the following density with respect to the Wiener measure:

$$y_{0:t} \mapsto \psi_t(Y_{0:t}, X_0, X_t) = \frac{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t, Y_{0:t} \right)}{\mathbb{E}^{\widehat{\mathbb{P}}} \left(\left. \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}} \right|_{\mathcal{F}_t} \middle| X_0, X_t \right)}$$

We have

$$\begin{aligned} \frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t} &= \exp\left(F(X_1) - F(X_0) - \frac{1}{2}\int_0^t f'(X_s)ds - \frac{1}{2}\int_0^1 f(X_s)^2 ds \right. \\ &\quad \left. + \int_0^t hX_s dY_s - \frac{1}{2}\int_0^t h^2 X_s^2 ds\right) \\ \mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t\right) &= \mathbb{E}^{\widehat{\mathbb{P}}}\left(\exp\left(F(X_1) - F(X_0) - \frac{1}{2}\int_0^t f'(X_s)ds - \frac{1}{2}\int_0^t f(X_s)^2 ds\right)\Big|X_0, X_1\right), \end{aligned} \quad (7.2)$$

so

$$\exp\left(-M|X_t - X_0| - \frac{t(M + M^2)}{2}\right) \leq \mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t\right) \leq \exp\left(M|X_t - X_0| + \frac{tM}{2}\right).$$

Using the above calculations (Equation (7.2)), we can write:

$$\begin{aligned} \exp\left(-M|X_t - X_0| - \frac{t(M + M^2)}{2}\right) \widehat{\psi}_t(Y_{0:t}, x_0, x_1) &\leq \\ \mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t, Y_{0:t}\right) &\leq \exp\left(M|X_t - X_0| + \frac{tM}{2}\right) \widehat{\psi}(Y_{0:t}, x_0, x_1). \end{aligned}$$

So we have the result. \square

Proof of Lemma 1.1. We define a new probability $\check{\mathbb{P}}$ by

$$\frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t} = \exp\left(\int_0^t hX_s dY_s - \frac{1}{2}\int_0^t h^2 X_s^2 ds\right), \forall t \geq 0.$$

By Girsanov's Theorem, (Y_t) is a Brownian motion under $\check{\mathbb{P}}$. For all bounded continuous φ and all $t \geq 0$, we have (Kallianpur-Striebel, see [BC09], p.57)

$$\mathbb{E}(\varphi(X_t)|Y_{0:t}) = \frac{\mathbb{E}^{\check{\mathbb{P}}}\left(\varphi(X_t) \frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}\right)}{\mathbb{E}^{\check{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}\right)},$$

and

$$\mathbb{E}^{\check{\mathbb{P}}}\left(\varphi(X_t) \frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}\right) = \mathbb{E}^{\check{\mathbb{P}}}\left(\varphi(X_t) \mathbb{E}^{\check{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}, X_0, X_t\right)\Big|Y_{0:t}\right),$$

and

$$\begin{aligned} \mathbb{E}^{\check{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\check{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}, X_0, X_t\right) &= \frac{\mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t} \frac{d\check{\mathbb{P}}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}, X_0, X_t\right)}{\mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\check{\mathbb{P}}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}, X_0, X_t\right)} \\ &= \frac{\psi_t(Y_{0:t}, X_0, X_t) \mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t\right)}{\mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|Y_{0:t}, X_0, X_t\right)} \\ &= \psi_t(Y_{0:t}, X_0, X_t) \times \frac{\mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t\right)}{\mathbb{E}^{\widehat{\mathbb{P}}}\left(\frac{d\mathbb{P}}{d\widehat{\mathbb{P}}}\Big|_{\mathcal{F}_t}\Big|X_0, X_t\right)} \end{aligned}$$

$$= \psi_t(Y_{0:t}, X_0, X_t).$$

As the law of $(X_s)_{s \geq 0}$ is the same under \mathbb{P} or $\check{\mathbb{P}}$, we get the desired result. \square

7.2 Proofs of Section 2

We first prove two technical Lemmas.

Lemma 7.1. *For any $t > 0$, for any function $g : \mathbb{R} \mapsto \mathbb{R}$ which is measurable with respect to the Lebesgue measure and such that $\int_0^t g(s)^2 ds < \infty$, we have :*

$$\int_0^t g(s) dB_s = \int_0^t \left(g(s) - \theta e^{\theta s} \int_s^t e^{-\theta u} g(u) du \right) d\beta_s.$$

Proof. Under \mathbb{Q} , B is an Ornstein-Uhlenbeck process (see Equation (2.2)). We can write B as the strong solution of (2.2):

$$B_t = e^{-\theta t} \int_0^t e^{\theta s} d\beta_s, \forall t \geq 0. \quad (7.3)$$

We use the integration by parts formula to compute:

$$\begin{aligned} & \int_0^t \left(g(s) - \theta e^{\theta s} \int_s^t e^{-\theta u} g(u) du \right) d\beta_s \\ &= \int_0^t \left(g(s) - \theta e^{\theta s} \int_0^t e^{-\theta u} g(u) du \right) d\beta_s + \int_0^t \left(\theta e^{\theta s} \int_0^s e^{-\theta u} g(u) du \right) d\beta_s \\ &= \int_0^t g(s) d\beta_s - \left(\int_0^t e^{-\theta u} g(u) du \right) \left(\int_0^t \theta e^{\theta s} d\beta_s \right) + \int_0^t \left(\theta e^{\theta s} \int_0^s e^{-\theta u} g(u) du \right) d\beta_s \\ &= \int_0^t g(s) d\beta_s - \int_0^t e^{-\theta u} g(u) \left(\int_0^u \theta e^{\theta s} d\beta_s \right) du - \int_0^t \left(\int_0^s e^{-\theta u} g(u) du \right) \theta e^{\theta s} d\beta_s \\ &\quad + \int_0^t \left(\theta e^{\theta s} \int_0^s e^{-\theta u} g(u) du \right) d\beta_s \\ &= \int_0^t g(s) d\beta_s - \int_0^t \theta g(u) B_u du = \int_0^t g(s) dB_s. \end{aligned}$$

\square

Lemma 7.2. *We have, for all $s, t \geq 0$,*

$$g(s) - \theta e^{\theta s} \int_s^t e^{-\theta u} g(u) du = \begin{cases} e^{\theta(s-t)} & \text{if } g(u) = 1, \forall u, \\ \left(t + \frac{1}{\theta} \right) e^{\theta(s-t)} - \frac{1}{\theta} & \text{if } g(u) = u, \forall u, \\ \left(t^2 + \frac{2t}{\theta} + \frac{2}{\theta^2} \right) e^{\theta(s-t)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2} \right) & \text{if } g(u) = u^2, \forall u. \end{cases}$$

Proof. The proof in the case $g(u) = 1$ is straightforward. We compute, for all $s, t \geq 0$:

$$\begin{aligned} s - \theta e^{\theta s} \int_s^t u e^{-\theta u} du &= s - \theta e^{\theta s} \left[\left(-\frac{u}{\theta} - \frac{1}{\theta^2} \right) e^{-\theta u} \right]_s^t \\ &= s - \theta \left(-\frac{t}{\theta} - \frac{1}{\theta^2} \right) e^{\theta(s-t)} + \theta \left(-\frac{s}{\theta} - \frac{1}{\theta^2} \right) \\ &= \left(t + \frac{1}{\theta} \right) e^{\theta(s-t)} - \frac{1}{\theta}, \end{aligned}$$

$$s^2 - \theta e^{\theta s} \int_s^t u^2 e^{-\theta u} du = s^2 - \theta e^{\theta s} \left[\left(-\frac{u^2}{\theta} - \frac{2u}{\theta^2} - \frac{2}{\theta^3} \right) e^{-\theta u} \right]_s^t$$

$$\begin{aligned}
&= s^2 - \theta e^{\theta s} \left(-\frac{t^2}{\theta} - \frac{2t}{\theta^2} - \frac{2}{\theta^3} \right) e^{-\theta t} + \theta e^{\theta s} \left(-\frac{s^2}{\theta} - \frac{2s}{\theta^2} - \frac{2}{\theta^3} \right) e^{-\theta s} \\
&= \left(t^2 + \frac{2t}{\theta} + \frac{2}{\theta^2} \right) e^{\theta(s-t)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2} \right).
\end{aligned}$$

□

Proof of Lemma 2.3. Lemma 7.1 tells us that the variables G_1, G_2, G_3, G_4 are centered Gaussians under \mathbb{Q} . Using Lemma 7.2, we compute the following expectations:

$$\begin{aligned}
\mathbb{E}^{\mathbb{Q}}(G_1^2) &= \int_0^1 e^{2\theta(s-1)} ds \\
&= \frac{1 - e^{-2\theta}}{2\theta},
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}^{\mathbb{Q}}(G_2^2) &= \int_0^1 \left(\left(1 + \frac{1}{\theta}\right) e^{\theta(s-1)} - \frac{1}{\theta} \right)^2 ds \\
&= \int_0^1 \left(1 + \frac{1}{\theta}\right)^2 e^{2\theta(s-1)} + \frac{1}{\theta^2} - \frac{2}{\theta} \left(1 + \frac{1}{\theta}\right) e^{\theta(s-1)} ds \\
&= \left[\left(1 + \frac{1}{\theta}\right)^2 \frac{e^{2\theta(s-1)}}{2\theta} + \frac{s}{\theta^2} - \left(1 + \frac{1}{\theta}\right) \frac{2e^{\theta(s-1)}}{\theta^2} \right]_0^1 \\
&= \left(1 + \frac{1}{\theta}\right)^2 \frac{(1 - e^{-2\theta})}{2\theta} + \frac{1}{\theta^2} - \left(\frac{2}{\theta^2} + \frac{2}{\theta^3}\right) (1 - e^{-\theta}),
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}^{\mathbb{Q}}(G_3^2) &= \int_0^1 \left(\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{\theta(s-1)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) \right)^2 ds = \\
&\int_0^1 \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right)^2 e^{2\theta(s-1)} + \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right)^2 - 2 \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{\theta(s-1)} \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) ds = \\
&\left[\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right)^2 \frac{e^{2\theta(s-1)}}{2\theta} + \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right)^2 \frac{\theta}{6} - 2 \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) \left(\frac{2s}{\theta^2}\right) e^{\theta(s-1)} \right]_0^1 = \\
&\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right)^2 \frac{(1 - e^{-2\theta})}{2\theta} + \left(\frac{2}{\theta} + \frac{2}{\theta^2}\right)^2 \frac{\theta}{6} - \frac{8}{6\theta^3} - \frac{4}{\theta^2} \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right),
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}^{\mathbb{Q}}(G_1 G_2) &= \int_0^1 e^{\theta(s-1)} \times \left(\left(1 + \frac{1}{\theta}\right) e^{\theta(s-1)} - \frac{1}{\theta} \right) ds \\
&= \left[\left(1 + \frac{1}{\theta}\right) \frac{e^{2\theta(s-1)}}{2\theta} - \frac{e^{\theta(s-1)}}{\theta^2} \right]_0^1 = \left(\frac{1}{2\theta} + \frac{1}{2\theta^2}\right) (1 - e^{-2\theta}) + \frac{e^{-\theta} - 1}{\theta^2},
\end{aligned}$$

$$\begin{aligned}
\mathbb{E}^{\mathbb{Q}}(G_1 G_3) &= \int_0^1 e^{\theta(s-1)} \left(\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{\theta(s-1)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) \right) ds \\
&= \left[\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) \frac{e^{2\theta(s-1)}}{2\theta} - \frac{2s}{\theta^2} e^{\theta(s-1)} \right]_0^1 = \left(\frac{1}{2\theta} + \frac{1}{\theta^2} + \frac{1}{\theta^3}\right) (1 - e^{-2\theta}) - \frac{2}{\theta^2},
\end{aligned}$$

$$\mathbb{E}^{\mathbb{Q}}(G_2 G_3) = \int_0^1 \left(\left(1 + \frac{1}{\theta}\right) e^{\theta(s-1)} - \frac{1}{\theta} \right) \times \left(\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{\theta(s-1)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) \right) ds$$

$$\begin{aligned}
&= \int_0^1 \left(1 + \frac{1}{\theta}\right) \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{2\theta(s-1)} - \frac{1}{\theta} \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) e^{\theta(s-1)} \\
&\quad - \left(1 + \frac{1}{\theta}\right) e^{\theta(s-1)} \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) + \frac{1}{\theta} \left(\frac{2s}{\theta} + \frac{2}{\theta^2}\right) ds \\
&= \left[\left(1 + \frac{1}{\theta}\right) \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2}\right) \frac{e^{2\theta(s-1)}}{2\theta} - \left(\frac{1}{\theta} + \frac{2}{\theta^2} + \frac{2}{\theta^3}\right) \frac{e^{\theta(s-1)}}{\theta} \right. \\
&\quad \left. - \left(1 + \frac{1}{\theta}\right) \frac{2s}{\theta^2} e^{\theta(s-1)} + \left(\frac{s^2}{\theta^2} + \frac{2s}{\theta^3}\right) \right]_0^1
\end{aligned}$$

□

Proof of Lemma 2.4. The coefficient of x^2 in P is

$$\begin{aligned}
-A_2(\theta) &= -h\frac{\theta}{6} + \frac{\sigma_1^2}{2}h\theta^3 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a\right)^2 \\
&\quad + \frac{\sigma_2^2}{2}h\theta^3 \left(b - \frac{\gamma}{2} - \frac{\theta^2\alpha\gamma}{2}\sigma_1^2 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a\right)\right)^2 + \frac{1}{2}h\theta^3c^2. \quad (7.4)
\end{aligned}$$

We compute (using [WR15] software):

$$\alpha = \frac{1}{\sqrt{2\theta}} + o\left(\frac{1}{\theta^n}\right), \forall n \geq 1, \quad (7.5)$$

$$\text{Cov}^{\mathbb{Q}}(G_1, G_3) = \frac{1}{2\theta} - \frac{1}{\theta^2} + \frac{1}{\theta^3} + o\left(\frac{1}{\theta^3}\right), \quad (7.6)$$

$$\beta = \frac{1}{\sqrt{2\theta}} - \frac{\sqrt{2}}{\theta^{3/2}} + \frac{\sqrt{2}}{\theta^{5/2}} + o\left(\frac{1}{\theta^{5/2}}\right), \quad (7.7)$$

$$\beta^2 = \frac{1}{2\theta} - \frac{2}{\theta^2} + \frac{4}{\theta^3} - \frac{4}{\theta^4} + o\left(\frac{1}{\theta^4}\right), \quad (7.8)$$

$$\text{Var}^{\mathbb{Q}}(G_3) = \frac{1}{2\theta} - \frac{2}{3\theta^2} + o\left(\frac{1}{\theta^4}\right), \quad (7.9)$$

$$\gamma = \frac{2}{\theta\sqrt{3}} - \frac{\sqrt{3}}{\theta^2} + \frac{\sqrt{3}}{4\theta^3} + \frac{3\sqrt{3}}{8\theta^4} + o\left(\frac{1}{\theta^4}\right), \quad (7.10)$$

$$\text{Cov}^{\mathbb{Q}}(G_1, G_2) = \frac{1}{2\theta} - \frac{1}{2\theta^2} + o\left(\frac{1}{\theta^3}\right), \quad (7.11)$$

$$\sigma_1^2 = \frac{6}{\theta} + \frac{18}{\theta^2} + \frac{18}{\theta^3} - \frac{54}{\theta^4} + o\left(\frac{1}{\theta^4}\right), \quad (7.12)$$

$$a = \frac{1}{\sqrt{2\theta}} - \frac{1}{\theta^{3/2}\sqrt{2}} + o\left(\frac{1}{\theta^{5/2}}\right), \quad (7.13)$$

$$\text{Var}^{\mathbb{Q}}(G_2) = \frac{1}{2\theta} - \frac{3}{2\theta^3} + o\left(\frac{1}{\theta^3}\right), \quad (7.14)$$

$$b = \frac{\sqrt{3}}{2\theta} - \frac{\sqrt{3}}{4\theta^2} - \frac{9\sqrt{3}}{16\theta^3} + o\left(\frac{1}{\theta^3}\right), \quad (7.15)$$

$$c^2 = \frac{1}{4\theta^2} - \frac{5}{4\theta^3} + o\left(\frac{1}{\theta^3}\right), \quad (7.16)$$

$$\sigma_2^2 = \frac{1}{3\theta^2} - \frac{1}{\theta} + 2 + o\left(\frac{1}{\theta^3}\right). \quad (7.17)$$

From which we deduce

$$\begin{aligned}
b - \frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) &= \frac{\sqrt{3}}{2\theta^3} + o\left(\frac{1}{\theta^3}\right), \\
\frac{\sigma_2^2}{2} h \theta^3 \left(b - \frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \right)^2 &\xrightarrow{\theta \rightarrow +\infty} 0, \\
\frac{1}{2} h \theta^3 c^2 &= h \left(\frac{\theta}{8} - \frac{5}{8} + o(1) \right), \\
\frac{\sigma_1^2}{2} h \theta^3 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right)^2 &= h \left(\frac{\theta}{24} + \frac{1}{8} + o(1) \right), \\
A_2(\theta) &\xrightarrow{\theta \rightarrow +\infty} \frac{h}{2}.
\end{aligned} \tag{7.18}$$

The coefficient of z^2 in P is

$$-B_2(\theta) = -h \frac{\theta}{6} + \frac{\sigma_1^2}{2} h \theta^3 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right)^2 + \frac{\sigma_2^2}{2} h \theta^3 \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right)^2. \tag{7.19}$$

We have:

$$\begin{aligned}
\frac{\sigma_2^2}{2} h \theta^3 \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right)^2 &\xrightarrow{\theta \rightarrow +\infty} 0, \\
\frac{\sigma_1^2}{2} h \theta^3 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right)^2 &= h \left(\frac{\theta}{6} - \frac{1}{2} + o(1) \right), \\
B_2(\theta) &\xrightarrow{\theta \rightarrow +\infty} \frac{h}{2}.
\end{aligned} \tag{7.20}$$

The coefficient of xz in P is

$$\begin{aligned}
C_1(\theta) &= -h \frac{\theta}{6} + \sigma_1^2 h \theta^3 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \\
&\quad + \sigma_2^2 h \theta^3 \left(b - \frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \right) \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right)
\end{aligned} \tag{7.21}$$

(it does not depend on $y_{0:\tau}$). We have:

$$\begin{aligned}
\sigma_2^2 h \theta^3 \left(b - \frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \right) \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma}{2} \sigma_1^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right) &\xrightarrow{\theta \rightarrow +\infty} 0, \\
\sigma_1^2 h \theta^3 \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) &= h \left(\frac{\theta}{6} - \frac{3}{2\theta} + o\left(\frac{1}{\theta}\right) \right), \\
C_1(\theta) &= \frac{3h}{2\theta} + o\left(\frac{1}{\theta}\right).
\end{aligned} \tag{7.22}$$

□

We need the following Lemma before going into the proof of Lemma 2.7.

Lemma 7.3. *For all $k \in \mathbb{N}$, $s \in [k, k+1]$*

$$\left| Y_{\tau s} - \int_k^{k+1} Y_{\tau u} du - Y_{\tau(s+1)} + \int_k^{k+1} Y_{\tau(u+1)} du \right| \leq h\tau^2 M + h\tau \mathcal{V}_{k\tau, (k+1)\tau} + \mathcal{W}_{k\tau, (k+1)\tau},$$

$$\left| \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau k + \tau s} - \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau(k+1) + \tau s} \right| \leq \frac{M\tau + \mathcal{V}_{k\tau, (k+1)\tau} + \mathcal{W}_{k\tau, (k+1)\tau}}{\theta}.$$

And, for all $s \in [0, 1]$,

$$\left| Y_{\tau s} - \int_0^1 Y_{\tau u} du \right| \leq h\tau^2 M + h\tau \mathcal{V}_{0, \tau} + \mathcal{W}_{0, \tau},$$

$$\left| \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau s} \right| \leq \frac{M\tau + \mathcal{V}_{0, \tau} + \mathcal{W}_{0, \tau}}{\theta}.$$

Proof. We write the proof only for the first two formulas. We have, for all $k \in \mathbb{N}$, $s \in [k, k+1]$,

$$\begin{aligned} & \left| Y_{\tau s} - \int_k^{k+1} Y_{\tau u} du - Y_{\tau(s+1)} + \int_k^{k+1} Y_{\tau(u+1)} du \right| \\ &= \left| - \int_{\tau s}^{\tau(s+1)} hX_u du + W_{\tau s} - W_{\tau(s+1)} - \int_k^{k+1} \left(-h \int_{\tau u}^{\tau(u+1)} X_v dv + W_{\tau u} - W_{\tau(u+1)} \right) du \right| \\ &\leq \left| \int_k^{k+1} h \left(\int_{\tau s}^{\tau(s+1)} X_v dv - \int_{\tau u}^{\tau(u+1)} X_v dv \right) du \right| + \mathcal{W}_{k\tau, (k+2)\tau} \\ &= \left| h \int_k^{k+1} \left(\int_{\tau s}^{\tau(s+1)} X_v - X_{v+\tau(u-s)} dv \right) du \right| + \mathcal{W}_{k\tau, (k+2)\tau} \\ &= \left| h \int_k^{k+1} \left(\int_{\tau s}^{\tau(s+1)} \int_{v+\tau(u-s)}^v f(X_t) dt + V_v - V_{v+\tau(u-s)} dv \right) du \right| + \mathcal{W}_{k\tau, (k+2)\tau} \\ &\leq h\tau^2 M + h\tau \mathcal{V}_{k\tau, (k+2)\tau} + \mathcal{W}_{k\tau, (k+2)\tau}, \end{aligned}$$

and (using integration by parts)

$$\begin{aligned} & \left| \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau k + \tau s} - \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau(k+1) + \tau s} \right| \\ &= \left| \int_0^{\tau} \frac{e^{-\theta} \sinh(hs)}{\theta} (h(X_{\tau k + s} - X_{\tau(k+1) + s}) ds + dW_{\tau k + s} - dW_{\tau(k+1) + s}) \right| \\ &\leq \left| \int_0^{\tau} \frac{e^{-\theta} \sinh(hs)}{\theta} h(X_{\tau k + s} - X_{\tau(k+1) + s}) ds \right| \\ &\quad + \left| \frac{e^{-\theta} \sinh(\theta)}{\theta} (W_{\tau(k+1)} - W_{\tau(k+2)}) \right| \\ &\quad + \left| \int_0^{\tau} (W_{\tau k + s} - W_{\tau(k+1) + s}) \frac{e^{-\theta} \cosh(hs)}{\tau} ds \right| \\ &\leq \int_0^{\tau} \frac{e^{-\theta} \sinh(hs)}{\theta} h(\tau M + \mathcal{V}_{k\tau, (k+2)\tau}) ds + \frac{e^{-\theta} \sinh(\theta)}{\theta} \mathcal{W}_{k\tau, (k+2)\tau} + \int_0^{\tau} \mathcal{W}_{k\tau, (k+2)\tau} \frac{e^{-\theta} \cosh(hs)}{\tau} ds \\ &\leq \frac{h(M\tau + \mathcal{V}_{k\tau, (k+2)\tau})}{h\theta} + \frac{\mathcal{W}_{k\tau, (k+2)\tau}}{\theta}. \end{aligned}$$

□

Proof of Lemma 2.7. We write the proof in the case $k = 0$. From (2.3), (2.15), we deduce

$$\begin{aligned} B_1(Y_{0:\tau}, \theta) &= -\sigma_1^2 h\theta^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \lambda_1(Y_{0:\tau}) + h \int_0^1 s dY_{\tau s} \\ &\quad + \sigma_2^2 h\theta^2 \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right) \left(-\lambda_2(Y_{0:\tau}) + \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \lambda_1(Y_{0:\tau}) \right). \end{aligned} \quad (7.23)$$

For further use, we also write the formula for $A_1(Y_{0:\tau}, \theta)$:

$$\begin{aligned} A_1(Y_{0:\tau}, \theta) &= h \int_0^1 (1-s) dY_{s\tau} + \sigma_1^2 h \theta^2 \left(\frac{\alpha}{3} + \frac{\beta}{2} - a \right) \lambda_1(Y_{0:\tau}) \\ &+ \sigma_2^2 h \theta^2 \left(b - \frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \left(-\frac{\alpha}{3} - \frac{\beta}{2} + a \right) \right) \left(-\lambda_2(Y_{0:\tau}) + \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \lambda_1(Y_{0:\tau}) \right) - h \theta^2 \lambda_3(Y_{0:\tau}) c. \end{aligned} \quad (7.24)$$

We have to remember here that $\lambda_1, \lambda_2, \lambda_3$ are functions of $y_{0:\tau}$. So we might write $\lambda_1(y_{0:\tau}), \dots$ to stress this dependency (and the same goes for other quantities). From Lemmas 7.1, 7.2, we get (g_1, g_2 defined below)

$$\begin{aligned} \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}) &= \int_0^1 e^{\theta(s-1)} \times \left(g_1(s) - \theta e^{\theta s} \int_s^1 e^{-\theta u} g_1(u) du \right) ds \\ &= \int_0^1 g_1(s) e^{\theta(s-1)} ds - \int_0^1 e^{-\theta u} g_1(u) \int_0^u \theta e^{2\theta s - \theta} ds du \\ &= \int_0^1 g_1(s) e^{\theta(s-1)} ds - \int_0^1 e^{-\theta u} g_1(u) \left(\frac{e^{2\theta u} - 1}{2} \right) e^{-\theta} du \\ &= \int_0^1 g_1(s) e^{-\theta} \cosh(\theta s) ds, \end{aligned} \quad (7.25)$$

$$\text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{\tau:2\tau}) = \int_0^1 g_2(s) e^{-\theta} \cosh(\theta s) ds, \quad (7.26)$$

with

$$g_1(s) = Y_{\tau s} - \int_0^1 Y_{\tau u} du, \quad g_2(s) = Y_{\tau(s+1)} - \int_0^1 Y_{\tau(u+1)} du,$$

and

$$\begin{aligned} \text{Cov}(G_3, G_4)(Y_{0:\tau}) &= \int_0^1 \left(\left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} \right) e^{\theta(s-1)} - \left(\frac{2s}{\theta} + \frac{2}{\theta^2} \right) \right) \\ &\quad \times \left(g_1(s) - \theta e^{\theta s} \int_s^1 e^{-\theta u} g_1(u) du \right) ds \\ &= \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} \right) \text{Cov}(G_1, G_4)(Y_{0:\tau}) - \int_0^1 \frac{2s}{\theta} g_1(s) ds \\ &\quad + \int_0^1 e^{-\theta u} g_1(u) \int_0^u \left(2s + \frac{2}{\theta} \right) e^{\theta s} ds du \\ &= \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} \right) \text{Cov}(G_1, G_4)(Y_{0:\tau}) - \int_0^1 \frac{2s}{\theta} g_1(s) ds \\ &\quad + \int_0^1 e^{-\theta u} g_1(u) \frac{2u}{\theta} e^{\theta u} du \\ &= \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} \right) \text{Cov}(G_1, G_4)(Y_{0:\tau}), \end{aligned} \quad (7.27)$$

$$\text{Cov}(G_3, G_4)(Y_{\tau:2\tau}) = \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} \right) \text{Cov}(G_1, G_4)(Y_{\tau:2\tau}). \quad (7.28)$$

From (2.8), (7.5)-(7.17), we deduce (using again [WR15])

$$-\sigma_1^2 h \theta^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \lambda_1(Y_{0:\tau}) = -h(2\theta + O(1)) \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}),$$

$$\sigma_2^2 h \theta^2 \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right) = h O(\theta),$$

$$\begin{aligned} -\lambda_2(Y_{0:\tau}) + \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \lambda_1(Y_{0:\tau}) &= \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}) \left(-\frac{1}{\gamma} \left(1 + \frac{2}{\theta} + \frac{2}{\theta^2} - \frac{\beta}{\alpha} \right) + \frac{\beta}{\alpha \gamma} + \frac{\theta^2 \gamma \sigma_1^2}{2} \right) \\ &= \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}) \times O\left(\frac{1}{\theta}\right). \end{aligned} \quad (7.29)$$

So we get

$$\begin{aligned} -\sigma_1^2 h \theta^2 \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \lambda_1(Y_{0:\tau}) + \sigma_2^2 h \theta^2 \left(\frac{\gamma}{2} - \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \left(-\frac{\alpha}{6} + \frac{\beta}{2} \right) \right) \left(-\lambda_2(Y_{0:\tau}) + \frac{\theta^2 \alpha \gamma \sigma_1^2}{2} \lambda_1(Y_{0:\tau}) \right) \\ = -2h(\theta + O(1)) \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}). \end{aligned} \quad (7.30)$$

We have

$$\begin{aligned} \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}) &= \int_0^1 \left(Y_{\tau s} - \int_0^1 Y_{\tau u} du \right) e^{-\theta} \cosh(\theta s) ds \\ &= \left(Y_{\tau} - \int_0^1 Y_{\tau u} du \right) \frac{e^{-\theta} \sinh(\theta)}{\theta} - \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau s} \\ &= \int_0^1 s dY_{\tau s} \times \frac{e^{-\theta} \sinh(\theta)}{\theta} - \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau s}, \end{aligned} \quad (7.31)$$

and so

$$\begin{aligned} -2h(\theta + O(1)) \text{Cov}^{\mathbb{Q}}(G_1, G_4)(Y_{0:\tau}) + h \int_0^1 s dY_{\tau s} &= -2h(\theta + O(1)) \left(\int_0^1 s dY_{\tau s} \right) \left(\frac{1}{2\theta} - \frac{e^{-2\theta}}{2\theta} \right) \\ &+ h \int_0^1 s dY_{\tau s} + h(\theta + O(1)) \times 2 \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau s} \\ &= h \left(\int_0^1 s dY_{\tau s} \right) \times O\left(\frac{1}{\theta}\right) - h(\theta + O(1)) \times 2 \int_0^1 \frac{e^{-\theta} \sinh(\theta s)}{\theta} dY_{\tau s} \end{aligned} \quad (7.32)$$

And so, using Lemma 7.3, Equations (7.30), (7.31), (7.32) (as similar formulas of the ones above are valid if we replace $Y_{0:\tau}$ by $Y_{0:2\tau}$), we get

$$\begin{aligned} |B_1(Y_{0:\tau}, \theta) - B_1(Y_{\tau:2\tau}, \theta)| &\leq \frac{1}{\tau} (h\tau^2 M + h\tau \mathcal{V}_{0,2\tau} + \mathcal{W}_{0,2\tau}) + h\theta \left(\frac{M\tau + \mathcal{V}_{0,2\tau} + \mathcal{W}_{0,2\tau}}{\theta} \right) \\ &\leq Mh\tau + h\mathcal{V}_{0,2\tau} + \left(h + \frac{1}{\tau} \right) \mathcal{W}_{0,2\tau}. \end{aligned}$$

□

7.3 Technical Lemmas used in Section 4

Lemma 7.4. *We have $\lambda_1 = \int_0^\tau f_1(s) dy_{\tau s}$ for some deterministic function f_1 . And the same is true for λ_2, λ_3 .*

Proof. We write the proof only for λ_1 . Using Equation (2.8) and Lemmas 7.1, 7.2 and integrations by parts, we get

$$\begin{aligned} \alpha \lambda_1 &= \text{Cov}^{\mathbb{Q}}(G_1, G_4) \\ &= \text{Cov}^{\mathbb{Q}} \left(B_1, \int_0^1 y_{\tau s} - \left(\int_0^1 y_{\tau u} du \right) dB_s \right) \end{aligned}$$

$$\begin{aligned}
&= \text{Cov}^{\mathbb{Q}} \left(\int_0^1 e^{\theta(s-1)} d\beta_s, \int_0^1 y_{\tau s} - \left(\int_0^1 y_{\tau u} du \right) - \theta e^{\theta s} \int_s^1 e^{-\theta u} \left(y_{\tau u} - \left(\int_0^1 y_{\tau v} dv \right) \right) dud\beta_s \right) \\
&= \int_0^1 e^{\theta(s-1)} \left(y_{\tau s} - \left(\int_0^1 y_{\tau u} du \right) - \theta e^{\theta s} \int_s^1 e^{-\theta u} \left(y_{\tau u} - \left(\int_0^1 y_{\tau v} dv \right) \right) du \right) ds \\
&= \int_0^1 e^{\theta(s-1)} y_{\tau s} ds - \int_0^1 e^{\theta(s-1)} ds \times \int_0^1 y_{\tau u} du - \int_0^1 e^{-\theta u} y_{\tau u} \int_u^1 \theta e^{\theta(2s-1)} ds du \\
&\quad - \int_0^1 \theta e^{\theta(2s-1)} \frac{(e^{-\theta s} - e^{-\theta})}{\theta} ds \times \int_0^1 y_{\tau u} du \\
&= \frac{y_{\tau}}{\theta} - \int_0^1 \frac{e^{\theta(s-1)}}{\theta} dy_{\tau s} - \frac{(1 - e^{-\theta})}{\theta} \left(y_{\tau} - \int_0^1 u dy_{\tau u} \right) - \int_0^1 y_{\tau u} \sinh(\theta(1-u)) du \\
&\quad - \left(\frac{1 - e^{-\theta}}{\theta} - \frac{(1 - e^{-2\theta})}{2\theta} \right) \left(y_{\tau} - \int_0^1 u dy_{\tau u} \right) \\
&= \frac{y_{\tau}}{\theta} - \int_0^1 \frac{e^{\theta(s-1)}}{\theta} dy_{\tau s} - \frac{(1 - e^{-\theta})}{\theta} \left(y_{\tau} - \int_0^1 u dy_{\tau u} \right) + \frac{y_{\tau}}{\theta} - \int_0^1 \frac{\cosh(\theta(1-u))}{\theta} dy_{\tau u} \\
&\quad - \left(\frac{1 - e^{-\theta}}{\theta} - \frac{(1 - e^{-2\theta})}{2\theta} \right) \left(y_{\tau} - \int_0^1 u dy_{\tau u} \right).
\end{aligned}$$

□

Lemma 7.5. *We can write*

$$A_1^{Y_0:\tau} = \int_0^{\tau} (f_1(s)dW_s + f_2(s)dX_s)$$

for some deterministic functions f_1, f_2 .

Proof. From Lemma 7.4 and Equation (7.24), we know there exists a deterministic function g_1 such that

$$A_1^{Y_0:\tau} = \int_0^1 g_1(s) dY_{\tau s}.$$

So, integrating by parts, we get (where G is the antiderivative of g_1 such that $G(0) = 0$)

$$\begin{aligned}
A_1^{Y_0:\tau} &= \int_0^1 g_1(s)(hX_{\tau s}\tau ds + dW_{\tau s}) \\
&= G(1)hX_{\tau}\tau - \int_0^1 G(s)h\tau dX_{\tau s} + \int_0^1 g_1(s)dW_{\tau s} \\
&= \int_0^{\tau} \left(G(1)h - G\left(\frac{s}{\tau}\right)h \right) dX_s + \int_0^{\tau} g_1\left(\frac{s}{\tau}\right) dW_s.
\end{aligned}$$

□

Lemma 7.6. *For any measurable \mathcal{B} , subset of \mathbb{R}^2 and Φ a Gaussian density,*

$$\begin{aligned}
&\int_{(t_1, t_2) \in \mathcal{B}} \Phi(t_1, t_2) \Psi_1(t_1, t_2) dt_1 dt_2 \\
&\leq \int_{\mathcal{B}} \Phi(t_1, t_2) C_1'(h, \tau) \exp(2M|(1, -1)\kappa^{-1}(t_1, t_2)^T| + \tau(2M + M^2)) dt_1 dt_2
\end{aligned}$$

(the constant $C_1'(h, \tau)$ coming from Lemma 4.3).

Proof. For any $\epsilon' < \epsilon$ (ϵ comes from Lemma 4.3), we can write $\mathcal{B} = \sqcup_{i \in I} \mathcal{B}_i$, where $I \subset \mathbb{N}$, \sqcup means that this is a partition, and for all i , the set \mathcal{B}_i is a subset of $B(2(x_i, z_i)\kappa, \epsilon')$ (for some x_i, z_i). For all i , for almost all (t_1, t_2) in $B(2(x_i, z_i)\kappa, \epsilon')$, we have, by Lemma 4.3,

$$\Psi_1(t_1, t_2) \leq C_1'(h, \tau) \exp(4M|x_i - z_i| + \tau(2M + M^2))$$

$$\begin{aligned}
&= C'_1(h, \tau) \exp(4M|(x_i, z_i)(1, -1)^T| + \tau(2M + M^2)) \\
&= C'_1(h, \tau) \exp(4M|2(x_i, z_i)\frac{1}{2}\kappa^{-1}(1, -1)^T| + \tau(2M + M^2)) \\
&\leq C'_1(h, \tau) \exp(4M|(t_1, t_2)\frac{1}{2}\kappa^{-1}(1, -1)^T| + C'\epsilon' + \tau(2M + M^2)),
\end{aligned}$$

for some constant C' . So we have

$$\begin{aligned}
&\int_{(t_1, t_2) \in \mathcal{B}} \Phi(t_1, t_2) \Psi_1(t_1, t_2) dt_1 dt_2 = \sum_{i \in I} \int_{(t_1, t_2) \in \mathcal{B}_i} \Phi(t_1, t_2) \Psi_1(t_1, t_2) dt_1 dt_2 \\
&\leq \sum_{i \in I} \int_{(t_1, t_2) \in \mathcal{B}_i} \Phi(t_1, t_2) C'_1(h, \tau) \exp(4M|(t_1, t_2)\frac{1}{2}\kappa^{-1}(1, -1)^T| + C'\epsilon' + \tau(2M + M^2)) dt_1 dt_2 \\
&= \int_{(t_1, t_2) \in \mathcal{B}} \Phi(t_1, t_2) C'_1(h, \tau) \exp(4M|(t_1, t_2)\frac{1}{2}\kappa^{-1}(1, -1)^T| + C'\epsilon' + \tau(2M + M^2)) dt_1 dt_2.
\end{aligned}$$

And by taking $\epsilon' \rightarrow 0$, we get the desired result. \square

7.4 Technical Lemma used in Section 6

Lemma 7.7. *Suppose that f_1 and f_2 are two probability densities on \mathbb{R}_+ and that φ is a non-decreasing function from \mathbb{R}_+ to \mathbb{R}_+ . Suppose the distribution functions F_1 and F_2 (associated respectively to f_1, f_2) are such that*

$$1 - F_1(x) \leq K(1 - F_2(x)), \text{ for all } x \geq 0,$$

for some constant $K \geq 1$. Then, for X_1 a random variable of density f_1 and X_2 a random variable of density f_2 , we have

$$\mathbb{E}(\varphi(X_1)) \leq K\mathbb{E}(\varphi(X_2)).$$

Proof. Let U be a random variable of uniform law on $[0, 1]$. Let m be such that $F_2(m) = 1 - 1/K$. Let $(K(1 - F_2))^{-1}$ be the inverse of $\widehat{F}_2 : x \in [m, +\infty[\mapsto K(1 - F_2(x)) \in [0, 1]$. We have:

$$\begin{aligned}
\mathbb{E}(\varphi(X_1)) &= \mathbb{E}(\varphi((1 - F_1)^{-1}(U))) \\
&\leq \mathbb{E}(\varphi((K(1 - F_2))^{-1}(U))) \\
&= \mathbb{E}(\varphi(X_2) | X_2 \geq m) \\
&= \mathbb{E}(\varphi(X_2) \mathbb{1}_{[m, +\infty[}(X_2)) \times K \\
&\leq K\mathbb{E}(\varphi(X_2)).
\end{aligned}$$

\square

THANKS : The authors would like to thank the following colleagues, whose help was greatly appreciated: Dan Crisan, François Delarue.

References

- [Ata98] Rami Atar. Exponential stability for nonlinear filtering of diffusion processes in a non-compact domain. *Ann. Probab.*, 26(4):1552–1574, 1998.
- [AZ97] Rami Atar and Ofer Zeitouni. Exponential stability for nonlinear filtering. *Ann. Inst. H. Poincaré Probab. Statist.*, 33(6):697–725, 1997.
- [BC09] Alan Bain and Dan Crisan. *Fundamentals of stochastic filtering*, volume 60 of *Stochastic Modelling and Applied Probability*. Springer, New York, 2009.

- [BCL04] Peter Baxendale, Pavel Chigansky, and Robert Liptser. Asymptotic stability of the Wonham filter: ergodic and nonergodic signals. *SIAM J. Control Optim.*, 43(2):643–669, 2004.
- [Ben81] V. E. Beneš. Exact finite-dimensional filters for certain diffusions with nonlinear drift. *Stochastics*, 5(1-2):65–92, 1981.
- [CR11] D. Crisan and B. Rozovskiĭ. Introduction. In *The Oxford handbook of nonlinear filtering*, pages 1–15. Oxford Univ. Press, Oxford, 2011.
- [CvH07] Pavel Chigansky and Ramon van Handel. Model robustness of finite state nonlinear filtering over the infinite time horizon. *Ann. Appl. Probab.*, 17(2):688–715, 2007.
- [DG01] Pierre Del Moral and Alice Guionnet. On the stability of interacting processes with applications to filtering and genetic algorithms. *Ann. Inst. H. Poincaré Probab. Statist.*, 37(2):155–194, 2001.
- [KV08] M. L. Kleptsyna and A. Yu. Veretennikov. On discrete time ergodic filters with wrong initial data. *Probability Theory and Related Fields*, 141(3):411–444, 2008.
- [LO03] François LeGland and Nadia Oudjane. A robustification approach to stability and to uniform particle approximation of nonlinear filters: the example of pseudo-mixing signals. *Stochastic Process. Appl.*, 106(2):279–316, 2003.
- [MY08] Roger Mansuy and Marc Yor. *Aspects of Brownian motion*. Universitext. Springer-Verlag, Berlin, 2008.
- [Oco99] D. L. Ocone. Asymptotic stability of Beneš filters. *Stochastic Anal. Appl.*, 17(6):1053–1074, 1999.
- [OR05] Nadia Oudjane and Sylvain Rubenthaler. Stability and uniform particle approximation of nonlinear filters in case of non ergodic signals. *Stoch. Anal. Appl.*, 23(3):421–448, 2005.
- [Oud00] Nadia Oudjane. *Stabilité et approximations particulières en filtrage non linéaire application au pistage*. PhD thesis, 2000. Thèse de doctorat Mathématiques et application Rennes 1.
- [Sta05] Wilhelm Stannat. Stability of the filter equation for a time-dependent signal on \mathbb{R}^d . *Appl. Math. Optim.*, 52(1):39–71, 2005.
- [WR15] Inc. Wolfram Research. Mathematica, 2015.