



**HAL**  
open science

## A note on some constants related to the zeta–function and their relationship with the Gregory coefficients

Iaroslav V Blagouchine, Marc-Antoine Coppo

► **To cite this version:**

Iaroslav V Blagouchine, Marc-Antoine Coppo. A note on some constants related to the zeta–function and their relationship with the Gregory coefficients. *Ramanujan Journal (The)*, 2018, 47 (2), pp.457-473. hal-01499011v3

**HAL Id: hal-01499011**

**<https://hal.univ-cotedazur.fr/hal-01499011v3>**

Submitted on 13 Nov 2017

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A note on some constants related to the zeta–function and their relationship with the Gregory coefficients

Iaroslav V. Blagouchine\*

*Steklov Institute of Mathematics at St.-Petersburg, Russia.*

Marc–Antoine Coppo

*Université Côte d’Azur, CNRS, LJAD (UMR 7351), France.*

---

## Abstract

In this article, new series for the first and second Stieltjes constants (also known as generalized Euler’s constant), as well as for some closely related constants are obtained. These series contain rational terms only and involve the so–called Gregory coefficients, which are also known as (reciprocal) logarithmic numbers, Cauchy numbers of the first kind and Bernoulli numbers of the second kind. In addition, two interesting series with rational terms for Euler’s constant  $\gamma$  and the constant  $\ln 2\pi$  are given, and yet another generalization of Euler’s constant is proposed and various formulas for the calculation of these constants are obtained. Finally, we mention in the paper that almost all the constants considered in this work admit simple representations via the Ramanujan summation.

*Keywords:* Stieltjes constants, Generalized Euler’s constants, Series expansions, Ramanujan summation, Harmonic product of sequences, Gregory’s coefficients, Logarithmic numbers, Cauchy numbers, Bernoulli numbers of the second kind, Stirling numbers of the first kind, Harmonic numbers.

---

## I. Introduction and definitions

The zeta-function

$$\zeta(s) \equiv \sum_{n=1}^{\infty} n^{-s} = \prod_{n=1}^{\infty} (1 - p_n^{-s})^{-1}, \quad \operatorname{Re} s > 1$$

$p_n \in \mathbb{P} = \{2, 3, 5, 7, 11, \dots\}$

is of fundamental and long-standing importance in analytic number theory, modern analysis, theory of  $L$ –functions, prime number theory and in a variety of other fields. The  $\zeta$ –function is a meromorphic function on the entire complex plane, except at the point  $s = 1$  at which it has one simple pole with residue 1. The coefficients of the regular part of its Laurent series, denoted  $\gamma_m$ ,

$$\zeta(s) = \frac{1}{s-1} + \gamma + \sum_{m=1}^{\infty} \frac{(-1)^m (s-1)^m}{m!} \gamma_m, \quad s \neq 1. \quad (1)$$

---

\*Corresponding author.

*Email addresses:* iaroslav.blagouchine@pdmi.ras.ru (Iaroslav V. Blagouchine), coppo@unice.fr (Marc–Antoine Coppo)

where  $\gamma$  is Euler's constant<sup>1</sup>, and those of the Maclaurin series  $\delta_m$

$$\zeta(s) = \frac{1}{s-1} + \frac{1}{2} + \sum_{m=1}^{\infty} \frac{(-1)^m s^m}{m!} \delta_m, \quad s \neq 1. \quad (2)$$

are of special interest and have been widely studied in the literature, see e.g. [25], [1, vol. I, letter 71 and following], [23, p. 166 *et seq.*], [28, 29, 24, 26, 2, 11, 5, 33, 27, 34, 20, 6, 32, 37, 19]. The coefficients  $\gamma_m$  are usually called *Stieltjes constants* or *generalized Euler's constants* (both names being in use), while  $\delta_m$  do not possess a special name.<sup>2</sup> It may be shown with the aid of the Euler–MacLaurin summation that  $\gamma_m$  and  $\delta_m$  may be also given by the following asymptotic representations

$$\gamma_m = \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n \frac{\ln^m k}{k} - \frac{\ln^{m+1} n}{m+1} \right\}, \quad m = 1, 2, 3, \dots, \quad (3)$$

and

$$\delta_m = \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n \ln^m k - n m! \sum_{k=0}^m \frac{(-1)^{m+k} \ln^k n}{k!} + (-1)^m m! - \frac{\ln^m n}{2} \right\}, \quad m = 1, 2, 3, \dots, \quad (4)$$

These representations may be translated into these simple expressions

$$\gamma_m = \sum_{k \geq 1}^{\mathcal{R}} \frac{\ln^m k}{k}, \quad \delta_m = \sum_{k \geq 1}^{\mathcal{R}} \ln^m k, \quad m = 1, 2, 3, \dots$$

where  $\sum^{\mathcal{R}}$  stands for the sum of the series in the sense of the Ramanujan summation of divergent series<sup>5</sup>. Due to the reflection formula for the zeta-function  $\zeta(1-s) = 2\zeta(s)\Gamma(s)(2\pi)^{-s} \cos \frac{1}{2}\pi s$ , the numbers  $\delta_m$  and  $\gamma_m$  are related to each other polynomially and also involve Euler's constant  $\gamma$  and the values of the  $\zeta$ -function at naturals. For the first values of  $m$ , this gives

$$\delta_1 = \frac{1}{2} \ln 2\pi - 1 = -0.08106146679 \dots$$

$$\delta_2 = \gamma_1 + \frac{1}{2}\gamma^2 - \frac{1}{2}\ln^2 2\pi - \frac{1}{24}\pi^2 + 2 = -0.006356455908 \dots \quad (6)$$

$$\delta_3 = -\frac{3}{2}\gamma_2 - 3\gamma_1\gamma - \gamma^3 - (3\gamma_1 + \frac{3}{2}\gamma^2 - \frac{1}{8}\pi^2) \ln 2\pi + \zeta(3) + \frac{1}{2}\ln^3 2\pi - 6 = +0.004711166862 \dots$$

and conversely

$$\gamma_1 = \delta_2 + 2\delta_1^2 + 4\delta_1 - \frac{1}{2}\gamma + \frac{1}{24}\pi^2 = -0.07281584548 \dots$$

$$\begin{aligned} \gamma_2 &= -\frac{2}{3}\delta_3 - 2\delta_2(\gamma + 2) - 4\delta_1\delta_2 - \frac{16}{3}\delta_1^3 - 4\delta_1^2(\gamma + 4) - 8\delta_1(\gamma + 1) - \frac{1}{12}\gamma\pi^2 + \frac{1}{3}\gamma^3 + \frac{2}{3}\zeta(3) - \frac{4}{3} \\ &= -0.009690363192 \dots \end{aligned}$$

The relationships between higher-order coefficients become very cumbersome, but may be found via a semi-recursive procedure described in [3]. Although there exist numerous representations for

<sup>1</sup>We recall that  $\gamma = \lim_{n \rightarrow \infty} (H_n - \ln n) = -\Gamma'(1) = 0.5772156649 \dots$ , where  $H_n$  is the  $n$ th harmonic number.

<sup>2</sup>It follows from (2) that  $\delta_m = (-1)^m \{ \zeta^{(m)}(0) + m! \}$

<sup>3</sup>This representation is very old and was already known to Adolf Pilz, Stieltjes, Hermite and Jensen [6, p. 366].

<sup>4</sup>A slightly different expression for  $\delta_m$  was given earlier by Lehmer [32, Eq. (5), p. 266], Sitaramachandrarao [37, Theorem 1], Finch [23, p. 168 *et seq.*] and Connon [19, Eqs. (2.15), (2.19)]. The formula given by these writers differ from our (4) by the presence of the definite integral of  $\ln^m x$  taken over  $[1, n]$ , which in fact may be reduced to a finite combination of logarithms and factorials.

<sup>5</sup>For more details on the *Ramanujan summation*, see [15, 13, 16, 12] and [4, Ch. 6].

<sup>6</sup>This expression for  $\delta_2$  was found by Ramanujan, see e.g. [4, (18.2)].

$\gamma_m$  and  $\delta_m$ , no convergent series with rational terms only are known for them (unlike for Euler's constant  $\gamma$ , see e.g. [6, Sect. 3], or for various expressions containing it [8, p. 413, Eqs. (41), (45)–(47)]). Recently, divergent envelopping series for  $\gamma_m$  containing rational terms only have been obtained in [6, Eqs. (46)–(47)]. In this paper, by continuing the same line of investigation, we derive convergent series representations with rational coefficients for  $\gamma_1, \delta_1, \gamma_2$  and  $\delta_2$ , and also find two new series of the same type for Euler's constant  $\gamma$  and  $\ln 2\pi$  respectively. These series are not simple and involve a product of *Gregory coefficients*  $G_n$ , which are also known as (*reciprocal*) *logarithmic numbers*, *Bernoulli numbers of the second kind*  $b_n$ , and *normalized Cauchy numbers of the first kind*  $C_{1,n}$ . Similar expressions for higher-order constants  $\gamma_m$  and  $\delta_m$  may be obtained by the same procedure, using the harmonic product of sequences introduced in [14], but are quite cumbersome. Since the Stieltjes constants  $\gamma_m$  generalize Euler's constant  $\gamma$  and since our series contain the product of  $G_n$ , these new series may also be seen as the generalization of the famous Fontana–Mascheroni series

$$\gamma = \sum_{n=1}^{\infty} \frac{|G_n|}{n} = \frac{1}{2} + \frac{1}{24} + \frac{1}{72} + \frac{19}{2880} + \frac{3}{800} + \frac{863}{362880} + \frac{275}{169344} + \dots \quad (5)$$

which is the first known series representation for Euler's constant having rational terms only, see [8, pp. 406, 413, 429], [6, p. 379]. In Appendix A, we introduce yet another set of constants  $\kappa_m = \sum_{n \geq 1} |G_n| n^{-m-1}$ , which also generalize Euler's constant  $\gamma$ . These numbers, similarly to  $\gamma_m$ , coincide with Euler's constant at  $m = 0$  and have various interesting series and integral representations, none of them are reducible to classical mathematical constants.

## II. Series expansions

### II.1. Preliminaries

Since the results, that we come to present here, are essentially based on the Gregory coefficients and Stirling numbers, it may be useful to briefly recall their definition and properties. Gregory numbers, denoted below  $G_n$ , are rational alternating  $G_1 = +1/2$ ,  $G_2 = -1/12$ ,  $G_3 = +1/24$ ,  $G_4 = -19/720$ ,  $G_5 = +3/160$ ,  $G_6 = -863/60480, \dots$ , decreasing in absolute value, and are also closely related to the theory of finite differences; they behave as  $(n \ln^2 n)^{-1}$  at  $n \rightarrow \infty$  and may be bounded from below and above according to [8, Eqs. (55)–(56)]. They may be defined either via their generating function

$$\frac{z}{\ln(1+z)} = 1 + \sum_{n=1}^{\infty} G_n z^n, \quad |z| < 1, \quad (6)$$

or recursively by

$$G_n = \frac{(-1)^{n+1}}{n+1} + \sum_{l=1}^{n-1} \frac{(-1)^{n+1-l} G_l}{n+1-l}, \quad G_1 = \frac{1}{2}, \quad n = 2, 3, 4, \dots \quad (7)$$

or explicitly by<sup>7</sup>

$$G_n = \frac{1}{n!} \int_0^1 x(x-1)(x-2) \cdots (x-n+1) dx, \quad n = 1, 2, 3, \dots \quad (8)$$

---

<sup>7</sup>For more information about  $G_n$ , see [8, pp. 410–415], [6, p. 379], [9], and the literature given therein (nearly 50 references).

Throughout the paper, we also use the Stirling numbers of the first kind, which we denote below by  $S_1(n, l)$ . Since there are different definitions and notations for them<sup>8</sup>, we specify that in our definition they are simply the coefficients in the expansion of falling factorial

$$x(x-1)(x-2)\cdots(x-n+1) = \sum_{l=1}^n S_1(n, l) \cdot x^l, \quad n = 1, 2, 3, \dots \quad (9)$$

and may equally be defined via the generating function

$$\frac{\ln^l(1+z)}{l!} = \sum_{n=l}^{\infty} \frac{S_1(n, l)}{n!} z^n = \sum_{n=0}^{\infty} \frac{S_1(n, l)}{n!} z^n, \quad |z| < 1, \quad l = 0, 1, 2, \dots \quad (10)$$

It is important to note that  $\text{sgn}[S_1(n, l)] = (-1)^{n-l}$ . The Stirling numbers of the first kind and the Gregory coefficients are linked by the relation

$$G_n = \frac{1}{n!} \sum_{l=1}^n \frac{S_1(n, l)}{l+1}, \quad n = 1, 2, 3, \dots \quad (11)$$

which follows directly from (8) and (9).

## II.2. Some auxiliary lemmas

Before we proceed with the series expansions for  $\delta_m$  and  $\gamma_m$ , we need to prove several useful lemmas.

**Lemma 1.** For each natural number  $k$ , let

$$\sigma_k := \sum_{n=1}^{\infty} \frac{|G_n|}{n+k}.$$

Then the following equality holds

$$\sigma_k = \frac{1}{k} + \sum_{m=1}^k (-1)^m \binom{k}{m} \ln(m+1), \quad k = 1, 2, 3, \dots \quad (12)$$

*Proof.* By using (11) and by making use of the generating equation for the Stirling numbers of the first kind (10), we obtain

$$\begin{aligned} \sum_{n=1}^{\infty} \frac{|G_n|}{n+k} &= \sum_{n=1}^{\infty} \frac{1}{n!} \sum_{l=1}^n \frac{(-1)^{l+1} |S_1(n, l)|}{l+1} \cdot \underbrace{\int_0^1 x^{n+k-1} dx}_{1/(n+k)} = - \sum_{l=1}^{\infty} \frac{1}{(l+1)!} \int_0^1 x^{k-1} \ln^l(1-x) dx \\ &= \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{(l+1)!} \int_0^{\infty} (1-e^{-t})^{k-1} t^l e^{-t} dt = \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{(l+1)!} \sum_{m=0}^{k-1} (-1)^m \binom{k-1}{m} \underbrace{\int_0^{\infty} e^{-t(m+1)} t^l dt}_{l!(m+1)^{-l-1}} \\ &= \sum_{l=1}^{\infty} \frac{(-1)^{l+1}}{l+1} \sum_{m=0}^{k-1} (-1)^m \binom{k-1}{m} \frac{1}{(m+1)^{l+1}} = \sum_{m=0}^{k-1} (-1)^m \binom{k-1}{m} \left\{ \frac{1}{m+1} - \ln \frac{m+2}{m+1} \right\}, \end{aligned} \quad (13)$$

<sup>8</sup>More information and references (more than 60) on the Stirling numbers of the first kind may be found in [8, Sect. 2.1] and [6, Sect. 1.2]. We also note that our definitions for the Stirling numbers agree with those adopted by MAPLE or MATHEMATICA: our  $S_1(n, l)$  equals to  $\text{Stirling1}(n, l)$  from the former and to  $\text{StirlingS1}[n, l]$  from the latter.

where at the last stage we made a change of variable  $x = 1 - e^{-t}$  and used the well-known formula for the  $\Gamma$ -function. But since

$$\sum_{m=0}^{k-1} \frac{(-1)^m}{m+1} \binom{k-1}{m} = \frac{1}{k}, \quad \text{and} \quad \binom{k-1}{m} + \binom{k-1}{m-1} = \binom{k}{m},$$

the last finite sum in (13) reduces to (12)<sup>9</sup>. □

**Remark 1.** One may show <sup>10</sup>that  $\sigma_k$  may also be written in terms of the Ramanujan summation:

$$\sigma_k = \sum_{n \geq 1}^{\mathcal{R}} \frac{\Gamma(k+1)\Gamma(n)}{\Gamma(n+k+1)} = \sum_{n \geq 1}^{\mathcal{R}} B(k+1, n), \quad (14)$$

where  $B$  stands for the Euler beta-function.

**Lemma 2.** Let  $a = (a(1), a(2), \dots, a(n), \dots)$  be a sequence of complex numbers. The following identity is true for all nonnegative integers  $n$ :

$$\sum_{l=0}^n (-1)^l \binom{n}{l} \frac{a(l+1)}{l+1} = \frac{1}{n+1} \sum_{k=0}^n \sum_{l=0}^k (-1)^l \binom{k}{l} a(l+1). \quad (15)$$

In particular, if  $a = \ln^m$  for any natural  $m$ , then this identity reduces to

$$\sum_{l=0}^n (-1)^l \binom{n}{l} \frac{\ln^m(l+1)}{l+1} = \frac{1}{n+1} \sum_{k=1}^n \sum_{l=1}^k (-1)^l \binom{k}{l} \ln^m(l+1). \quad (16)$$

*Proof.* Formula (15) is an explicit translation of [14, Proposition 7]. □

**Lemma 3.** For all natural  $m$

$$\gamma_m = \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{k=1}^n \sum_{l=1}^k (-1)^l \binom{k}{l} \ln^m(l+1), \quad (17)$$

$$\delta_m = \sum_{n=1}^{\infty} |G_{n+1}| \sum_{l=1}^n (-1)^l \binom{n}{l} \ln^m(l+1). \quad (18)$$

*Proof.* Using the representation for the  $\zeta$ -function

$$\zeta(s) = \frac{1}{s-1} + \sum_{n=0}^{\infty} |G_{n+1}| \sum_{k=0}^n (-1)^k \binom{n}{k} (k+1)^{-s}, \quad s \neq 1,$$

see e.g. [6, pp. 382–383], [7], we first have

$$\gamma_m = \sum_{n=0}^{\infty} |G_{n+1}| \sum_{l=0}^n (-1)^l \binom{n}{l} \frac{\ln^m(l+1)}{l+1},$$

and

$$\delta_m = \sum_{n=0}^{\infty} |G_{n+1}| \sum_{l=0}^n (-1)^l \binom{n}{l} \ln^m(l+1).$$

Then formula (17) follows from (16). □

<sup>9</sup>The result appeared without proof in [8]. For a slightly more general result, see [21, Proposition 1].

<sup>10</sup>See [12] Eq. (4.31).

II.3. Series with rational terms for the first Stieltjes constant  $\gamma_1$  and for the coefficient  $\delta_1$

**Theorem 1.** The first Stieltjes constant  $\gamma_1$  may be given by the following series

$$\begin{aligned}\gamma_1 &= \frac{3}{2} - \frac{\pi^2}{6} + \sum_{n=2}^{\infty} \left[ \frac{|G_n|}{n^2} + \sum_{k=1}^{n-1} \frac{|G_k G_{n+1-k}| (H_n - H_k)}{n+1-k} \right] \\ &= \frac{3}{2} - \frac{\pi^2}{6} + \frac{1}{32} + \frac{5}{432} + \frac{1313}{207360} + \frac{42169}{10368000} + \frac{137969}{48384000} + \frac{1128119}{533433600} + \dots\end{aligned}\quad (19)$$

containing  $\pi^2$  and positive rational coefficients only. Using Euler's formula  $\pi^2 = 6 \sum n^{-2}$ , the latter may be reduced to a series with rational terms only.

*Proof.* By (17) with  $m = 1$ , one has

$$\gamma_1 = \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{k=1}^n \sum_{m=1}^k (-1)^m \binom{k}{m} \ln(m+1).$$

Using (12), we find that

$$\frac{1}{n+1} \sum_{k=1}^n \sum_{m=1}^k (-1)^m \binom{k}{m} \ln(m+1) = \frac{1}{n+1} \sum_{k=1}^n \sigma_k - \frac{H_{n+1}}{n+1} + \frac{1}{(n+1)^2}.$$

Thus

$$\begin{aligned}\gamma_1 &= \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{k=1}^n \sigma_k - \sum_{n=0}^{\infty} \frac{|G_{n+1}| H_{n+1}}{n+1} + \sum_{n=0}^{\infty} \frac{|G_{n+1}|}{(n+1)^2} \\ &= \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{|G_{n+1} G_m| (H_{m+n} - H_m)}{n+1} - \zeta(2) + 1 + \sum_{n=1}^{\infty} \frac{|G_n|}{n^2}\end{aligned}$$

since

$$\sum_{n=1}^{\infty} \frac{|G_n| H_n}{n} = \zeta(2) - 1.$$

For the latter, see e.g. [41, p. 2952, Eq. (1.3)], [17, p. 20, Eq. (3.6)], [13, p. 307, Eq. for  $F_0(2)$ ], [8, p. 413, Eq. (44)]. Rearranging the double absolutely convergent series as follows

$$\sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{|G_{n+1} G_m| (H_{m+n} - H_m)}{n+1} = \sum_{n=2}^{\infty} \sum_{k=1}^{n-1} \frac{|G_k G_{n+1-k}| (H_n - H_k)}{n+1-k},$$

we finally arrive at (19).  $\square$

**Remark 2.** It seems that the sum  $\kappa_1 := \sum |G_n| n^{-2} = 0.5290529699\dots$ <sup>11</sup> cannot be reduced to the "standard" mathematical constants. However, it admits several interesting representations, which we give in Appendix A.

**Theorem 2.** The first MacLaurin coefficient  $\delta_1 = \frac{1}{2} \ln 2\pi - 1$  admits a series representation similar to that for  $\gamma_1$ , namely

$$\delta_1 = \sum_{n=1}^{\infty} \frac{1}{n} \sum_{k=1}^n |G_k G_{n+1-k}| + \frac{1 - \ln 2\pi}{2}. \quad (20)$$

<sup>11</sup>For more digits, see OEIS A270859.

*Proof.* Proceeding analogously to the previous case and recalling that

$$\sum_{n=2}^{\infty} \frac{|G_n|}{n-1} = -\frac{\gamma+1-\ln 2\pi}{2},$$

see e.g. [8, p. 413, Eq. (41)], [42, Corollary 9], we have

$$\begin{aligned} \delta_1 &= \sum_{n=0}^{\infty} |G_{n+1}| \sum_{l=0}^n (-1)^l \binom{n}{l} \ln(l+1) \\ &= \sum_{n=1}^{\infty} |G_{n+1}| \left( \sigma_n - \frac{1}{n} \right) = \sum_{n=1}^{\infty} |G_{n+1}| \sigma_n - \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n} \\ &= \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{|G_{n+1} G_k|}{n+k} + \frac{\gamma+1-\ln 2\pi}{2} = \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \frac{|G_n G_k|}{n+k-1} + \frac{1-\ln 2\pi}{2} \\ &= \sum_{n=1}^{\infty} \frac{1}{n} \sum_{k=1}^n |G_k G_{n+1-k}| + \frac{1-\ln 2\pi}{2}, \end{aligned} \quad (21)$$

where in (21) we eliminated  $\gamma$  by using the fact that  $G_1 = 1/2$  and that the sum of  $|G_n|/n$  over all natural  $n$  equals precisely Euler's constant, see (5).  $\square$

**Corollary 1.** *The constant  $\ln 2\pi$  has the following beautiful series representation with rational terms only and containing a product of Gregory coefficients*

$$\ln 2\pi = \frac{3}{2} + \sum_{n=1}^{\infty} \frac{1}{n} \sum_{k=1}^n |G_k G_{n+1-k}| = \frac{3}{2} + \frac{1}{4} + \frac{1}{24} + \frac{7}{432} + \frac{1}{120} + \frac{43}{8640} + \frac{79}{24192} + \dots \quad (22)$$

*This result directly follows from (20). It is worth noting that one can also readily derive a series with rational coefficients only for  $\ln \pi$  (for instance, with the aids of the Mercator series).*

**Corollary 2.** *Euler's constant  $\gamma$  admits the following series representation with rational terms*

$$\begin{aligned} \gamma &= 2 \ln 2\pi - 3 - 2 \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{k=1}^n |G_k G_{n+2-k}| = 2 \ln 2\pi - 3 - \frac{1}{24} - \frac{1}{54} - \frac{29}{2880} - \\ &\quad - \frac{67}{10800} - \frac{1507}{362880} - \frac{3121}{1058400} - \frac{12703}{15806080} - \frac{164551}{97977600} - \dots \end{aligned} \quad (23)$$

*This result seems to be undiscovered yet. It follows from (5), (21), and the transformation*

$$\sum_{k=1}^{\infty} \sum_{n=1}^{\infty} \frac{|G_{k+1} G_n|}{n+k} = \sum_{n=1}^{\infty} \frac{1}{n+1} \sum_{k=1}^n |G_k G_{n+2-k}|.$$

#### II.4. Generalizations to the second-order coefficients $\delta_2$ and $\gamma_2$ via an application of the harmonic product

We recall the main properties of the harmonic product of sequences which are stated and proved in [14]. If  $a = (a(1), a(2), \dots)$  and  $b = (b(1), b(2), \dots)$  are two sequences in  $\mathbb{C}^{\mathbb{N}^*}$ , then the harmonic product  $a \bowtie b$  admits the explicit expression:

$$(a \bowtie b)(m+1) = \sum_{0 \leq l \leq k \leq m} (-1)^{k-l} \binom{m}{k} \binom{k}{l} a(k+1) b(m+1-l), \quad m = 0, 1, 2, \dots \quad (24)$$



For small values of  $m$ , this gives:

$$\begin{aligned}(a \rtimes b)(1) &= a(1)b(1), \\(a \rtimes b)(2) &= a(2)b(1) + a(1)b(2) - a(2)b(2), \\(a \rtimes b)(3) &= a(3)b(1) + a(1)b(3) + 2a(2)b(2) - 2a(3)b(2) - 2a(2)b(3) + a(3)b(3).\end{aligned}$$

The harmonic product  $\rtimes$  is associative and commutative.

Let  $D$  be the operator (known as the binomial transform) defined by

$$D(a)(m+1) = \sum_{j=0}^m (-1)^j \binom{m}{j} a(j+1), \quad m = 0, 1, 2, \dots$$

then  $D = D^{-1}$  and the harmonic product satisfies the following property:

$$D(ab) = D(a) \rtimes D(b). \quad (25)$$

In particular, if  $a = \ln$ , then  $D(a)(1) = \ln 1 = 0$ . By (12), we find that

$$D(\ln)(m+1) = \sum_{j=1}^m (-1)^j \binom{m}{j} \ln(j+1) = \sigma_m - \frac{1}{m}, \quad m = 1, 2, 3, \dots \quad (26)$$

Therefore, if  $a = \ln^2$  then, by (24), (25), and (26), the following identity holds

$$D(\ln^2)(m+1) = \sum_{\substack{0 \leq l \leq k \leq m \\ k \neq 0 \\ l \neq m}} (-1)^{k-l} \binom{m}{k} \binom{k}{l} \left( \sigma_k - \frac{1}{k} \right) \left( \sigma_{m-l} - \frac{1}{m-l} \right). \quad (27)$$

From this identity, we deduce the following theorem:

**Theorem 3.** *The second coefficients  $\gamma_2$  and  $\delta_2$  may be given by the following series*

$$\gamma_2 = \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{\substack{0 \leq l \leq k \leq m \leq n \\ k \neq 0 \\ l \neq m}} (-1)^{k-l} \binom{m}{k} \binom{k}{l} \left( \sigma_k - \frac{1}{k} \right) \left( \sigma_{m-l} - \frac{1}{m-l} \right),$$

and

$$\delta_2 = \sum_{m=1}^{\infty} |G_{m+1}| \sum_{\substack{0 \leq l \leq k \leq m \\ k \neq 0 \\ l \neq m}} (-1)^{k-l} \binom{m}{k} \binom{k}{l} \left( \sigma_k - \frac{1}{k} \right) \left( \sigma_{m-l} - \frac{1}{m-l} \right).$$

*Proof.* Using (17) and equation (27), we get the following equalities:

$$\begin{aligned}\gamma_2 &= \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{m=1}^n \sum_{j=1}^m (-1)^j \binom{m}{j} \ln^2(j+1) \\ &= \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{m=1}^n D(\ln^2)(m+1) \\ &= \sum_{n=1}^{\infty} \frac{|G_{n+1}|}{n+1} \sum_{m=1}^n \sum_{\substack{0 \leq l \leq k \leq m \\ k \neq 0 \\ l \neq m}} (-1)^{k-l} \binom{m}{k} \binom{k}{l} \left( \sigma_k - \frac{1}{k} \right) \left( \sigma_{m-l} - \frac{1}{m-l} \right).\end{aligned}$$

Similarly  $\delta_2$  is

$$\begin{aligned}
\delta_2 &= \sum_{n=0}^{\infty} |G_{n+1}| \sum_{j=0}^n (-1)^j \binom{n}{j} \ln^2(j+1) \\
&= \sum_{n=1}^{\infty} |G_{n+1}| D(\ln^2)(n+1) \\
&= \sum_{m=1}^{\infty} |G_{m+1}| \sum_{\substack{0 \leq l \leq k \leq m \\ k \neq 0 \\ l \neq m}} (-1)^{k-l} \binom{m}{k} \binom{k}{l} \left( \sigma_k - \frac{1}{k} \right) \left( \sigma_{m-l} - \frac{1}{m-l} \right).
\end{aligned}$$

□

By following the same method, one may also obtain expressions for higher-order constants  $\gamma_m$  and  $\delta_m$ . However, the resulting expressions are more theoretical than practical.

### Appendix A. Yet another generalization of Euler's constant

The numbers  $\kappa_p := \sum |G_n| n^{-p-1}$ , where the summation extends over  $n = [1, \infty)$ , may also be regarded as one of the possible generalizations of Euler's constant (since  $\kappa_0 = \gamma_0 = \gamma$  and  $\kappa_{-1} = \gamma_{-1} = 1$ ).<sup>12,13</sup> These constants, which do not seem to be reducible to the "classical mathematical constants", admit several interesting representations as stated in the following proposition.

**Proposition 1.** *Generalized Euler's constants  $\kappa_p := \sum |G_n| n^{-p-1}$ , where the summation extends over  $n = [1, \infty)$ , admit the following representations:*

$$\kappa_p = \frac{(-1)^p}{\Gamma(p+1)} \int_0^1 \left\{ \frac{1}{\ln(1-x)} + \frac{1}{x} \right\} \ln^p x \, dx, \quad \text{Re } p > -1, \quad (28)$$

$$\underbrace{\int_0^1 \cdots \int_0^1}_{p\text{-fold}} \left\{ \text{li} \left( 1 - \prod_{k=1}^p x_k \right) + \gamma + \sum_{k=1}^p \ln x_k \right\} \frac{dx_1 \cdots dx_p}{x_1 \cdots x_p}, \quad p = 1, 2, 3, \dots \quad (29)$$

$$= \sum_{k=2}^{\infty} \frac{(-1)^k}{k} \sum_{n=p+1}^{\infty} \frac{|S_1(n, p+1)|}{n! n^{k-1}}, \quad p = 0, 1, 2, \dots \quad (30)$$

$$= \sum_{k=2}^{\infty} \frac{(-1)^k}{k} \sum_{n=p}^{\infty} \frac{P_p(H_n^{(1)}, -H_n^{(2)}, \dots, (-1)^{p-1} H_n^{(p)})}{(n+1)^k}, \quad p = 0, 1, 2, \dots \quad (31)$$

$$= \sum_{n \geq 1} \frac{\mathcal{R}}{n} \sum_{n \geq n_1 \geq \dots \geq n_p \geq 1} \frac{1}{n_1 \cdots n_p} = \sum_{n \geq 1} \frac{\mathcal{R}}{n} \frac{P_p(H_n^{(1)}, H_n^{(2)}, \dots, H_n^{(p)})}{n}, \quad p = 1, 2, 3, \dots \quad (32)$$

<sup>12</sup>Numbers  $\kappa_0$  and  $\kappa_{-1}$  are found for the values to which Fontana–Mascheroni and Fontana series converge respectively [8, pp. 406, 410].

<sup>13</sup>Other possible generalizations of Euler's constant were proposed by Briggs, Lehmer, Dilcher and some other authors [10, 31, 38, 35, 39, 22].

where  $\text{li}$  denotes the integral logarithm function,  $H_n^{(m)} := \sum_{k=1}^n k^{-m}$  stands for the generalized harmonic number and the sequence of polynomials  $(P_n)$  is

$$\begin{aligned} P_0 &= 1, & P_1(x_1) &= x_1, & P_2(x_1, x_2) &= \frac{1}{2} (x_1^2 + x_2), \\ P_3(x_1, x_2, x_3) &= \frac{1}{6} (x_1^3 + 3x_1x_2 + 2x_3), & \dots & \end{aligned} \quad 14$$

In particular, for the series  $\kappa_1$  mentioned in Theorem 1 and Remark 2, this gives

$$\kappa_1 = \sum_{n=1}^{\infty} \frac{|G_n|}{n^2} = - \int_0^1 \left\{ \frac{1}{\ln(1-x)} + \frac{1}{x} \right\} \ln x \, dx \quad (33)$$

$$= \int_0^1 \frac{-\text{li}(1-x) + \gamma + \ln x}{x} \, dx = \int_0^{\infty} \left\{ -\text{li}(1-e^{-x}) + \gamma - x \right\} \, dx \quad (34)$$

$$= \sum_{k=2}^{\infty} \frac{(-1)^k}{k} \sum_{n=1}^{\infty} \frac{H_n}{(n+1)^k} = \sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n}. \quad (35)$$

Moreover, we have

$$\kappa_1 = \gamma_1 + \frac{\gamma^2}{2} - \frac{\pi^2}{12} + \int_0^1 \frac{\Psi(x+1) + \gamma}{x} \, dx \quad (36)$$

$$= \frac{\gamma^2}{2} + \frac{\pi^2}{12} - \frac{1}{2} + \frac{1}{2} \int_0^1 \Psi^2(x+1) \, dx, \quad (37)$$

where  $\Psi$  denotes the digamma function (logarithmic derivative of the  $\Gamma$ -function).

*Proof of formula (28)*

We first write the generating equation for Gregory's coefficients, Eq. (6), in the following form

$$\frac{1}{\ln(1-x)} + \frac{1}{x} = \sum_{n=1}^{\infty} |G_n| x^{n-1}, \quad |x| < 1. \quad (38)$$

Multiplying both sides by  $\ln^p x$ , integrating over the unit interval and changing the order of summation and integration<sup>15</sup> yields:

$$\int_0^1 \left\{ \frac{1}{\ln(1-x)} + \frac{1}{x} \right\} \ln^p x \, dx = \sum_{n=1}^{\infty} |G_n| \int_0^1 x^{n-1} \ln^p x \, dx, \quad \text{Re } p > -1. \quad (39)$$

<sup>14</sup>More generally, these polynomials, called *the modified Bell polynomials* are defined by the generating function:  
 $\exp\left(\sum_{k=1}^{\infty} x_k \frac{z^k}{k}\right) = \sum_{n=0}^{\infty} P_n(x_1, \dots, x_n) z^n.$

<sup>15</sup>The series being uniformly convergent.

The last integral may be evaluated as follows. Considering Legendre's integral  $\Gamma(p+1) = \int t^p e^{-t} dt$  taken over  $[0, \infty)$  and making a change of variable  $t = -(s+1) \ln x$ , we have

$$\int_0^1 x^s \ln^p x \, dx = (-1)^p \frac{\Gamma(p+1)}{(s+1)^{p+1}}, \quad \operatorname{Re} s > -1, \operatorname{Re} p > -1. \quad (40)$$

Inserting this formula into (39) and setting  $n-1$  instead of  $s$ , yields (28).

*Proof of formula (29)*

Putting in (38)  $x = x_1 x_2 \cdots x_{p+1}$  and integrating over the volume  $[0, 1]^{p+1}$ , where  $p \in \mathbb{N}$ , on the one hand, we have

$$\underbrace{\int_0^1 \cdots \int_0^1}_{(p+1)\text{-fold}} \sum_{n=1}^{\infty} |G_n| (x_1 x_2 \cdots x_{p+1})^{n-1} dx_1 \cdots dx_{p+1} = \sum_{n=1}^{\infty} \frac{|G_n|}{n^{p+1}}. \quad (41)$$

On the other hand

$$\int_0^1 \left\{ \frac{1}{\ln(1-xy)} + \frac{1}{xy} \right\} dx = -\frac{\operatorname{li}(1-y) - \gamma - \ln y}{y}.$$

Taking instead of  $y$  the product  $x_1 x_2 \cdots x_p$  and setting  $x = x_{p+1}$ , and then integrating  $p$  times over the unit hypercube and equating the result with (41) yields (29).

*Proof of formulas (30) and (31)*

Writing in the generating equation (10)  $-x$  instead of  $z$ , multiplying it by  $\ln^m x/x$  and integrating over the unit interval, we obtain the following relation<sup>16</sup>

$$\Omega(k, m) = (-1)^{m+k} m! k! \sum_{n=k}^{\infty} \frac{|S_1(n, k)|}{n! n^{m+1}},$$

where

$$\Omega(k, m) := \int_0^1 \frac{\ln^k(1-x) \ln^m x}{x} dx, \quad k, m \in \mathbb{N}.$$

By integration by parts, it may be readily shown that

$$\Omega(k, m) = \frac{k}{m+1} \Omega(m+1, k-1),$$

and thus, we deduce the duality formula:

$$\sum_{n=k}^{\infty} \frac{|S_1(n, k)|}{n! n^{m+1}} = \sum_{n=m+1}^{\infty} \frac{|S_1(n, m+1)|}{n! n^k}.$$

---

<sup>16</sup>See also [40, Theorem 2.7].

Furthermore, since  $1 - x = e^{\ln(1-x)}$  for  $x < 1$ , the Maclaurin series expansion of  $e^z$  with  $z = \ln(1 - x)$  gives

$$x = 1 - \sum_{k=0}^{\infty} \frac{\ln^k(1-x)}{k!},$$

or equivalently,

$$x + \ln(1-x) = - \sum_{k=1}^{\infty} \frac{\ln^{k+1}(1-x)}{(k+1)!},$$

from which we deduce

$$\begin{aligned} \int_0^1 \left\{ \frac{1}{\ln(1-x)} + \frac{1}{x} \right\} \ln^m x \, dx &= - \int_0^1 \sum_{k=1}^{\infty} \frac{\ln^{k+1}(1-x)}{(k+1)!} \cdot \frac{\ln^m x}{\ln(1-x)} \cdot \frac{dx}{x} \\ &= - \sum_{k=1}^{\infty} \frac{\Omega(k, m)}{(k+1)!} = (-1)^m m! \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{(k+1)} \sum_{n=m+1}^{\infty} \frac{|S_1(n, m+1)|}{n! n^k}, \end{aligned}$$

which is identical with (30) if setting  $m = p$ . Moreover, it is well known that

$$\frac{|S_1(n+1, m+1)|}{n!} = P_m(H_n^{(1)}, -H_n^{(2)}, \dots, (-1)^{m-1} H_n^{(m)}),$$

see [18, p. 217], [36, p. 1395], [30, p. 425, Eq. (43)], [6, Eq. (16)], which immediately gives (31) and completes the proof.

*Proof of formula (32)*

This formula straightforwardly follows from the fact that  $\kappa_p = F_p(1)$ , see [13, p. 307, 318 *et seq.*], where  $F_p(s)$  is the special function defined by

$$F_p(s) := \sum_{n=1}^{\infty} \frac{|G_n|}{n^p} D\left(\frac{1}{x^s}\right)(n) = \sum_{n=0}^{\infty} \frac{|G_{n+1}|}{(n+1)^p} \sum_{k=0}^n (-1)^k \binom{n}{k} (k+1)^{-s}.$$

*Proof of formulas (36) and (37)*

These formulas immediately follow from [12, Eqs. (3.21) and (3.23)] and (35).

## Acknowledgments

The authors warmly thank the referee for his valuable comments. They are also grateful to Vladimir V. Reshetnikov for his kind help and useful remarks.

## References

- [1] Correspondance d'Hermitte et de Stieltjes. Vol. 1 and 2, Gauthier-Villars, Paris, 1905.
- [2] Collected papers of Srinivasa Ramanujan, Cambridge, 1927.
- [3] T. M. Apostol, Formulas for higher derivatives of the Riemann zeta function, *Mathematics of Computation*, vol. 44, no. 169, pp. 223–232 (1985).
- [4] B. C. Berndt, *Ramanujan's Notebooks, Part I*. Springer-Verlag, 1985.

- [5] *B. C. Berndt*, On the Hurwitz Zeta–function, *Rocky Mountain Journal of Mathematics*, vol. 2, no. 1, pp. 151–157 (1972).
- [6] *Ia. V. Blagouchine*, Expansions of generalized Euler’s constants into the series of polynomials in  $\pi^{-2}$  and into the formal enveloping series with rational coefficients only, *Journal of Number Theory* (Elsevier), vol. 158, pp. 365–396 and vol. 173, pp. 631–632, arXiv:1501.00740 (2016).
- [7] *Ia. V. Blagouchine*, Three notes on Ser’s and Hasse’s representations for the zeta-functions, arXiv:1606.02044 (2016).
- [8] *Ia. V. Blagouchine*, Two series expansions for the logarithm of the gamma function involving Stirling numbers and containing only rational coefficients for certain arguments related to  $\pi^{-1}$ , *Journal of Mathematical Analysis and Applications* (Elsevier), vol. 442, pp. 404–434, arXiv:1408.3902 (2016).
- [9] *Ia. V. Blagouchine*, A note on some recent results for the Bernoulli numbers of the second kind, *J. Integer Seq.*, vol. 20, no. 3, Article 17.3.8, pp. 1-7, 2017, arXiv:1612.03292 (2017).
- [10] *W. E. Briggs*, The irrationality of  $\gamma$  or of sets of similar constants, *Vid. Selsk. Forh. (Trondheim)*, vol. 34, pp. 25–28 (1961).
- [11] *W. E. Briggs* and *S. Chowla*, The power series coefficients of  $\zeta(s)$ , *The American Mathematical Monthly*, vol. 62, pp. 323–325 (1955).
- [12] *B. Candelpergher*, Ramanujan summation of divergent series, *Lecture Notes in Mathematics series*, Vol. 2185, Springer, 2017, hal-01150208.
- [13] *B. Candelpergher* and *M.-A. Coppo*, A new class of identities involving Cauchy numbers, harmonic numbers and zeta values, *The Ramanujan Journal*, vol. 27, pp. 305–328 (2012).
- [14] *B. Candelpergher* and *M.-A. Coppo*, Le produit harmonique des suites, *Enseign. Math.*, vol. 59, pp. 39–72 (2013).
- [15] *B. Candelpergher*, *M.-A. Coppo* and *E. Delabaere*, La sommation de Ramanujan, *Enseign. Math.*, vol. 43, pp. 93–132 (1997).
- [16] *B. Candelpergher*, *H. Gadiyar* and *R. Padma*, Ramanujan summation and the exponential generating function  $\sum_{k=0}^{\infty} \frac{z^k}{k!} \zeta'(-k)$ , *The Ramanujan Journal*, vol. 21, pp. 99–122 (2010).
- [17] *M. W. Coffey*, Certain logarithmic integrals, including solution of monthly problem #tbd, zeta values, and expressions for the Stieltjes constants, arXiv:1201.3393v1 (2012).
- [18] *L. Comtet*, *Advanced Combinatorics. The art of Finite and Infinite Expansions* (revised and enlarged edition), D. Reidel Publishing Company, Dordrecht, Holland, 1974.
- [19] *D. F. Connon*, Some possible approaches to the Riemann hypothesis via the Li/Keiper constants, arXiv:1002.3484 (2010).
- [20] *M.-A. Coppo*, Nouvelles expressions des constantes de Stieltjes, *Expositiones Mathematicae*, vol. 17, pp. 349–358 (1999).

- [21] *M.-A. Coppo and P. T. Young*, On shifted Mascheroni series and hyperharmonic numbers, *Journal of Number Theory*, vol. 169, pp. 1–20 (2016).
- [22] *K. Dilcher*, Generalized Euler constants for arithmetical progressions, *Mathematics of Computation*, vol. 59, pp. 259–282 (1992).
- [23] *S. R. Finch*, *Mathematical constants*, Cambridge University Press, USA, 2003.
- [24] *J. Franel*, Note n° 245, *L'Intermédiaire des mathématiciens*, tome II, pp. 153–154 (1895).
- [25] *J. P. Gram*, Note sur le calcul de la fonction  $\zeta(s)$  de Riemann, *Oversigt. K. Danske Vidensk. (Selskab Forhandlingar)*, pp. 305–308 (1895).
- [26] *G. H. Hardy*, Note on Dr. Vacca's series for  $\gamma$ , *The Quarterly journal of pure and applied mathematics*, vol. 43, pp. 215–216 (1912).
- [27] *M. I. Israilov*, On the Laurent decomposition of Riemann's zeta function [in Russian], *Trudy Mat. Inst. Akad. Nauk. SSSR*, vol. 158, pp. 98–103 (1981).
- [28] *J. L. W. V. Jensen*, Sur la fonction  $\zeta(s)$  de Riemann, *Comptes-rendus de l'Académie des sciences*, tome 104, pp. 1156–1159 (1887).
- [29] *J. L. W. V. Jensen*, Note n° 245. Deuxième réponse. Remarques relatives aux réponses de MM. Franel et Kluyver, *L'Intermédiaire des mathématiciens*, tome II, pp. 346–347 (1895).
- [30] *V. Kowalenko*, Properties and applications of the reciprocal logarithm numbers, *Acta Applicandæ Mathematicæ*, vol. 109, pp. 413–437 (2010).
- [31] *D. H. Lehmer*, Euler constants for arithmetical progressions, *Acta Arithmetica*, vol. 27, pp. 125–142 (1975).
- [32] *D. H. Lehmer*, The sum of like powers of the zeros of the Riemann zeta function, *Mathematics of Computation*, vol. 50, no. 181, pp. 265–273 (1988).
- [33] *J. J. Y. Liang and J. Todd*, The Stieltjes constants, *Journal of Research of the National Bureau of Standards—Mathematical Sciences*, vol. 76B, nos. 3–4, pp. 161–178 (1972).
- [34] *Z. Nan-You and K. S. Williams*, Some results on the generalized Stieltjes constant, *Analysis*, vol. 14, pp. 147–162 (1994).
- [35] *T. H. Pilehrood and K. H. Pilehrood*, Criteria for irrationality of generalized Euler's constant, *Journal of Number Theory*, vol. 108, pp. 169–185 (2004).
- [36] *L.-C. Shen*, Remarks on some integrals and series involving the Stirling numbers and  $\zeta(n)$ , *The Transactions of the American Mathematical Society*, vol. 347, no. 4, pp. 1391–1399 (1995).
- [37] *R. Sitaramachandrarao*, Maclaurin coefficients of the Riemann zeta function, *Abstracts of Papers Presented to the American Mathematical Society*, vol. 7, no. 4, p. 280, \*86T-11-236 (1986).
- [38] *T. Tasaka*, Note on the generalized Euler constants, *Mathematical Journal of Okayama University*, vol. 36, pp. 29–34 (1994).

- [39] *L. Xia*, The parameterized-Euler-constant function  $\gamma_a(z)$ , *Journal of Number Theory*, vol. 133, no. 1, pp. 1–11 (2013).
- [40] *C. Xu, Y. Yan and Z. Shi*, Euler sums and integrals of polylogarithm functions, *Journal of Number Theory*, vol. 165, pp. 84–108 (2016).
- [41] *P. T. Young*, A 2-adic formula for Bernoulli numbers of the second kind and for the Nörlund numbers, *Journal of Number Theory*, vol. 128, pp. 2951–2962 (2008).
- [42] *P. T. Young*, Rational series for multiple zeta and log gamma functions, *Journal of Number Theory*, vol. 133, pp. 3995–4009 (2013).