



HAL
open science

New identities involving Cauchy numbers, harmonic numbers and zeta values

Marc-Antoine Coppo

► **To cite this version:**

Marc-Antoine Coppo. New identities involving Cauchy numbers, harmonic numbers and zeta values. 2021. hal-03131735v2

HAL Id: hal-03131735

<https://hal.univ-cotedazur.fr/hal-03131735v2>

Preprint submitted on 10 Feb 2021 (v2), last revised 3 Sep 2021 (v10)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

New identities involving Cauchy numbers, harmonic numbers and zeta values

Marc-Antoine Coppo*

Université Côte d'Azur, CNRS, LJAD (UMR 7351), Nice, France

Abstract In this article, we present a number of series identities linking together Cauchy numbers (also known as Bernoulli numbers of the second kind), the values of zeta and its derivative, and a generalization of Roman harmonic numbers.

Keywords: Cauchy numbers, Roman harmonic numbers, binomial identities, series with special numbers, zeta values.

Introduction

Several years ago, we proposed in [3] a method based on the Ramanujan summation of series which enabled us to generate a large number of identities involving Cauchy numbers, harmonic numbers and zeta values. By means of new formulas recently proved in our last paper [6], we are in a position to complete these results by showing new identities of the same kind (see Propositions 1 and 2). In order to do this, we make use of a rather natural generalization of Roman harmonic numbers introduced in [4] (see Definition 2). A notable novelty is the appearance in our formulas of certain alternating series involving the values of zeta that have recently been considered in [5] and thoroughly studied in [7] (see Definition 3).

1 Preliminaries : reminder of the main definitions and results

We first recall a number of definitions and results that appear in previous work.

*Corresponding author. *Email address:* coppo@unice.fr

1.1 Harmonic numbers and harmonic sums

Definition 1. The generalized harmonic numbers $H_n^{(r)}$ are defined for all natural numbers n and r by

$$H_0^{(r)} = 0 \quad \text{and} \quad H_n^{(r)} = \sum_{j=1}^n \frac{1}{j^r} \quad \text{for } n \geq 1. \quad (1)$$

For $r = 1$, they reduce to classical harmonic numbers $H_n = H_n^{(1)}$. The sums

$$\mathcal{S}_{p,q} = \sum_{n=1}^{\infty} \frac{H_n^{(p)}}{n^q}$$

for positive integers p and q with $q \geq 2$ are called linear Euler sums. We have the classical Euler's formula (cf. [6, 9]):

$$2\mathcal{S}_{1,p} = (p+2)\zeta(p+1) - \sum_{j=1}^{p-2} \zeta(p-j)\zeta(j+1) \quad (p \geq 2).$$

Definition 2 (cf. [4]). The harmonic sums $H_{n,k}^{(r)}$ are defined for all natural numbers n, r and k with $n \geq 1, r \geq 1$ and $k \geq 0$ by

$$H_{n,0}^{(r)} = \frac{1}{n^{r-1}} \quad \text{and} \quad H_{n,k}^{(r)} = \sum_{n \geq j_1 \geq \dots \geq j_k \geq 1} \frac{1}{j_1 j_2 \dots j_k^r} \quad \text{for } k \geq 1. \quad (2)$$

In particular, for $k = 1$, they reduce to classical generalized harmonic numbers

$$H_{n,1}^{(r)} = H_n^{(r)},$$

and for $k = 2$, we have

$$H_{n,2}^{(r)} = \sum_{j=1}^n \frac{H_j^{(r)}}{j}.$$

Remark 1. For $r = 1$, the harmonic sums $H_{n,k} := H_{n,k}^{(1)}$ are nothing else than the ordinary Roman harmonic numbers $c_n^{(k)}$ (cf. [8]). In particular, we have

$$H_{n,0} = 1, \quad H_{n,1} = H_n, \quad H_{n,2} = \sum_{j=1}^n \frac{H_j}{j}, \quad \text{etc.}$$

In the general case, it can be shown (cf. [3, Eq. (18)] and [8, Eq. (29)]) that

$$H_{n,k} = P_k(H_n, \dots, H_n^{(k)})$$

where $P_k(x_1, \dots, x_k)$ are the modified Bell polynomials (cf. [3, Definition 2]).

Example 1.

$$H_{n,2} = \sum_{j=1}^n \frac{H_j}{j} = \frac{1}{2}(H_n)^2 + \frac{1}{2}H_n^{(2)}. \quad (3)$$

The harmonic sums $H_{n,k}^{(r)}$ verify the following identity (cf. [4, Eq. (4.7)]):

$$H_{n,k}^{(r)} = \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} \frac{H_{j,r-1}}{j^k} \quad (k \geq 1). \quad (4)$$

Example 2. For $r = 1$, formula (4) reduces to

$$H_{n,k} = \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} \frac{1}{j^k} \quad (5)$$

which is a classical property of Roman harmonic numbers (cf. [8, Eq. (20)]), and for $r = 2$, it is written as follows:

$$H_{n,k}^{(2)} = \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} \frac{H_j}{j^k}. \quad (6)$$

By inverse binomial transform¹, the identity (4) also admits a kind of reciprocal:

$$\frac{H_{n,r-1}}{n^k} = \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} H_{j,k}^{(r)} \quad (k \geq 1). \quad (7)$$

1.2 Alternating series with zeta values

Definition 3 (cf. [5, 6, 7]). the conditionally convergent series τ_p are defined for any positive integer p by

$$\tau_p = \sum_{k=1}^{\infty} (-1)^{k+p} \frac{\zeta(k+p)}{k}. \quad (8)$$

For $p \geq 2$, they verify the relation

$$\sum_{n=1}^{\infty} \frac{\ln(n+1)}{n^p} = -\zeta'(p) - (-1)^p \tau_p, \quad (9)$$

and for $p = 1$, we have

$$\tau_1 = \int_0^1 \frac{\psi(x+1) + \gamma}{x} dx \quad (10)$$

where ψ is the digamma function.

1. If $b(n) = \frac{1}{n} \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} a(j)$, then $a(n) = \frac{1}{n} \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} j b(j)$ (cf. [4, Definition 5 and Corollary 1]).

Remark 2. The series τ_p defined above are linked to the series σ_p introduced and studied in [7] by the elementary relation

$$\tau_p = (-1)^{p-1} \sigma_p.$$

1.3 Cauchy numbers

Definition 4 (cf. [1, 2, 3]). The *non-alternating Cauchy numbers* λ_n are defined for all integers $n \geq 1$ by the recurrence relation

$$\sum_{k=1}^{n-1} \frac{\lambda_k}{k!(n-k)} = \frac{1}{n} \quad \text{for } n \geq 2.$$

The first values are

$$\lambda_1 = \frac{1}{2}, \lambda_2 = \frac{1}{6}, \lambda_3 = \frac{1}{4}, \lambda_4 = \frac{19}{30}, \lambda_5 = \frac{9}{4}, \lambda_6 = \frac{863}{84}, \text{ etc.}$$

Remark 3. The λ_n are linked to the classical Cauchy numbers denoted c_n in [1] by the relation

$$\lambda_n = (-1)^{n-1} c_n = (-1)^{n-1} \int_0^1 x(x-1) \cdots (x-n+1) dx \quad \text{for } n \geq 1.$$

We recall the transformation formula that links Cauchy numbers to the Ramanujan summation of series (cf. [2, Theorem 18]) : if a is a function analytic in the half-plane $P = \{\text{Re}(z) > 0\}$ such that there exists a constant $C > 0$ with

$$|a(z)| < C 2^{|z|} \quad \text{for all } z \in P,$$

then

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} \sum_{j=1}^n (-1)^{j-1} \binom{n}{j} j a(j) = \sum_{n \geq 1}^{\mathcal{R}} a(n) \quad (11)$$

where $\sum_{n \geq 1}^{\mathcal{R}}$ denotes the \mathcal{R} -sum of the series (i.e. the sum of the series in the sense of Ramanujan's summation method).

2 Series with Cauchy numbers and harmonic sums

Proposition 1. Let p be an integer with $p \geq 1$, then

a)

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_{n,p-1} = \begin{cases} \gamma & \text{for } p = 1 \\ \zeta(p) - \frac{1}{p-1} & \text{for } p > 1 \end{cases} \quad (12)$$

where $\gamma = -\psi(1)$ is Euler's constant.

b)

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_{n,p-1}^{(2)} = \begin{cases} \frac{1}{2}\gamma^2 - \frac{1}{2}\zeta(2) + \gamma_1 + \tau_1 & \text{for } p = 1 \\ \mathcal{S}_{1,p} - \sum_{j=1}^{p-2} \frac{(-1)^{p-j}}{j} \zeta(p-j) + (-1)^p \zeta'(p) + \tau_p & \text{for } p > 1 \end{cases} \quad (13)$$

where γ_1 is the first Stieltjes constant.

Proof. It follows from formulas (4) and (11) that

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_{n,k}^{(r)} = \sum_{n \geq 1}^{\mathcal{R}} \frac{H_{n,r-1}}{n^{k+1}} \quad (k \geq 0, r \geq 1).$$

Applying the above formula with $k = p - 1$, we obtain

a) for $r = 1$,

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_{n,p-1} = \sum_{n \geq 1}^{\mathcal{R}} \frac{1}{n^p},$$

b) for $r = 2$,

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_{n,p-1}^{(2)} = \sum_{n \geq 1}^{\mathcal{R}} \frac{H_n}{n^p}.$$

Hence, formula (12) results from [2, Eqs. (1.22) and (1.24)] while formula (13) results from [6, Proposition 1 and Eq. (13)]. \square

Example 3. a) For $p = 1$, formulas (12)–(13) translate into

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} = \gamma, \quad (14)$$

and

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n^2} = \frac{1}{2}\gamma^2 - \frac{1}{2}\zeta(2) + \gamma_1 + \tau_1. \quad (15)$$

b) For $p = 2$, formulas (12)–(13) translate into

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_n = \zeta(2) - 1, \quad (16)$$

and

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n!n} H_n^{(2)} = 2\zeta(3) + \zeta'(2) + \tau_2, \quad (17)$$

Remark 4. Formulas (14) and (16) are well-known classical series representations for γ and $\zeta(2)$ (cf. [1, 2, 3]).

Example 4. For $p = 3$, formulas (12)–(13) translate into

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} \sum_{j=1}^n \frac{H_j}{j} = \zeta(3) - \frac{1}{2}, \quad (18)$$

and

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} \sum_{j=1}^n \frac{H_j^{(2)}}{j} = \frac{5}{4} \zeta(4) - \zeta(2) - \zeta'(3) + \tau_3. \quad (19)$$

Remark 5. a) Thanks to the identity (3), formula (18) can be rewritten under the following form:

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} (H_n)^2 + \sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} H_n^{(2)} = 2\zeta(3) - 1. \quad (20)$$

Subtracting (17) from (20) we deduce yet another interesting identity:

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} (H_n)^2 = -\zeta'(2) - \tau_2 - 1. \quad (21)$$

b) By means of [6, Eq. (14)] we can also easily prove the following formula dual of (19):

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} \sum_{j=1}^n \frac{H_j}{j^2} = \frac{7}{4} \zeta(4) + \zeta(2) + 2\zeta'(3) - 2\tau_3 - 1. \quad (22)$$

Hence, by adding (19) and (22) we obtain

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} \sum_{j=1}^n \left\{ \frac{H_j}{j^2} + \frac{H_j^{(2)}}{j} \right\} = 3\zeta(4) + \zeta'(3) - \tau_3 - 1. \quad (23)$$

Proposition 2. Let p be an integer with $p \geq 2$, then

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n^2} H_{n,p-1} = \gamma \zeta(p) + \zeta(p+1) - \mathcal{S}_{1,p} - \zeta'(p) - (-1)^p \tau_p - \sigma_p \quad (24)$$

with $\sigma_2 = 1$ and

$$\sigma_p = \frac{1 + (-1)^p}{p} + \sum_{j=1}^{p-2} (-1)^j \zeta(p-j) \left[\frac{(j-1)!(p-1-j)!}{(p-1)!} - \frac{1}{j} \right] \quad \text{for } p \geq 3.$$

Proof. It follows from formulas (7) and (11) that

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n^{k+1}} H_{n,r-1} = \sum_{n \geq 1}^{\mathcal{R}} \frac{H_{n,k}^{(r)}}{n} \quad (k \geq 0, r \geq 1).$$

The above formula applied for $k = 1$ and $r = p$ gives

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n^2} H_{n,p-1} = \sum_{n \geq 1}^{\mathcal{R}} \frac{H_n^{(p)}}{n}.$$

Formula (24) then results from [6, Proposition 2]. □

Example 5. Formula (24) applied with $p = 2$ gives the following identity:

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n^2} H_n = \gamma \zeta(2) - \zeta(3) - \zeta'(2) - \tau_2 - 1. \quad (25)$$

Moreover, subtracting (25) from (21) and replacing $H_n - \frac{1}{n}$ by H_{n-1} leads to the nice expression

$$\sum_{n=1}^{\infty} \frac{\lambda_n}{n! n} H_n H_{n-1} = \zeta(3) - \gamma \zeta(2). \quad (26)$$

Example 6. Formula (24) applied with $p = 3$ gives the relation:

$$\frac{1}{2} \sum_{n=1}^{\infty} \frac{\lambda_n}{n! n^2} \left\{ (H_n)^2 + H_n^{(2)} \right\} = \gamma \zeta(3) - \frac{1}{4} \zeta(4) - \frac{1}{2} \zeta(2) - \zeta'(3) + \tau_3. \quad (27)$$

References

- [1] K. N. Boyadzhiev, New series identities with Cauchy, Stirling, and harmonic numbers, and Laguerre polynomials, 2020, arXiv:1911.00186v2.
- [2] B. Candelpergher, *Ramanujan Summation of Divergent Series*, Lecture Notes in Math. 2185, Springer, 2017.
- [3] B. Candelpergher, M-A. Coppo, A new class of identities involving Cauchy numbers, harmonic numbers and zeta values, *Ramanujan J.* 27 (2012), 305–328.
- [4] B. Candelpergher, M-A. Coppo, Le produit harmonique des suites, *L'enseignement Mathématique*, 59 (2013), 39–72.
- [5] B. Candelpergher, M-A. Coppo, Laurent expansion of harmonic zeta functions, *J. Math. Anal. App.* 491 (2020), Article 124309.

- [6] M-A. Coppo, B. Candelpergher, New reflection formulas for Euler sums, 2021. hal-02921391v3.
- [7] A. Dil, K. N. Boyadzhiev, and I. A. Aliev, On values of the Riemann zeta function at positive integers, *Lithuanian Math. J.* **60** (2020), 9–24.
- [8] J. Sesma, The Roman harmonic numbers revisited, *J. Number Theory* **180** (2017), 544–565.
- [9] W. Wang, Y. Lyu, Euler sums and Stirling sums, *J. Number Theory* **185** (2018), 160–193.