



**HAL**  
open science

# Generalized Glaisher-Kinkelin constants and Blagouchine's integrals

Marc-Antoine Coppo

► **To cite this version:**

Marc-Antoine Coppo. Generalized Glaisher-Kinkelin constants and Blagouchine's integrals. 2022.  
hal-03197403v17

**HAL Id: hal-03197403**

**<https://hal.univ-cotedazur.fr/hal-03197403v17>**

Preprint submitted on 2 Mar 2022 (v17), last revised 3 Jun 2024 (v23)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Generalized Glaisher-Kinkelin constants and Blagouchine's integrals

Marc-Antoine Coppo

*Université Côte d'Azur, CNRS, LJAD (UMR 7351), Nice, France*

*Courriel : coppo@unice.fr*

**Abstract.** The main purpose of this short article is to highlight the existence of a close connection between a family of complex integrals introduced by Blagouchine and some notable mathematical constants, namely the generalized Glaisher-Kinkelin constants (also known as the Bendersky constants) which occur quite naturally in number theory and analysis.

**Keywords.** Glaisher-Kinkelin constants, Bendersky generalized gamma function, infinite series with zeta values, complex integration.

## 1 Introduction

The aim of this article is to establish a link between the family of complex integrals  $\{\mathcal{J}_{k,p}\}$  (for integers  $k \geq 0$  and  $p \geq 1$  with  $p$  odd) defined by

$$\mathcal{J}_{k,p} = \int_{-\infty}^{+\infty} \frac{\zeta(\frac{p}{2} + ix)}{(2k + p + 2ix) \cosh(\pi x)} dx,$$

and a sequence of notable mathematical constants, namely the generalized Glaisher-Kinkelin constants (also known as the Bendersky constants) which occur quite naturally in analysis and number theory (see Definition 1). Blagouchine [3] introduced these integrals in the cases  $p = 1$  and  $p = 3$ . To show this close connection, we make use of a relation between the integral  $\mathcal{J}_{k,p}$  and the alternating series

$$\sum_{n=N_p}^{\infty} (-1)^n \frac{\zeta(n)}{n+k} \quad \text{with } N_p = \max(2, \frac{p+1}{2})$$

which can be deduced from the residue theorem (see Proposition 1). These series have been studied in detail in our previous article [7]. In particular, this enables us

to give a general expression of the integrals  $\mathcal{J}_{k,1}$  and  $\mathcal{J}_{k,3}$  in terms of the generalized Glaisher-Kinkelin constants for all positive integers  $k$  (see Corollary 1).

Recently, Candelpergher [5] has generalized the relation between  $\mathcal{J}_{k,p}$  and the alternating series with zeta values by using another method (see Proposition 2). As a corollary, this allows us to give a representation of the Dirichlet series

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n^s}$$

which is valid for any complex number  $s$  with  $\operatorname{Re}(s) > \frac{1}{2}$  (see Corollary 2).

## 2 Generalized Glaisher-Kinkelin constants

**Definition 1** ([1, 9, 11]). For any integer  $k \geq 0$ , the constant  $A_k$  are usually defined by

$$\begin{aligned} \ln A_0 &= \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N \ln n - \left( N + \frac{1}{2} \right) \ln N + N \right\}, \\ \ln A_1 &= \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N n \ln n - \left( \frac{N^2}{2} + \frac{N}{2} + \frac{1}{12} \right) \ln N + \frac{N^2}{4} \right\}, \\ \ln A_2 &= \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N n^2 \ln n - \left( \frac{N^3}{3} + \frac{N^2}{2} + \frac{N}{6} \right) \ln N + \frac{N^3}{9} - \frac{N}{12} \right\}, \end{aligned}$$

and more generally

$$\ln A_k = \lim_{N \rightarrow \infty} \left\{ \sum_{n=1}^N n^k \ln n - P_k(N) \ln N + Q_k(N) \right\},$$

where  $P_k$  and  $Q_k$  are polynomials of degree  $k+1$  that can be explicitly computed (see e.g. [11, Eq. (1.1)]). The numbers  $A_k$  for  $k = 0, 1, 2, \dots$  are called the *generalized Glaisher-Kinkelin constants* (sometimes called the *Bendersky constants*). Adamchik [1, Proposition 4] has given an alternative expression of the constants  $A_k$  in terms of the derivatives of the Riemann zeta function. More precisely, we have the expression

$$\ln A_k = \frac{H_k B_{k+1}}{k+1} - \zeta'(-k), \quad (1)$$

where  $H_n$  are the harmonic numbers (with the usual convention  $H_0 = 0$ ) and  $B_n$  are the Bernoulli numbers.

**Example 1.** The constant  $A_0$  is the Stirling constant:

$$A_0 = \exp(-\zeta'(0)) = \sqrt{2\pi},$$

the constant  $A_1$  is the classical Glaisher-Kinkelin constant:

$$A_1 = \exp \left\{ \frac{1}{12} - \zeta'(-1) \right\} = \exp \left\{ \frac{1}{12}(\gamma + \ln 2\pi) - \frac{\zeta'(2)}{2\pi^2} \right\},$$

$\gamma$  being the Euler constant, and

$$A_2 = \exp(-\zeta'(-2)) = \exp \left\{ \frac{\zeta(3)}{4\pi^2} \right\}.$$

More generally, the following relations:

$$\zeta'(-2k) = (-1)^k \frac{(2k)!}{2(2\pi)^{2k}} \zeta(2k+1) \quad (k \geq 1),$$

and

$$\zeta'(1-2k) = (-1)^{k+1} \frac{(2k)!}{k(2\pi)^{2k}} \zeta'(2k) + \frac{B_{2k}}{2k} (H_{2k-1} - \gamma - \ln 2\pi) \quad (k \geq 1),$$

which are easily derived by differentiation of the functional equation for the zeta function, enable to deduce from Adamchik's formula (1) the expressions

$$\ln A_{2k-1} = (-1)^k \frac{(2k)!}{k(2\pi)^{2k}} \zeta'(2k) + \frac{B_{2k}}{2k} (\gamma + \ln 2\pi) \quad (k \geq 1), \quad (2)$$

and

$$\ln A_{2k} = (-1)^{k+1} \frac{(2k)!}{2(2\pi)^{2k}} \zeta(2k+1) \quad (k \geq 1). \quad (3)$$

*Remark 1.* Bendersky [2] introduced for the first time the sequence of numbers  $A_k$  without any consideration of their relation with the  $\zeta$ -function. From the point of view of the summation of divergent series, the constants  $A_k$  can be interpreted as follows: let  $\sum_{n \geq 1}^{\mathcal{R}} n^k \ln n$  denotes the  $\mathcal{R}$ -sum of the divergent series  $\sum_{n \geq 1} n^k \ln n$  (i.e. the sum of the series in the sense of Ramanujan's summation method [4]), then, for any integer  $k \geq 0$ , we have (see [4, p. 68] and [2, p. 280]):

$$\begin{aligned} \sum_{n \geq 1}^{\mathcal{R}} n^k \ln n &= -\zeta'(-k) - \frac{1}{(k+1)^2} \\ &= \ln A_k - \frac{H_k B_{k+1}}{k+1} - \frac{1}{(k+1)^2} \\ &= \int_0^1 \ln \Gamma_k(x+1) dx, \end{aligned}$$

where  $\Gamma_k$  is the Bendersky generalized gamma function [2, p. 279]. This function verifies

$$\Gamma_k(n+1) = 1^{1^k} 2^{2^k} \cdots n^{n^k} \quad \text{for any integer } n \geq 1.$$

In particular, we have  $\Gamma_0 = \Gamma$ , and  $\Gamma_1 = K$ , where  $K$  stands for the Kinkelin-Bendersky hyperfactorial function which can be defined (see e.g. [10, Definition 3]) by the relation

$$\ln K(x) = \frac{x^2 - x}{2} - \frac{x}{2} \ln 2\pi + \int_0^x \ln \Gamma(u) du \quad (x \geq 0).$$

Unaware of Bendersky's work and following an idea of Milnor, Kurokawa and Ochiai [8, Theorem 2] have given an expression of the function  $\Gamma_k$  in terms of the derivative of the Hurwitz zeta function  $\zeta(s, x)$  at  $s = -k$ . Precisely, they show that

$$\ln \Gamma_k(x) = \zeta'(-k, x) - \zeta'(-k) \quad \text{for } x > 0 \text{ and } k \geq 0,$$

a formula which generalizes a classical formula for  $\Gamma$  (see [6, Definition 9.6.13]) and can be seen as the analogue of Adamchik's formula for  $A_k$ .

### 3 Blagouchine's integrals and series with zeta values

**Definition 2.** For any non-negative integer  $k$  and positive odd integer  $p$ , the integral  $\mathcal{J}_{k,p}$  is defined by

$$\mathcal{J}_{k,p} = \int_{-\infty}^{+\infty} \frac{\zeta(\frac{p}{2} + ix)}{(2k + p + 2ix) \cosh(\pi x)} dx.$$

**Proposition 1.** We have the following relations:

$$\mathcal{J}_{k,1} = \frac{\gamma}{k+1} - \frac{1}{(k+1)^2} - \sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n+k}, \quad (4)$$

and

$$\mathcal{J}_{k,p} = (-1)^{\frac{p+1}{2}} \sum_{n=\frac{p+1}{2}}^{\infty} (-1)^n \frac{\zeta(n)}{n+k} \quad \text{for } p = 3, 5, 7, \dots \quad (5)$$

*Proof.* For  $k \geq 0$ , let us consider the function

$$f_k(z) = \frac{\zeta(z)}{(k+z) \sin(\pi z)}.$$

The function  $f_k$  has poles at integers  $n \in \mathbb{Z}$ . For  $n \geq 2$ , the residue of  $f_k$  at  $z = n$  is

$$\operatorname{Res}(f_k; n) = \frac{(-1)^n \zeta(n)}{(n+k)\pi}.$$

For  $n = 1$ ,  $f_k$  has a double pole and

$$\operatorname{Res}(f_k; 1) = -\frac{1}{\pi} \left( \frac{\gamma}{k+1} - \frac{1}{(k+1)^2} \right).$$

Applying the residue theorem, we get

$$-\frac{1}{2i\pi} \int_{\operatorname{Re}(z)=p/2} f_k(z) dz = \sum_{n > \frac{p}{2}} \operatorname{Res}(f_k; n).$$

This leads to formulas (4) and (5).  $\square$

**Corollary 1.** We have

$$\mathcal{J}_{k,1} = \sum_{j=0}^{k-1} (-1)^j \binom{k}{j} \ln A_j - \frac{1}{k} - \frac{1}{(k+1)^2} \quad (k \geq 1), \quad (6)$$

and

$$\mathcal{J}_{k,3} = \frac{\gamma}{k+1} - \frac{1}{(k+1)^2} - \mathcal{J}_{k,1} = \frac{\gamma}{k+1} + \frac{1}{k} - \sum_{j=0}^{k-1} (-1)^j \binom{k}{j} \ln A_j \quad (k \geq 1). \quad (7)$$

*Proof.* From [7, Proposition 1], we have

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n+k} = \frac{\gamma}{k+1} + \sum_{j=0}^{k-1} (-1)^j \binom{k}{j} \zeta'(-j) + \frac{1}{k} + \sum_{j=0}^{k-1} \binom{k}{j} \frac{B_{j+1} H_j}{j+1} \quad (k \geq 1).$$

By Adamchik's formula (1), this expression may be rewritten as follows:

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n+k} = \frac{\gamma}{k+1} + \frac{1}{k} - \sum_{j=0}^{k-1} (-1)^j \binom{k}{j} \ln A_j \quad (k \geq 1).$$

Then, using the relation (4) and (5), we get the expressions (6) and (7).  $\square$

**Example 2.** For  $p = 1$  and the first values of  $k$ , we have

$$\begin{aligned} \mathcal{J}_{0,1} &= -1, \\ \mathcal{J}_{1,1} &= \frac{1}{2} \ln 2\pi - \frac{5}{4}, \\ \mathcal{J}_{2,1} &= \frac{1}{3} \ln 2\pi - \frac{1}{6} \gamma - \frac{11}{18} + \frac{\zeta'(2)}{\pi^2}, \\ \mathcal{J}_{3,1} &= \frac{1}{4} \ln 2\pi - \frac{1}{4} \gamma - \frac{19}{48} + \frac{3}{2\pi^2} \zeta'(2) + \frac{3}{4\pi^2} \zeta(3). \end{aligned}$$

## 4 Further generalization

Using a Fourier transform method, Candelpergher [5, Eq. (7)] recently proved the following beautiful relation which is a natural generalization of (4).

**Proposition 2.** for  $k \geq 0$  and  $\operatorname{Re}(s) > \frac{1}{2}$ , we have

$$2^{s-1} \mathcal{J}_{k,1}(s) = \frac{\gamma}{(k+1)^s} - \frac{s}{(k+1)^{s+1}} - \sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{(n+k)^s}. \quad (8)$$

with

$$\mathcal{J}_{k,1}(s) := \int_{-\infty}^{+\infty} \frac{\zeta(\frac{1}{2} + ix)}{(2k+1+2ix)^s \cosh(\pi x)} dx$$

Applying (8) with  $k=0$  allows us to deduce the following identity:

**Corollary 2.**

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n^s} = \gamma - s - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{\zeta(\frac{1}{2} + ix)}{(\frac{1}{2} + ix)^s \cosh(\pi x)} dx \quad (\operatorname{Re}(s) > \frac{1}{2}). \quad (9)$$

**Example 3.** For  $s=1$ , the representation

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n} = \gamma$$

is regained (since  $\mathcal{J}_{0,1}(1) = -1$ ), and for  $s=2$ , formula (9) gives the relation

$$\sum_{n=2}^{\infty} (-1)^n \frac{\zeta(n)}{n^2} = \gamma - 2 - \frac{1}{2} \int_{-\infty}^{+\infty} \frac{\zeta(\frac{1}{2} + ix)}{(\frac{1}{2} + ix)^2 \cosh(\pi x)} dx. \quad (10)$$

## References

- [1] V. Adamchik, Polygamma functions of negative order, *J. Comput. Appl. Math.* **100** (1998), 191–199.
- [2] L. Bendersky, Sur la fonction gamma généralisée, *Acta Math.* **61** (1933), 263–322.
- [3] I. V. Blagouchine, A complement to a recent paper on some infinite sums with the zeta values, preprint, 2020. Available at <https://arxiv.org/abs/2001.00108>
- [4] B. Candelpergher, *Ramanujan Summation of Divergent Series*, Lecture Notes in Math. 2185, Springer, 2017.

- [5] B. Candelpergher, An expansion of the Riemann Zeta function on the critical line, preprint, 2021. Available at <https://hal.archives-ouvertes.fr/hal-03271709>
- [6] H. Cohen, *Number Theory, Volume II: Analytic and Modern Tools*, Graduate Texts in Math., vol. 240, Springer, 2007.
- [7] M-A. Coppo, A note on some alternating series involving zeta and multiple zeta values, *J. Math. Anal. Appl.* **475** (2019), 1831–1841.
- [8] N. Kurokawa and H. Ochiai, Generalized Kinkelin’s formulas *Kodai Math. J.* **30** (2007), 195–212.
- [9] M. Perkins and R. A. Van Gorder, Closed-form calculation of infinite products of Glaisher-type related to Dirichlet series, *Ramanujan J.* **49** (2019), 371–389.
- [10] J. Sondow and P. Hadjicostas, The generalized-Euler-constant function  $\gamma(z)$  and a generalization of Somos’s quadratic recurrence constant, *J. Math. Anal. Appl.* **332** (2007), 292–314.
- [11] W. Wang, Some asymptotic expansions of hyperfactorial functions and generalized Glaisher-Kinkelin constants, *Ramanujan J.* **43** (2017), 513–533.