



**HAL**  
open science

# A Complement to Laurent expansion of harmonic zeta functions

Marc-Antoine Coppo, Bernard Candelpergher

► **To cite this version:**

Marc-Antoine Coppo, Bernard Candelpergher. A Complement to Laurent expansion of harmonic zeta functions. 2022. hal-03602568v1

**HAL Id: hal-03602568**

**<https://hal.univ-cotedazur.fr/hal-03602568v1>**

Preprint submitted on 9 Mar 2022 (v1), last revised 28 May 2024 (v3)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# A Complement to Laurent expansion of harmonic zeta functions

Marc-Antoine Coppo\* , Bernard Candelpergher

**Abstract** We complement an earlier article dedicated to harmonic zeta functions by outlining a method for obtaining closed-form expressions of the Laurent series coefficients of the harmonic zeta function  $\zeta_H$  about its pole at  $s = 1$ . These coefficients are named harmonic Stieltjes constants by analogy with the classical case.

## 1 Two representations of the harmonic zeta function

We recall that the harmonic zeta function  $\zeta_H$  (noted  $h$  in [2]) is defined by

$$\zeta_H(s) := \sum_{n=1}^{\infty} \frac{H_n}{n^s} \quad \text{for } \operatorname{Re}(s) > 1,$$

where  $H_n$  are the classical harmonic numbers

$$H_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}.$$

We consider the function  $F$  defined by the integral representation

$$F(s) := \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1} e^z}{e^z - 1} \operatorname{Log} \left( \frac{e^z - 1}{z} \right) dz,$$


where  $\operatorname{Log}$  denotes the principal logarithm and  $L$  is the Hankel contour defined by the parametrization

$$z = \operatorname{Log}(1 + e^{ix}) \quad \text{with } x \in ] -\pi, \pi[ \quad \text{for all } z \in L.$$

---

\*Corresponding author.

*E-mail address:* coppo@unice.fr (M-A. Coppo)



This function appears in [2] and plays a central role in the study of the function  $\zeta_H$  around its poles at the negative integers. The following representation:

$$\zeta_H(s) = \pi \cot(\pi s) \zeta(s) + \zeta(s+1) - \psi(1-s)\zeta(s) - \zeta'(s) + F(s) \quad \text{for } s \text{ in } \mathbb{C} \setminus \mathbb{Z}, \quad (1)$$

where  $\psi(s) = \Gamma'(s)/\Gamma(s)$  is the digamma function, is a direct consequence of [2, Theorem 1] (see [2, Eqs. (8), (12) and (25)]).

On the other hand, differentiating the Hankel integral representation of  $\zeta$ :

$$\zeta(s) = \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1}e^z}{1-e^z} dz$$

leads to the following identity:

$$\zeta'(s) + \psi(1-s)\zeta(s) = \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1}e^z \text{Log}(z)}{1-e^z} dz \quad \text{for } s \neq 1, 2, 3, \dots \quad (2)$$

(see [1, Eqs. (2.4)–(2.7)]). Furthermore, the above integral splits into two parts:

$$\int_L \frac{z^{s-1}e^z \text{Log}(z)}{1-e^z} dz = \int_L \frac{z^{s-1}e^z}{e^z-1} \text{Log}\left(\frac{e^z-1}{z}\right) dz - \int_L \frac{z^{s-1}e^z}{e^z-1} \text{Log}(e^z-1) dz,$$

allowing a rewriting of formula (2) as follows:

$$\zeta'(s) + \psi(1-s)\zeta(s) = F(s) - G(s), \quad (3)$$

with

$$G(s) := \frac{\Gamma(1-s)}{2i\pi} \int_L \frac{z^{s-1}e^z}{e^z-1} \text{Log}(e^z-1) dz.$$

Finally, a substitution of (3) in formula (1) leads to another simpler representation of  $\zeta_H$  involving the function  $G$ . We have

$$\zeta_H(s) = \pi \cot(\pi s) \zeta(s) + \zeta(s+1) + G(s) \quad \text{for } s \text{ in } \mathbb{C} \setminus \mathbb{Z}. \quad (4)$$

## 2 Laurent series expansions at $s = 1$

The splitting of  $\zeta_H(s)$  into three parts given by formula (4) above is the key formula for obtaining the Laurent series expansion of the harmonic zeta function about its (double) pole at  $s = 1$ . To show this, we will make use of the following expansions:

- a) The Laurent expansions of  $\pi \cot(\pi s)$  and  $\zeta(s)$  at  $s = 1$  are known. They are respectively

$$\pi \cot(\pi s) = \frac{1}{s-1} - 2\zeta(2)(s-1) - 2\zeta(4)(s-1)^3 - \dots,$$

and

$$\zeta(s) = \frac{1}{s-1} + \gamma - \gamma_1(s-1) + \frac{1}{2}\gamma_2(s-1)^2 + \dots,$$

where  $\gamma_n$  are the classical Stieltjes constants. The expansion of  $\pi \cot(\pi s) \zeta(s)$  is then deduced by Cauchy's product as follows:

$$\pi \cot(\pi s) \zeta(s) = \frac{1}{(s-1)^2} + \frac{\gamma}{s-1} - 2\zeta(2) - \gamma_1 + \left( \frac{1}{2}\gamma_2 - 2\gamma\zeta(2) \right) (s-1) + \dots \quad (5)$$

- b) The Taylor series expansion of  $\zeta(s+1)$  at  $s = 1$  is given by

$$\zeta(s+1) = \zeta(2) + \zeta'(2)(s-1) + \frac{1}{2}\zeta''(2)(s-1)^2 + \dots \quad (6)$$

Moreover, we will use afterwards a well-known expression of  $\zeta'(2)$ :

$$\zeta'(2) = \zeta(2)\gamma + \zeta(2) \log(2\pi) - 2\pi^2 \log(A),$$

where  $A$  is the Glaisher-Kinkelin constant defined by

$$\log A = \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n k \log k - \left( \frac{n^2}{2} + \frac{n}{2} + \frac{1}{12} \right) \log n + \frac{n^2}{4} \right\}.$$

- c) The Laurent expansion of  $\Gamma(z)$  at  $z = 0$  is given by

$$\Gamma(z) = \sum_{k \geq 0} P_k(-\gamma, \dots, (-1)^k \zeta(k)) z^{k-1},$$

where the polynomials  $P_k$  are the modified Bell polynomials defined by

$$\begin{aligned} \exp\left(\sum_{k \geq 1} x_k \frac{z^k}{k}\right) &= \sum_{k \geq 0} P_k(x_1, \dots, x_k) z^k \\ &= 1 + x_1 z + \left(\frac{1}{2}x_2 + \frac{1}{2}x_1^2\right) z^2 + \left(\frac{1}{3}x_3 + \frac{1}{2}x_1 x_2 + \frac{1}{6}x_1^3\right) z^3 + \dots \end{aligned}$$

The Laurent expansion of  $\Gamma(1-s)$  at  $s=1$  is easily deduced from this expression by setting  $z=1-s$ . We have

$$\Gamma(1-s) = -\frac{1}{s-1} - \gamma - \left(\frac{1}{2}\zeta(2) + \frac{1}{2}\gamma^2\right)(s-1) - \left(\frac{1}{3}\zeta(3) + \frac{1}{2}\gamma\zeta(2) + \frac{1}{6}\gamma^3\right)(s-1)^2 + \dots \quad (7)$$

d) Let us consider the function

$$g(s) := G(s)/\Gamma(1-s) = \frac{1}{2i\pi} \int_L \frac{z^{s-1}e^z}{e^z-1} \text{Log}(e^z-1) dz.$$

This is an entire function of  $s$ , and the change of variables

$$z = \text{Log}(1 + e^{ix}) \quad \text{with } x \in ]-\pi, \pi[$$

leads to the integral representation

$$g(s) = -\frac{1}{2i\pi} \int_{-\pi}^{\pi} x (\text{Log}(1 + e^{ix}))^{s-1} dx.$$

If we now define  $K_n$  by

$$K_n := g^{(n)}(1) = \frac{i\pi}{2} \int_{-1}^1 x \text{Log}^n(\text{Log}(1 + e^{i\pi x})) dx,$$

then  $K_0 = g(1) = 0$ , and the Taylor series expansion of  $g(s)$  at  $s=1$  is given by

$$g(s) = \sum_{n \geq 1} \frac{K_n}{n!} (s-1)^n. \quad (8)$$

e) The Laurent expansion of  $G(s)$  at  $s=1$  is deduced from (7) and (8) by Cauchy's product. The result is as follows:

$$G(s) = -K_1 - \left(\frac{K_2}{2} + \gamma K_1\right)(s-1) - \left(\frac{K_3}{6} + \gamma \frac{K_2}{2} + (\gamma^2 + \zeta(2)) \frac{K_1}{2}\right)(s-1)^2 + \dots \quad (9)$$

In particular, it results from the decomposition (4) and the expansions (5), (6), and (9) above that the constant term in the Laurent expansion of  $\zeta_H(s)$  at  $s=1$  is  $-\zeta(2) - \gamma_1 - K_1$ . However, by using yet another representation of  $\zeta_H$ , we have

shown (see [3, Corollary 1 and Lemma 2]) that this constant term is  $\frac{1}{2}\gamma^2 + \frac{1}{2}\zeta(2)$ . By comparing these two expressions, we derive the relation

$$-\zeta(2) - \gamma_1 - K_1 = \frac{1}{2}\gamma^2 + \frac{1}{2}\zeta(2)$$

which gives an expression of the integral  $K_1$  in terms of  $\zeta(2)$ ,  $\gamma$  and  $\gamma_1$ . We have

$$-K_1 = -\frac{i\pi}{2} \int_{-1}^1 x \operatorname{Log}(\operatorname{Log}(1 + e^{i\pi x})) dx = \frac{3}{2}\zeta(2) + \frac{\gamma^2}{2} + \gamma_1 = 2.561174\dots \quad (10)$$

*Remark 1.* Unfortunately, no such formula is known for the integral  $K_2$  whose decimal approximation is  $K_2 = -1.924491$ . More generally, it is not known whether the integrals  $K_n$  for  $n \geq 2$  may be expressed using classical constants.

### 3 Evaluation of the harmonic Stieltjes constants

The expansions (5), (6), and (9) above and the decomposition (4) enable us to write the Laurent series expansion of the harmonic zeta function  $\zeta_H$  about its double pole  $s = 1$ . We have

$$\zeta_H(s) = \frac{1}{(s-1)^2} + \frac{\gamma}{(s-1)} + \tilde{\gamma}_0 - \tilde{\gamma}_1(s-1) + \sum_{n=2}^{\infty} \frac{(-1)^n}{n!} \tilde{\gamma}_n (s-1)^n \quad (0 < |s-1| < 1),$$

where the coefficients  $\tilde{\gamma}_n$  are the *harmonic Stieltjes constants* which are so called by analogy with the classical Stieltjes constants (see [3, Remark 3]). Moreover, this provides a method to evaluate these constants. Thus, for the first two ones, we obtain the following closed-form expressions:

$$\tilde{\gamma}_0 = -\zeta(2) - \gamma_1 - K_1 = \frac{\gamma^2}{2} + \frac{\pi^2}{12} = 0,98905599\dots$$

and

$$\tilde{\gamma}_1 = \frac{1}{2}K_2 + 2\pi^2 \log A - \frac{1}{2}\gamma_2 - \frac{\pi^2}{12}\gamma - \frac{\pi^2}{6} \log 2\pi - \frac{1}{2}\gamma^3 - \gamma\gamma_1 = 0,40076\dots$$

*Remark 2.* The expressions for  $\tilde{\gamma}_0$  and  $\tilde{\gamma}_1$  given above coincide with the asymptotic formula for  $\tilde{\gamma}_n$  mentioned in [3] (see [3, Remark 3 c]). In this order, we have

$$\tilde{\gamma}_0 = \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n \frac{H_k}{k} - \frac{1}{2} \log^2(n) - \gamma \log(n) \right\}$$

and

$$\tilde{\gamma}_1 = \lim_{n \rightarrow \infty} \left\{ \sum_{k=1}^n \frac{H_k}{k} \log k - \frac{1}{3} \log^3(n) - \frac{1}{2} \gamma \log^2(n) \right\}.$$

## References

- [1] K.N. Boyadzhiev, Evaluation of series with Hurwitz and Lerch zeta function coefficients by using Hankel contour integrals, *Applied Mathematics and Computation* 186, (2007).
- [2] K.N. Boyadzhiev, H. Gadiyar and R. Padma, The values of an Euler sum at the negative integers and a relation to a certain convolution of Bernoulli numbers, *Bulletin of the Korean Mathematical Society* 45, (2008).
- [3] B. Candelpergher and M-A. Coppo, Laurent expansion of harmonic zeta functions, *Journal of Mathematical Analysis and Applications* 491, (2020).